

Patient Name: 유돌화

Gender: F

Sample ID: N25-198

Primary Tumor Site: Lung

Collection Date: 2025.08.20

Sample Cancer Type: Lung Cancer

Table of Contents	Page	Report Highlights
Variant Details	2	5 Relevant Biomarkers
Biomarker Descriptions	3	17 Therapies Available
Alert Details	8	211 Clinical Trials
Relevant Therapy Summary	9	

Relevant Lung Cancer Findings

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	EGFR p.(L858R) c.2573T>G, EGFR amplification	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

Genomic Alteration	Finding
Tumor Mutational Burden	4.74 Mut/Mb measured

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	EGFR p.(L858R) c.2573T>G epidermal growth factor receptor Allele Frequency: 78.35% Locus: chr7:55259515 Transcript: NM_005228.5	afatinib <sup>1, 2 / I, II+</sup> amivantamab + lazertinib <sup>1, 2 / I, II+</sup> bevacizumab† + erlotinib <sup>2 / I, II+</sup> dacomitinib <sup>1, 2 / I, II+</sup> erlotinib <sup>2 / I, II+</sup> erlotinib + ramucirumab <sup>1, 2 / I, II+</sup> gefitinib <sup>1, 2 / I, II+</sup> osimertinib <sup>1, 2 / I, II+</sup> osimertinib + chemotherapy <sup>1, 2 / I</sup> amivantamab + chemotherapy <sup>1, 2 / II+</sup> BAT1706 + erlotinib <sup>2</sup> gefitinib + chemotherapy <sup>I</sup> atezolizumab + bevacizumab + chemotherapy <sup>II+</sup>	None*	201

\* Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO  
\* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO  
† Includes biosimilars/generics  
Line of therapy: I: First-line therapy, II+: Other line of therapy  
Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	<i>EGFR amplification</i> epidermal growth factor receptor Locus: chr7:55211010	None*	None*	8
IIC	<i>TP53 p.(L130Pfs*41) c.385_386dup</i> tumor protein p53 Allele Frequency: 20.70% Locus: chr17:7578543 Transcript: NM_000546.6	None*	None*	6
IIC	<i>FLT4 amplification</i> fms related tyrosine kinase 4 Locus: chr5:180030092	None*	None*	3
IIC	<i>FGFR4 amplification</i> fibroblast growth factor receptor 4 Locus: chr5:176517731	None*	None*	2

\* Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO  
\* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO  
† Includes biosimilars/generics  
Line of therapy: I: First-line therapy, II+: Other line of therapy  
Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

 Alerts informed by public data sources:  Contraindicated,  Resistance,  Breakthrough,  Fast Track

*EGFR p.(L858R) c.2573T>G*  *patritumab deruxtecan*<sup>1</sup>

Public data sources included in alerts: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources

*Microsatellite stable, RAD52 p.(S346\*) c.1037C>A, TPMT p.(Y240C) c.719A>G, HLA-A p.(L180\*) c.539T>A, Tumor Mutational Burden*

Variant Details

DNA Sequence Variants							
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
EGFR	p.(L858R)	c.2573T>G	COSM6224	chr7:55259515	78.35%	NM_005228.5	missense
TP53	p.(L130Pfs*41)	c.385_386dup	.	chr17:7578543	20.70%	NM_000546.6	frameshift Insertion
RAD52	p.(S346*)	c.1037C>A	.	chr12:1023218	30.97%	NM_134424.4	nonsense
TPMT	p.(Y240C)	c.719A>G	COSM4986703	chr6:18130918	25.68%	NM_000367.5	missense
HLA-A	p.(L180*)	c.539T>A	.	chr6:29911240	38.58%	NM_001242758.1	nonsense
MECOM	p.(E320Q)	c.958G>C	.	chr3:168840388	2.45%	NM_004991.4	missense
HLA-A	p.(I121R)	c.362_363delTAinsGG	.	chr6:29911063	37.83%	NM_001242758.1	missense
ETS1	p.([L381=;W382R])	c.1143_1144delATinsG C	.	chr11:128333502	3.03%	NM_001143820.2	synonymous, missense

Variant Details (continued)

DNA Sequence Variants (continued)							
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
CREBBP	p.(E1007K)	c.3019G>A	.	chr16:3819216	22.15%	NM_004380.3	missense
RAD51D	p.(C43R)	c.127T>C	.	chr17:33446147	29.86%	NM_133629.3	missense
SMARCA4	p.(E449K)	c.1345G>A	.	chr19:11101925	23.48%	NM_001128849.3	missense

Copy Number Variations			
Gene	Locus	Copy Number	CNV Ratio
EGFR	chr7:55211010	19.26	4.71
FLT4	chr5:180030092	5.67	1.79
FGFR4	chr5:176517731	6.19	1.9
NOTCH2	chr1:120457903	7.56	2.19
TGFBR2	chr3:30648337	5.05	1.65
PXDNL	chr8:52233342	0.53	0.69

Biomarker Descriptions

EGFR amplification, EGFR p.(L858R) c.2573T>G

epidermal growth factor receptor

Background: The EGFR gene encodes the epidermal growth factor receptor (EGFR), a member of the ERBB/human epidermal growth factor receptor (HER) tyrosine kinase family<sup>1</sup>. In addition to EGFR/ERBB1/HER1, other members of the ERBB/HER family include ERBB2/HER2, ERBB3/HER3, and ERBB4/HER4<sup>63</sup>. EGFR ligand-induced dimerization results in kinase activation and leads to stimulation of oncogenic signaling pathways, including the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways<sup>64</sup>. Activation of these pathways promotes cell proliferation, differentiation, and survival<sup>65,66</sup>.

Alterations and prevalence: Recurrent somatic mutations in the tyrosine kinase domain (TKD) of EGFR are observed in approximately 10-20% of lung adenocarcinoma, and at higher frequencies in never-smoker, female, and Asian populations<sup>8,9,67,68</sup>. The most common mutations occur near the ATP-binding pocket of the TKD and include short in-frame deletions in exon 19 (EGFR exon 19 deletion) and the L858R amino acid substitution in exon 21<sup>69</sup>. These mutations constitutively activate EGFR resulting in downstream signaling, and represent 80% of the EGFR mutations observed in lung cancer<sup>69</sup>. A second group of less prevalent activating mutations includes E709K, G719X, S768I, L861Q, and short in-frame insertion mutations in exon 20<sup>70,71,72,73</sup>. EGFR activating mutations in lung cancer tend to be mutually exclusive to KRAS activating mutations<sup>74</sup>. In contrast, a different set of recurrent activating EGFR mutations in the extracellular domain includes R108K, A289V and G598V and are primarily observed in glioblastoma<sup>69,75</sup>. Amplification of EGFR is observed in several cancer types including 44% of glioblastoma multiforme, 12% of esophageal adenocarcinoma, 10% of head and neck squamous cell carcinoma, 8% of brain lower grade glioma, 6% of lung squamous cell carcinoma, 5% of bladder urothelial carcinoma cancer, lung adenocarcinoma, and stomach adenocarcinoma, 3% of cholangiocarcinoma, and 2% of cervical squamous cell carcinoma, sarcoma, and breast invasive carcinoma<sup>8,9,17,68,75</sup>. Deletion of exons 2-7, encoding the extracellular domain of EGFR (EGFRvIII), results in overexpression of a ligand-independent constitutively active protein and is observed in approximately 30% of glioblastoma<sup>76,77,78</sup>. Alterations in EGFR are rare in pediatric cancers<sup>8,9</sup>. Somatic mutations are observed in 2% of bone cancer and glioma, 1% of leukemia (4 in 354 cases), and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 252 cases), peripheral nervous system cancers (1 in 1158 cases), and embryonal tumors (3 in 332 cases)<sup>8,9</sup>. Amplification of EGFR is observed in 2% of bone cancer and less than 1% of Wilms tumor (1 in 136 cases), B-lymphoblastic leukemia/lymphoma (2 in 731 cases), and leukemia (1 in 250 cases)<sup>8,9</sup>.

Potential relevance: Approved first-generation EGFR tyrosine kinase inhibitors (TKIs) include erlotinib<sup>79</sup> (2004) and gefitinib<sup>80</sup> (2015), which block the activation of downstream signaling by reversible interaction with the ATP-binding site. Although initially approved for advanced lung cancer, the discovery that drug sensitivity was associated with exon 19 and exon 21 activating mutations allowed first-generation TKIs to become subsequently approved for front-line therapy in lung cancer tumors containing exon 19 or exon 21

## Biomarker Descriptions (continued)

activating mutations<sup>81</sup>. Second-generation TKIs afatinib<sup>82</sup> (2013) and dacomitinib<sup>83</sup> (2018) bind EGFR and other ERBB/HER gene family members irreversibly and were subsequently approved. First- and second-generation TKIs afatinib, dacomitinib, erlotinib, and gefitinib are recommended for the treatment NSCLC harboring EGFR exon 19 insertions, exon 19 deletions, point mutations L861Q, L858R, S768I, and codon 719 mutations, whereas most EGFR exon 20 insertions, except p.A763\_Y764insFQEA, confer resistance to the same therapies<sup>84,85,86,87</sup>. However, BDTX-189<sup>88</sup> was granted a fast track designation (2020) for the treatment of solid tumors harboring an EGFR exon 20 insertion mutations. In 2022, the FDA granted breakthrough therapy designation to the irreversible EGFR inhibitors, CLN-081 (TPC-064)<sup>89</sup> and sunvozertinib<sup>90</sup>, for locally advanced or metastatic non-small cell lung cancer harboring EGFR exon 20 insertion mutations. In lung cancer containing EGFR exon 19 or 21 activating mutations, treatment with TKIs is eventually associated with the emergence of drug resistance<sup>91</sup>. The primary resistance mutation that emerges following treatment with first-generation TKI is T790M, accounting for 50-60% of resistant cases<sup>69</sup>. Third generation TKIs were developed to maintain sensitivity in the presence of T790M<sup>91</sup>. Osimertinib<sup>92</sup> (2015) is an irreversible inhibitor indicated for metastatic EGFR T790M positive lung cancer and for the first-line treatment of metastatic NSCLC containing EGFR exon 19 deletions or exon 21 L858R mutations. Like first-generation TKIs, treatment with osimertinib is associated with acquired resistance, specifically the C797S mutation, which occurs in 22-44% of cases<sup>91</sup>. The T790M and C797S mutations may be each selected following sequential treatment with a first-generation TKI followed by a third-generation TKI or vice versa<sup>93</sup>. T790M and C797S can occur in either cis or trans allelic orientation<sup>93</sup>. If C797S is observed following progression after treatment with a third-generation TKI in the first-line setting, sensitivity may be retained to first-generation TKIs<sup>93</sup>. If C797S co-occurs in trans with T790M following sequential treatment with first- and third-generation TKIs, patients may exhibit sensitivity to combination first- and third-generation TKIs, but resistance to third-generation TKIs alone<sup>93,94</sup>. However, C797S occurring in cis conformation with T790M, confers resistance to first- and third-generation TKIs<sup>93</sup>. Fourth-generation TKIs are in development to overcome acquired resistance mutations after osimertinib treatment, including BDTX-1535<sup>95</sup> (2024), a CNS-penetrating small molecule inhibitor, that received fast track designation from the FDA for the treatment of patients with EGFR C797S-positive NSCLC who have disease progression on or after a third-generation EGFR TKI. EGFR-targeting antibodies including cetuximab (2004), panitumumab (2006), and necitumumab (2016) are under investigation in combination with EGFR-targeting TKIs for efficacy against EGFR mutations<sup>96</sup>. The bispecific antibody, amivantamab<sup>97</sup> (2021), targeting EGFR and MET was approved for NSCLC tumors harboring EGFR exon 20 insertion mutations. A small molecule kinase inhibitor, lazertinib<sup>98</sup> (2024), was approved in combination with amivantamab as a first-line treatment for adult patients with locally advanced or metastatic NSCLC with EGFR exon 19 deletions or exon 21 L858R mutations. In 2024, a CNS penetrating small molecule, ERAS-801<sup>99</sup> received fast track designation for the treatment of adult patients with EGFR altered glioblastoma. HLX-42<sup>100</sup>, an anti-EGFR-antibody-drug conjugate (ADC) consisting of an anti-EGFR monoclonal antibody conjugated with a novel high potency DNA topoisomerase I (topo I) inhibitor, also received fast track designation (2024) for the treatment of patients with advanced or metastatic EGFR-mutated non-small cell lung cancer whose disease has progressed on a third-generation EGFR tyrosine kinase inhibitor. CPO301<sup>101</sup> (2023) received a fast track designation from the FDA for the treatment of EGFR mutations in patients with metastatic NSCLC who are relapsed/refractory or ineligible for EGFR targeting therapy such as 3rd-generation EGFR inhibitors, including osimertinib. The Oncoprex immunogene therapy quaratusugene ozeplasmid<sup>102</sup> (2020), in combination with osimertinib, received fast track designation from the FDA for NSCLC tumors harboring EGFR mutations that progressed on osimertinib alone. Amplification and mutations of EGFR commonly occur in H3-wild type IDH-wild type diffuse pediatric high-grade glioma<sup>103,104,105</sup>.

### TP53 p.(L130Pfs\*41) c.385\_386dup

#### tumor protein p53

**Background:** The TP53 gene encodes the tumor suppressor protein p53, which binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair<sup>1</sup>. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis<sup>12</sup>. Alterations in TP53 are required for oncogenesis as they result in loss of protein function and gain of transforming potential<sup>13</sup>. Germline mutations in TP53 are the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers<sup>14,15</sup>.

**Alterations and prevalence:** TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)<sup>8,9,16,17,18,19</sup>. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common, including substitutions at codons R158, R175, Y220, R248, R273, and R282<sup>8,9</sup>. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes<sup>20,21,22,23</sup>. Alterations in TP53 are also observed in pediatric cancers<sup>8,9</sup>. Somatic mutations are observed in 53% of non-Hodgkin lymphoma, 24% of soft tissue sarcoma, 19% of glioma, 13% of bone cancer, 9% of B-lymphoblastic leukemia/lymphoma, 4% of embryonal tumors, 3% of Wilms tumor and leukemia, 2% of T-lymphoblastic leukemia/lymphoma, and less than 1% of peripheral nervous system cancers (5 in 1158 cases)<sup>8,9</sup>. Biallelic loss of TP53 is observed in 10% of bone cancer, 2% of Wilms tumor, and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 731 cases) and leukemia (1 in 250 cases)<sup>8,9</sup>.

**Potential relevance:** The small molecule p53 reactivator, PC14586<sup>24</sup> (2020), received a fast track designation by the FDA for advanced tumors harboring a TP53 Y220C mutation. The FDA has granted fast track designation to the p53 reactivator, eprenetapopt<sup>25</sup>, (2019) and breakthrough designation<sup>26</sup> (2020) in combination with azacitidine or azacitidine and venetoclax for acute myeloid leukemia

## Biomarker Descriptions (continued)

patients (AML) and myelodysplastic syndrome (MDS) harboring a TP53 mutation, respectively. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation<sup>27,28</sup>. TP53 mutation are a diagnostic marker of SHH-activated, TP53-mutant medulloblastoma<sup>29</sup>. TP53 mutations confer poor prognosis and poor risk in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), and chronic lymphocytic leukemia (CLL), and acute lymphoblastic leukemia (ALL)<sup>30,31,32,33,34,35</sup>. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant<sup>36</sup>. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system<sup>37</sup>.

### FLT4 amplification

*fms related tyrosine kinase 4*

**Background:** The FLT4 gene encodes the *fms* related receptor tyrosine kinase 4, also known as VEGFR3<sup>1</sup>. FLT4 is a type 2 transmembrane cell surface receptor tyrosine kinase (RTK) and is a member of a family of cognate RTKs called vascular endothelial growth factor receptors (VEGFRs) that also includes VEGFR1 (FLT-1) and VEGFR2 (KDR)<sup>106,107</sup>. Ligand binding to FLT4, including by VEGF-C and VEGF-D, results in FLT4 activation and has been observed to promote tumor metastasis through lymphangiogenesis<sup>106,108</sup>.

**Alterations and prevalence:** Somatic mutations in FLT4 are observed in 12% of skin cutaneous melanoma, 8% of uterine corpus endometrial carcinoma, 6% of colorectal adenocarcinoma and stomach adenocarcinoma, 5% of kidney chromophobe, 4% of lung squamous cell carcinoma and lung adenocarcinoma, and 3% of cervical squamous cell carcinoma, pancreatic adenocarcinoma, adrenocortical carcinoma, and bladder urothelial carcinoma<sup>8,9</sup>. Amplifications are observed in 7% of kidney renal clear cell carcinoma and adrenocortical carcinoma, 4% of uterine carcinosarcoma, and 3% of sarcoma<sup>8,9</sup>.

**Potential relevance:** Currently, no therapies are approved for FLT4 aberrations. Although not approved for specific FLT4 aberrations, the tyrosine kinase inhibitor, sorafenib, is a potent inhibitor of FLT4 and has been approved for the treatment of unresectable hepatocellular carcinoma, advanced renal cell carcinoma, and locally recurrent or metastatic, progressive, differentiated thyroid carcinoma (DTC) refractory to radioactive iodine treatment<sup>109,110</sup>.

### FGFR4 amplification

*fibroblast growth factor receptor 4*

**Background:** The FGFR4 gene encodes fibroblast growth receptor 4, a member of the fibroblast growth-factor receptor (FGFR) family that also includes FGFR1, 2, and 3. These proteins are single-transmembrane receptors composed of three extracellular immunoglobulin (Ig)-type domains and an intracellular kinase domain. Upon FGF-mediated stimulation, FGFRs activate several oncogenic signaling pathways, including the RAS/RAF/MEK/ERK, PI3K/AKT/MTOR, PLC/PKC, and JAK/STAT pathways influencing cell proliferation, migration, and survival<sup>111,112,113</sup>. FGFR4 selectively binds the ligand FGF19, wherein FGF19-mediated aberrant signaling has been identified as an oncogenic driver in hepatocellular carcinoma<sup>114,115</sup>.

**Alterations and prevalence:** Aberrations most common to the FGFR family are amplifications, followed by mutations and fusions. The majority of these aberrations result in gain of function<sup>116</sup>. FGFR4 exhibits amplification in up to 15% of clear-cell renal cell carcinomas, with somatic mutations observed in up to 6% of melanomas and uterine cancer<sup>8,9</sup>.

**Potential relevance:** Currently, no targeted therapies are approved for FGFR4 aberrations. However, FDA-approved multi-kinase inhibitors known to inhibit FGFR family members, including regorafenib (2013), ponatinib (2012), lenvatinib (2015), nintedanib (2014), and pazopanib (2009), have demonstrated anti-tumor activity in select cancer types harboring FGFR alterations<sup>117,118,119,120,121,122,123</sup>. Selective, irreversible FGFR4 inhibitors, including BLU-554, have underwent clinical trial evaluation. In a phase-I clinical study of BLU-554 in patients with FGF19-positive advanced hepatocellular carcinoma, the overall response rate was 17%<sup>124</sup>.

### Microsatellite stable

**Background:** Microsatellites are short tandem repeats (STR) of 1 to 6 bases of DNA between 5 to 50 repeat units in length. There are approximately 0.5 million STRs that occupy 3% of the human genome<sup>38</sup>. Microsatellite instability (MSI) is defined as a change in the length of a microsatellite in a tumor as compared to normal tissue<sup>39,40</sup>. MSI is closely tied to the status of the mismatch repair (MMR) genes. In humans, the core MMR genes include MLH1, MSH2, MSH6, and PMS2<sup>41</sup>. Mutations and loss of expression in MMR genes, known as defective MMR (dMMR), lead to MSI. In contrast, when MMR genes lack alterations, they are referred to as MMR proficient (pMMR). Consensus criteria were first described in 1998 and defined MSI-high (MSI-H) as instability in two or more of the following five markers: BAT25, BAT26, D5S346, D2S123, and D17S250<sup>42</sup>. Tumors with instability in one of the five markers were defined as MSI-low (MSI-L) whereas, those with instability in zero markers were defined as MS-stable (MSS)<sup>42</sup>. Tumors classified as MSI-L are often phenotypically indistinguishable from MSS tumors and tend to be grouped with MSS<sup>43,44,45,46,47</sup>. MSI-H is a hallmark of Lynch syndrome (LS), also known as hereditary non-polyposis colorectal cancer, which is caused by germline mutations in the MMR genes<sup>40</sup>.

## Biomarker Descriptions (continued)

LS is associated with an increased risk of developing colorectal cancer, as well as other cancers, including endometrial and stomach cancer<sup>39,40,44,48</sup>.

**Alterations and prevalence:** The MSI-H phenotype is observed in 30% of uterine corpus endothelial carcinoma, 20% of stomach adenocarcinoma, 15-20% of colon adenocarcinoma, and 5-10% of rectal adenocarcinoma<sup>39,40,49,50</sup>. MSI-H is also observed in 5% of adrenal cortical carcinoma and at lower frequencies in other cancers such as esophageal, liver, and ovarian cancers<sup>49,50</sup>.

**Potential relevance:** Anti-PD-1 immune checkpoint inhibitors including pembrolizumab<sup>51</sup> (2014) and nivolumab<sup>52</sup> (2015) are approved for patients with MSI-H or dMMR colorectal cancer who have progressed following chemotherapy. Pembrolizumab<sup>51</sup> is also approved as a single agent, for the treatment of patients with advanced endometrial carcinoma that is MSI-H or dMMR with disease progression on prior therapy who are not candidates for surgery or radiation. Importantly, pembrolizumab is approved for the treatment of MSI-H or dMMR solid tumors that have progressed following treatment, with no alternative option and is the first anti-PD-1 inhibitor to be approved with a tumor agnostic indication<sup>51</sup>. Dostarlimab<sup>53</sup> (2021) is also approved for dMMR recurrent or advanced endometrial carcinoma or solid tumors that have progressed on prior treatment and is recommended as a subsequent therapy option in dMMR/MSI-H advanced or metastatic colon or rectal cancer<sup>45,54</sup>. The cytotoxic T-lymphocyte antigen 4 (CTLA-4) blocking antibody, ipilimumab<sup>55</sup> (2011), is approved alone or in combination with nivolumab in MSI-H or dMMR colorectal cancer that has progressed following treatment with chemotherapy. MSI-H may confer a favorable prognosis in colorectal cancer although outcomes vary depending on stage and tumor location<sup>45,56,57</sup>. Specifically, MSI-H is a strong prognostic indicator of better overall survival (OS) and relapse free survival (RFS) in stage II as compared to stage III colorectal cancer patients<sup>57</sup>. The majority of patients with tumors classified as either MSS or pMMR do not benefit from treatment with single-agent immune checkpoint inhibitors as compared to those with MSI-H tumors<sup>58,59</sup>. However, checkpoint blockade with the addition of chemotherapy or targeted therapies have demonstrated response in MSS or pMMR cancers<sup>58,59</sup>.

### **RAD52 p.(S346\*) c.1037C>A**

*RAD52 homolog, DNA repair protein*

**Background:** The RAD52 gene encodes the RAD52 homolog, DNA repair protein<sup>1</sup>. RAD52 binds to single- and double-stranded DNA and enables strand exchange for double-strand break (DSB) repair by binding to RAD51<sup>10</sup>. RAD52 also promotes DSB repair through homologous recombination repair (HRR) by recruiting BRCA1 to sites of DSBs, which leads to the removal of TP53BP1 and prevents DSB repair by non-homologous end joining (NHEJ)<sup>11</sup>.

**Alterations and prevalence:** Somatic mutations in RAD52 are observed in 2% of uterine corpus endometrial carcinoma, uterine carcinosarcoma, and skin cutaneous melanoma<sup>8,9</sup>.

**Potential relevance:** Currently, no therapies are approved for RAD52 aberrations.

### **TPMT p.(Y240C) c.719A>G**

*thiopurine S-methyltransferase*

**Background:** The TPMT gene encodes thiopurine S-methyltransferase, a cytosolic enzyme that methylates aromatic and heterocyclic sulfhydryl compounds such as thiopurines<sup>1,60,61</sup>. TPMT is the major enzyme responsible for the metabolic inactivation of thiopurine chemotherapeutic drugs used in the treatment of acute lymphoblastic leukemia (ALL), including, 6-mercaptopurine, 6-thioguanine, and azathioprine<sup>60,61,62</sup>. Inherited TPMT polymorphisms, including TPMT\*2, TPMT\*3A, TPMT\*3B, TPMT\*3C, and TPMT\*8, can result in TPMT deficiency, which is characterized by impaired enzymatic activity and confers an increased risk of severe toxicity to thiopurine drugs due to an increase in systemic drug exposure<sup>60,62</sup>.

**Alterations and prevalence:** Somatic mutations in TPMT are observed in 2% of uterine corpus endometrial carcinoma and colorectal adenocarcinoma<sup>8,9</sup>. Biallelic loss of TPMT is observed in 1% of stomach adenocarcinoma, esophageal adenocarcinoma, and adrenocortical carcinoma<sup>8,9</sup>. Amplification of TPMT is observed in 7% of ovarian serous cystadenocarcinoma, 6% of bladder urothelial carcinoma, 4% of diffuse large B-cell lymphoma, uveal melanoma, uterine carcinosarcoma, and skin cutaneous melanoma, 3% of cholangiocarcinoma, and 2% of breast invasive carcinoma, uterine corpus endometrial carcinoma, and liver hepatocellular carcinoma<sup>8,9</sup>.

**Potential relevance:** Currently, no therapies are approved for TPMT aberrations.

### **HLA-A p.(L180\*) c.539T>A**

*major histocompatibility complex, class I, A*

**Background:** The HLA-A gene encodes the major histocompatibility complex, class I, A<sup>1</sup>. MHC (major histocompatibility complex) class I molecules are located on the cell surface of nucleated cells and present antigens from within the cell for recognition by cytotoxic T

## Biomarker Descriptions (continued)

cells<sup>2</sup>. MHC class I molecules are heterodimers composed of two polypeptide chains,  $\alpha$  and B2M<sup>3</sup>. The classical MHC class I genes include HLA-A, HLA-B, and HLA-C and encode the  $\alpha$  polypeptide chains, which present short polypeptide chains, of 7 to 11 amino acids, to the immune system to distinguish self from non-self<sup>4,5,6</sup>. Downregulation of MHC class I promotes tumor evasion of the immune system, suggesting a tumor suppressor role for HLA-A<sup>7</sup>.

Alterations and prevalence: Somatic mutations in HLA-A are observed in 7% of diffuse large B-cell lymphoma (DLBCL), 4% of cervical squamous cell carcinoma and head and neck squamous cell carcinoma, 3% of colorectal adenocarcinoma, and 2% of uterine corpus endometrial carcinoma and stomach adenocarcinoma<sup>8,9</sup>. Biallelic loss of HLA-A is observed in 4% of DLBCL<sup>8,9</sup>.

Potential relevance: Currently, no therapies are approved for HLA-A aberrations.

Alerts Informed By Public Data Sources

Current FDA Information

Contraindicated   Not recommended   Resistance   Breakthrough   Fast Track

FDA information is current as of 2025-05-14. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

EGFR p.(L858R) c.2573T>G

patritumab deruxtecan

Cancer type: Non-Small Cell Lung Cancer   Variant class: EGFR L858R mutation or EGFRi sensitizing mutation

Supporting Statement:  
The FDA has granted Breakthrough Therapy designation to a potential first-in-class HER3 directed antibody-drug conjugate, patritumab deruxtecan, for metastatic or locally advanced, EGFR-mutant non-small cell lung cancer.

Reference:  
<https://www.cancernetwork.com/view/fda-grants-breakthrough-therapy-status-to-patritumab-deruxtecan-for-egfr-metastatic-nsccl>

Genes Assayed

Genes Assayed for the Detection of DNA Sequence Variants

ABL1, ABL2, ACVR1, AKT1, AKT2, AKT3, ALK, AR, ARAF, ATP1A1, AURKA, AURKB, AURKC, AXL, BCL2, BCL2L12, BCL6, BCR, BMP5, BRAF, BTK, CACNA1D, CARD11, CBL, CCND1, CCND2, CCND3, CCNE1, CD79B, CDK4, CDK6, CHD4, CSF1R, CTNNB1, CUL1, CYSLTR2, DDR2, DGCR8, DROSHA, E2F1, EGFR, EIF1AX, EPAS1, ERBB2, ERBB3, ERBB4, ESR1, EZH2, FAM135B, FGF7, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FOXL2, FOXO1, GATA2, GLI1, GNA11, GNAQ, GNAS, HIF1A, HRAS, IDH1, IDH2, IKBKB, IL6ST, IL7R, IRF4, IRS4, KCNJ5, KDR, KIT, KLF4, KLF5, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAPK1, MAX, MDM4, MECOM, MED12, MEF2B, MET, MITF, MPL, MTOR, MYC, MYCN, MYD88, MYOD1, NFE2L2, NRAS, NSD2, NT5C2, NTRK1, NTRK2, NTRK3, NUP93, PAX5, PCBP1, PDGFRA, PDGFRB, PIK3C2B, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PIK3R2, PIM1, PLCG1, PPP2R1A, PPP6C, PRKACA, PTPN11, PTPRD, PXDNL, RAC1, RAF1, RARA, RET, RGS7, RHEB, RHOA, RICTOR, RIT1, ROS1, RPL10, SETBP1, SF3B1, SIX1, SIX2, SLC01B3, SMC1A, SMO, SNCAIP, SOS1, SOX2, SPOP, SRC, SRSF2, STAT3, STAT5B, STAT6, TAF1, TERT, TGFBF1, TOP1, TOP2A, TPMT, TRRAP, TSHR, U2AF1, USP8, WAS, XPO1, ZNF217, ZNF429

Genes Assayed for the Detection of Copy Number Variations

ABCB1, ABL1, ABL2, ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AKT1, AKT2, AKT3, ALK, AMER1, APC, AR, ARAF, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AURKA, AURKC, AXIN1, AXIN2, AXL, B2M, BAP1, BARD1, BCL2, BCL2L12, BCL6, BCOR, BLM, BMPR2, BRAF, BRCA1, BRCA2, BRIP1, CARD11, CASP8, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDK4, CDK6, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHD4, CHEK1, CHEK2, CIC, CREBBP, CSMD3, CTCF, CTLA4, CTNND2, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, DAXX, DDR1, DDR2, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, EGFR, EIF1AX, ELF3, EMSY, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERBB2, ERBB3, ERBB4, ERCC2, ERCC4, ERFFI1, ESR1, ETV6, EZH2, FAM135B, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAT1, FBXW7, FGF19, FGF23, FGF3, FGF4, FGF9, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FUBP1, FYN, GATA2, GATA3, GLI3, GNA13, GNAS, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, IDH2, IGF1R, IKBKB, IL7R, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KDR, KEAP1, KIT, KLF5, KMT2A, KMT2B, KMT2C, KMT2D, KRAS, LARP4B, LATS1, LATS2, MAGOH, MAP2K1, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK1, MAPK8, MAX, MCL1, MDM2, MDM4, MECOM, MEF2B, MEN1, MET, MGA, MITF, MLH1, MLH3, MPL, MRE11, MSH2, MSH3, MSH6, MTAP, MTOR, MUTYH, MYC, MYCL, MYCN, MYD88, NBN, NCOR1, NF1, NF2, NFE2L2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, NRAS, NTRK1, NTRK3, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PCBP1, PDCD1, PDCD1LG2, PDGFRA, PDGFRB, PDIA3, PGD, PHF6, PIK3C2B, PIK3CA, PIK3CB, PIK3R1, PIK3R2, PIM1, PLCG1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R1A, PPP2R2A, PPP6C, PRDM1, PRDM9, PRKACA, PRKAR1A, PTCH1, PTEN, PTPN11, PTPRT, PXDNL, RAC1, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RAF1, RARA, RASA1, RASA2, RB1, RBM10, RECQL4, RET, RHEB, RICTOR, RIT1, RNASEH2A, RNASEH2B, RNF43, ROS1, RPA1, RPS6KB1, RPTOR, RUNX1, SDHA, SDHB, SDHD, SETBP1, SETD2, SF3B1, SLC01B3, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SMC1A, SMO, SOX9, SPEN, SPOP, SRC, STAG2, STAT3, STAT6, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TERT, TET2, TGFBF2,

Genes Assayed (continued)

Genes Assayed for the Detection of Copy Number Variations (continued)

TNFAIP3, TNFRSF14, TOP1, TP53, TP63, TPMT, TPP2, TSC1, TSC2, U2AF1, USP8, USP9X, VHL, WT1, XPO1, XRCC2, XRCC3, YAP1, YES1, ZFH3, ZMYM3, ZNF217, ZNF429, ZRSR2

Genes Assayed for the Detection of Fusions

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSP02, RSP03, TERT

Genes Assayed with Full Exon Coverage

ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AMER1, APC, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AXIN1, AXIN2, B2M, BAP1, BARD1, BCOR, BLM, BMPR2, BRCA1, BRCA2, BRIP1, CALR, CASP8, CBF3, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHEK1, CHEK2, CIC, CIITA, CREBBP, CSMD3, CTCF, CTLA4, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, CYP2D6, DAXX, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, ELF3, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERCC2, ERCC4, ERCC5, ERFF1, ETV6, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAS, FAT1, FBXW7, FUBP1, GATA3, GNA13, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, ID3, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KEAP1, KLHL13, KMT2A, KMT2B, KMT2C, KMT2D, LARP4B, LATS1, LATS2, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK8, MEN1, MGA, MLH1, MLH3, MRE11, MSH2, MSH3, MSH6, MTAP, MTUS2, MUTYH, NBN, NCOR1, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PDCD1, PDCD1LG2, PDIA3, PGD, PHF6, PIK3R1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R2A, PRDM1, PRDM9, PRKAR1A, PSMB10, PSMB8, PSMB9, PTCH1, PTEN, PTPRT, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RASA1, RASA2, RB1, RBM10, RECQL4, RNASEH2A, RNASEH2B, RNASEH2C, RNF43, RPA1, RPL22, RPL5, RUNX1, RUNX1T1, SDHA, SDHB, SDHC, SDHD, SETD2, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SOCS1, SOX9, SPEN, STAG2, STAT1, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TET2, TGFB2, TMEM132D, TNFAIP3, TNFRSF14, TP53, TP63, TPP2, TSC1, TSC2, UGT1A1, USP9X, VHL, WT1, XRCC2, XRCC3, ZBTB20, ZFH3, ZMYM3, ZRSR2

Relevant Therapy Summary

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

EGFR p.(L858R) c.2573T>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib					(III)
afatinib					(II)
dacomitinib					(II)
gefitinib					(II)
erlotinib + ramucirumab					
amivantamab + carboplatin + pemetrexed					
amivantamab + lazertinib					
osimertinib + chemotherapy + pemetrexed					
bevacizumab + erlotinib					
erlotinib					

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib + carboplatin + pemetrexed	✕	●	✕	✕	✕
osimertinib + cisplatin + pemetrexed	✕	●	✕	✕	✕
BAT1706 + erlotinib	✕	✕	●	✕	✕
bevacizumab (Allergan) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Biocon) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Celltrion) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Mabxience) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Pfizer) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Samsung Bioepis) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Stada) + erlotinib	✕	✕	●	✕	✕
atezolizumab + bevacizumab + carboplatin + paclitaxel	✕	✕	✕	●	✕
gefitinib + carboplatin + pemetrexed	✕	✕	✕	●	✕
adebreliumab, bevacizumab, chemotherapy	✕	✕	✕	✕	● (IV)
afatinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (IV)
befotertinib	✕	✕	✕	✕	● (IV)
bevacizumab, almonertinib, chemotherapy	✕	✕	✕	✕	● (IV)
catequentinib, toripalimab	✕	✕	✕	✕	● (IV)
EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (IV)
gefitinib, chemotherapy	✕	✕	✕	✕	● (IV)
gefitinib, endostatin	✕	✕	✕	✕	● (IV)
natural product, gefitinib, erlotinib, icotinib hydrochloride, osimertinib, almonertinib, furmonertinib	✕	✕	✕	✕	● (IV)
almonertinib, apatinib	✕	✕	✕	✕	● (III)
almonertinib, catequentinib	✕	✕	✕	✕	● (III)
almonertinib, chemotherapy	✕	✕	✕	✕	● (III)
almonertinib, radiation therapy	✕	✕	✕	✕	● (III)
almonertinib, radiation therapy, chemotherapy	✕	✕	✕	✕	● (III)
befotertinib, icotinib hydrochloride	✕	✕	✕	✕	● (III)
bevacizumab, osimertinib	✕	✕	✕	✕	● (III)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
BL-B01D1	✕	✕	✕	✕	● (III)
BL-B01D1, osimertinib	✕	✕	✕	✕	● (III)
CK-101, gefitinib	✕	✕	✕	✕	● (III)
datopotamab deruxtecan, osimertinib	✕	✕	✕	✕	● (III)
FHND9041, afatinib	✕	✕	✕	✕	● (III)
furmonertinib	✕	✕	✕	✕	● (III)
furmonertinib, osimertinib, chemotherapy	✕	✕	✕	✕	● (III)
gefitinib, afatinib, erlotinib, metformin hydrochloride	✕	✕	✕	✕	● (III)
icotinib hydrochloride, catequentinib	✕	✕	✕	✕	● (III)
icotinib hydrochloride, chemotherapy	✕	✕	✕	✕	● (III)
icotinib hydrochloride, radiation therapy	✕	✕	✕	✕	● (III)
JMT-101, osimertinib	✕	✕	✕	✕	● (III)
osimertinib, bevacizumab	✕	✕	✕	✕	● (III)
osimertinib, chemotherapy	✕	✕	✕	✕	● (III)
osimertinib, datopotamab deruxtecan	✕	✕	✕	✕	● (III)
sacituzumab tirumotecan	✕	✕	✕	✕	● (III)
sacituzumab tirumotecan, osimertinib	✕	✕	✕	✕	● (III)
savolitinib, osimertinib	✕	✕	✕	✕	● (III)
SH-1028	✕	✕	✕	✕	● (III)
targeted therapy	✕	✕	✕	✕	● (III)
TY-9591, osimertinib	✕	✕	✕	✕	● (III)
SCTB-14, chemotherapy	✕	✕	✕	✕	● (II/III)
ABSK-043, furmonertinib	✕	✕	✕	✕	● (II)
almonertinib	✕	✕	✕	✕	● (II)
almonertinib, adebrelimab, chemotherapy	✕	✕	✕	✕	● (II)
almonertinib, bevacizumab	✕	✕	✕	✕	● (II)
almonertinib, chemoradiation therapy	✕	✕	✕	✕	● (II)
almonertinib, dacomitinib	✕	✕	✕	✕	● (II)
amivantamab, chemotherapy	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

☒ In this cancer type    
 ☐ In other cancer type    
 ☒ In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
amivantamab, lazertinib, chemotherapy	✕	✕	✕	✕	● (II)
atezolizumab, bevacizumab, tiragolumab	✕	✕	✕	✕	● (II)
befotertinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
bevacizumab, afatinib	✕	✕	✕	✕	● (II)
bevacizumab, furmonertinib	✕	✕	✕	✕	● (II)
cadonilimab, chemotherapy, catequentinib	✕	✕	✕	✕	● (II)
camrelizumab, apatinib	✕	✕	✕	✕	● (II)
capmatinib, osimertinib, ramucirumab	✕	✕	✕	✕	● (II)
catequentinib, almonertinib	✕	✕	✕	✕	● (II)
chemotherapy, atezolizumab, bevacizumab	✕	✕	✕	✕	● (II)
dacomitinib, osimertinib	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor, osimertinib, chemotherapy	✕	✕	✕	✕	● (II)
EGFR tyrosine kinase inhibitor, radiation therapy	✕	✕	✕	✕	● (II)
erlotinib, chemotherapy	✕	✕	✕	✕	● (II)
erlotinib, OBI-833	✕	✕	✕	✕	● (II)
furmonertinib, bevacizumab	✕	✕	✕	✕	● (II)
furmonertinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
furmonertinib, catequentinib	✕	✕	✕	✕	● (II)
furmonertinib, chemotherapy	✕	✕	✕	✕	● (II)
furmonertinib, chemotherapy, bevacizumab	✕	✕	✕	✕	● (II)
furmonertinib, icotinib hydrochloride	✕	✕	✕	✕	● (II)
gefitinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
gefitinib, icotinib hydrochloride	✕	✕	✕	✕	● (II)
gefitinib, thalidomide	✕	✕	✕	✕	● (II)
icotinib hydrochloride	✕	✕	✕	✕	● (II)
icotinib hydrochloride, autologous RAK cell	✕	✕	✕	✕	● (II)
icotinib hydrochloride, osimertinib	✕	✕	✕	✕	● (II)
ivonescimab, chemotherapy	✕	✕	✕	✕	● (II)
lazertinib	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
lazertinib, bevacizumab	✕	✕	✕	✕	● (II)
lazertinib, chemotherapy	✕	✕	✕	✕	● (II)
lenvatinib, pembrolizumab	✕	✕	✕	✕	● (II)
osimertinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
osimertinib, chemoradiation therapy	✕	✕	✕	✕	● (II)
osimertinib, radiation therapy	✕	✕	✕	✕	● (II)
PLB-1004, bozitinib, osimertinib	✕	✕	✕	✕	● (II)
ramucirumab, erlotinib	✕	✕	✕	✕	● (II)
sacituzumab govitecan	✕	✕	✕	✕	● (II)
sacituzumab tirumotecan, chemotherapy, osimertinib	✕	✕	✕	✕	● (II)
sunvozertinib	✕	✕	✕	✕	● (II)
sunvozertinib, catequentinib	✕	✕	✕	✕	● (II)
sunvozertinib, golidocitinib	✕	✕	✕	✕	● (II)
tislelizumab, chemotherapy, bevacizumab	✕	✕	✕	✕	● (II)
toripalimab	✕	✕	✕	✕	● (II)
toripalimab, bevacizumab, Clostridium butyricum, chemotherapy	✕	✕	✕	✕	● (II)
toripalimab, chemotherapy	✕	✕	✕	✕	● (II)
TY-9591, chemotherapy	✕	✕	✕	✕	● (II)
zorifertinib, pirotinib	✕	✕	✕	✕	● (II)
AFM-24_I, atezolizumab	✕	✕	✕	✕	● (I/II)
almonertinib, icotinib hydrochloride	✕	✕	✕	✕	● (I/II)
BDTX-1535	✕	✕	✕	✕	● (I/II)
benmelstobart, catequentinib	✕	✕	✕	✕	● (I/II)
BH-30643	✕	✕	✕	✕	● (I/II)
bozitinib, osimertinib	✕	✕	✕	✕	● (I/II)
BPI-361175	✕	✕	✕	✕	● (I/II)
cetrelimab, amivantamab	✕	✕	✕	✕	● (I/II)
dacomitinib, catequentinib	✕	✕	✕	✕	● (I/II)
DAJH-1050766	✕	✕	✕	✕	● (I/II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
DB-1310, osimertinib	✕	✕	✕	✕	● (I/II)
dositinib	✕	✕	✕	✕	● (I/II)
FWD-1509	✕	✕	✕	✕	● (I/II)
H-002	✕	✕	✕	✕	● (I/II)
ifebemtiniib, furmonertinib	✕	✕	✕	✕	● (I/II)
MRTX0902	✕	✕	✕	✕	● (I/II)
necitumumab, osimertinib	✕	✕	✕	✕	● (I/II)
quaratusugene ozeplasmid, osimertinib	✕	✕	✕	✕	● (I/II)
RC-108, furmonertinib, toripalimab	✕	✕	✕	✕	● (I/II)
sotiburafusp alfa, HB-0030	✕	✕	✕	✕	● (I/II)
sunvozertinib, chemotherapy	✕	✕	✕	✕	● (I/II)
TAS-3351	✕	✕	✕	✕	● (I/II)
TQ-B3525, osimertinib	✕	✕	✕	✕	● (I/II)
TRX-221	✕	✕	✕	✕	● (I/II)
WSD-0922	✕	✕	✕	✕	● (I/II)
afatinib, chemotherapy	✕	✕	✕	✕	● (I)
almonertinib, midazolam	✕	✕	✕	✕	● (I)
ASKC-202	✕	✕	✕	✕	● (I)
AZD-9592	✕	✕	✕	✕	● (I)
BG-60366	✕	✕	✕	✕	● (I)
BPI-1178, osimertinib	✕	✕	✕	✕	● (I)
catequentinib, gefitinib, metformin hydrochloride	✕	✕	✕	✕	● (I)
DZD-6008	✕	✕	✕	✕	● (I)
EGFR tyrosine kinase inhibitor, catequentinib	✕	✕	✕	✕	● (I)
genolimzumab, fruquintinib	✕	✕	✕	✕	● (I)
IBI-318, lenvatinib	✕	✕	✕	✕	● (I)
KQB-198, osimertinib	✕	✕	✕	✕	● (I)
LAVA-1223	✕	✕	✕	✕	● (I)
MRX-2843, osimertinib	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ① In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib, carotuximab	✕	✕	✕	✕	● (I)
osimertinib, Minnelide	✕	✕	✕	✕	● (I)
osimertinib, tegatrabetan	✕	✕	✕	✕	● (I)
patritumab deruxtecan	✕	✕	✕	✕	● (I)
repotrectinib, osimertinib	✕	✕	✕	✕	● (I)
VIC-1911, osimertinib	✕	✕	✕	✕	● (I)
WJ13404	✕	✕	✕	✕	● (I)
WTS-004	✕	✕	✕	✕	● (I)
YH-013	✕	✕	✕	✕	● (I)
YL-202	✕	✕	✕	✕	● (I)

### EGFR amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
nimotuzumab	✕	✕	✕	✕	● (III)
ensartinib, sintilimab	✕	✕	✕	✕	● (II)
sacituzumab govitecan	✕	✕	✕	✕	● (II)
BBI-355, erlotinib	✕	✕	✕	✕	● (I/II)
DF-9001	✕	✕	✕	✕	● (I/II)
GB263T	✕	✕	✕	✕	● (I/II)
MCLA-129	✕	✕	✕	✕	● (I/II)
EGFR targeted DK210 diakine (Deka BioSciences), chemotherapy, radiation therapy, pembrolizumab, nivolumab	✕	✕	✕	✕	● (I)

### TP53 p.(L130Pfs\*41) c.385\_386dup

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
almonertinib, catequentinib	✕	✕	✕	✕	● (III)
osimertinib, chemotherapy	✕	✕	✕	✕	● (III)
osimertinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
sunvozertinib, catequentinib	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

FLT4 amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
pazopanib	×	×	×	×	● (II)
regorafenib	×	×	×	×	● (II)
sunitinib, regorafenib	×	×	×	×	● (II)

FGFR4 amplification

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
BBI-355, futibatinib	×	×	×	×	● (I/II)
ABSK-121	×	×	×	×	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

HRR Details

Gene/Genomic Alteration	Finding
LOH percentage	14.8%
BRCA2	LOH, 13q13.1(32890491-32972932)x2

Homologous recombination repair (HRR) genes were defined from published evidence in relevant therapies, clinical guidelines, as well as clinical trials, and include - BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, and RAD54L.

Thermo Fisher Scientific's Ion Torrent Oncomine Reporter software was used in generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on Oncomine Reporter (6.1.1 data version 2025.06(006)). The data presented here are from a curated knowledge base of publicly available information, but may not be exhaustive. FDA information was sourced from [www.fda.gov](http://www.fda.gov) and is current as of 2025-05-14. NCCN information was sourced from [www.nccn.org](http://www.nccn.org) and is current as of 2025-05-01. EMA information was sourced from [www.ema.europa.eu](http://www.ema.europa.eu) and is current as of 2025-05-14. ESMO information was sourced from [www.esmo.org](http://www.esmo.org) and is current as of 2025-05-01. Clinical Trials information is current as of 2025-05-01. For the most up-to-date information regarding a particular trial, search [www.clinicaltrials.gov](http://www.clinicaltrials.gov) by NCT ID or search local clinical trials authority website by local identifier listed in 'Other identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials listed in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

## References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Hulpke et al. The MHC I loading complex: a multitasking machinery in adaptive immunity. *Trends Biochem Sci.* PMID: 23849087
3. Adams et al. The adaptable major histocompatibility complex (MHC) fold: structure and function of nonclassical and MHC class I-like molecules. *Annu Rev Immunol.* 2013;31:529-61. PMID: 23298204
4. Rossjohn et al. T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol.* 2015;33:169-200. PMID: 25493333
5. Parham. MHC class I molecules and KIRs in human history, health and survival. *Nat Rev Immunol.* 2005 Mar;5(3):201-14. PMID: 15719024
6. Sidney et al. HLA class I supertypes: a revised and updated classification. *BMC Immunol.* 2008 Jan 22;9:1. PMID: 18211710
7. Cornel et al. MHC Class I Downregulation in Cancer: Underlying Mechanisms and Potential Targets for Cancer Immunotherapy. *Cancers (Basel).* 2020 Jul 2;12(7). PMID: 32630675
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
10. Jalan et al. Emerging Roles of RAD52 in Genome Maintenance. *Cancers (Basel).* 2019 Jul 23;11(7). PMID: 31340507
11. Yasuhara et al. Human Rad52 Promotes XPG-Mediated R-loop Processing to Initiate Transcription-Associated Homologous Recombination Repair. *Cell.* 2018 Oct 4;175(2):558-570.e11. PMID: 30245011
12. Nag et al. The MDM2-p53 pathway revisited. *J Biomed Res.* 2013 Jul;27(4):254-71. PMID: 23885265
13. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell.* 2014 Mar 17;25(3):304-17. PMID: 24651012
14. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol.* 2010 Jan;2(1):a001008. PMID: 20182602
15. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med.* 2017 Apr 3;7(4). PMID: 28270529
16. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature.* 2012 Sep 27;489(7417):519-25. PMID: 22960745
17. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature.* 2015 Jan 29;517(7536):576-82. PMID: 25631445
18. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet.* 2016 Jun;48(6):607-16. PMID: 27158780
19. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature.* 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061
20. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat.* 2002 Jun;19(6):607-14. PMID: 12007217
21. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer.* 2011 Apr;2(4):466-74. PMID: 21779514
22. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene.* 2007 Apr 2;26(15):2157-65. PMID: 17401424
23. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat.* 2014 Jun;35(6):766-78. PMID: 24729566
24. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
25. <https://ir.aprea.com/news-releases/news-release-details/aprea-therapeutics-receives-fda-fast-track-designation>
26. <http://vp280.alertir.com/en/pressreleases/karolinska-development%27s-portfolio-company-aprea-therapeutics-receives-fda-breakthrough-therapy-designation-1769167>
27. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol.* 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534
28. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci.* 2017 Nov;74(22):4171-4187. PMID: 28643165

## References (continued)

29. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol.* 2021 Aug 2;23(8):1231-1251. PMID: 34185076
30. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]
31. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
32. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 2.2025]
33. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 1.2025]
34. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 3.2025]
35. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 3.2024]
36. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 2.2025]
37. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med.* 2020 Aug 3. PMID: 32747829
38. Lander et al. Initial sequencing and analysis of the human genome. *Nature.* 2001 Feb 15;409(6822):860-921. PMID: 11237011
39. Baudrin et al. Molecular and Computational Methods for the Detection of Microsatellite Instability in Cancer. *Front Oncol.* 2018 Dec 12;8:621. doi: 10.3389/fonc.2018.00621. eCollection 2018. PMID: 30631754
40. Nojadeh et al. Microsatellite instability in colorectal cancer. *EXCLI J.* 2018;17:159-168. PMID: 29743854
41. Saeed et al. Microsatellites in Pursuit of Microbial Genome Evolution. *Front Microbiol.* 2016 Jan 5;6:1462. doi: 10.3389/fmicb.2015.01462. eCollection 2015. PMID: 26779133
42. Boland et al. A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res.* 1998 Nov 15;58(22):5248-57. PMID: 9823339
43. Halford et al. Low-level microsatellite instability occurs in most colorectal cancers and is a nonrandomly distributed quantitative trait. *Cancer Res.* 2002 Jan 1;62(1):53-7. PMID: 11782358
44. Imai et al. Carcinogenesis and microsatellite instability: the interrelationship between genetics and epigenetics. *Carcinogenesis.* 2008 Apr;29(4):673-80. PMID: 17942460
45. NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2025]
46. Pawlik et al. Colorectal carcinogenesis: MSI-H versus MSI-L. *Dis. Markers.* 2004;20(4-5):199-206. PMID: 15528785
47. Lee et al. Low-Level Microsatellite Instability as a Potential Prognostic Factor in Sporadic Colorectal Cancer. *Medicine (Baltimore).* 2015 Dec;94(50):e2260. PMID: 26683947
48. Latham et al. Microsatellite Instability Is Associated With the Presence of Lynch Syndrome Pan-Cancer. *J. Clin. Oncol.* 2019 Feb 1;37(4):286-295. PMID: 30376427
49. Cortes-Ciriano et al. A molecular portrait of microsatellite instability across multiple cancers. *Nat Commun.* 2017 Jun 6;8:15180. doi: 10.1038/ncomms15180. PMID: 28585546
50. Bonneville et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precis Oncol.* 2017;2017. PMID: 29850653
51. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125514s174lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125514s174lbl.pdf)
52. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125554s129lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125554s129lbl.pdf)
53. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/761174s009lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/761174s009lbl.pdf)
54. NCCN Guidelines® - NCCN-Rectal Cancer [Version 2.2025]
55. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125377s133lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125377s133lbl.pdf)
56. Ribic et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N. Engl. J. Med.* 2003 Jul 17;349(3):247-57. PMID: 12867608
57. Klingbiel et al. Prognosis of stage II and III colon cancer treated with adjuvant 5-fluorouracil or FOLFIRI in relation to microsatellite status: results of the PETACC-3 trial. *Ann. Oncol.* 2015 Jan;26(1):126-32. PMID: 25361982
58. Hermel et al. The Emerging Role of Checkpoint Inhibition in Microsatellite Stable Colorectal Cancer. *J Pers Med.* 2019 Jan 16;9(1). PMID: 30654522
59. Ciardiello et al. Immunotherapy of colorectal cancer: Challenges for therapeutic efficacy. *Cancer Treat. Rev.* 2019 Jun;76:22-32. PMID: 31079031
60. Katara et al. TPMT Polymorphism: When Shield Becomes Weakness. *Interdiscip Sci.* 2016 Jun;8(2):150-155. PMID: 26297310
61. Yong et al. The role of pharmacogenetics in cancer therapeutics. *Br J Clin Pharmacol.* 2006 Jul;62(1):35-46. PMID: 16842377

## References (continued)

62. McLeod et al. Genetic polymorphism of thiopurine methyltransferase and its clinical relevance for childhood acute lymphoblastic leukemia. *Leukemia*. 2000 Apr;14(4):567-72. PMID: 10764140
63. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science*. 1985 Sep 6;229(4717):974-6. PMID: 2992089
64. Liu et al. EGFR-TKIs resistance via EGFR-independent signaling pathways. *Mol Cancer*. 2018 Feb 19;17(1):53. PMID: 29455669
65. Zhixiang. ErbB Receptors and Cancer. *Methods Mol. Biol.* 2017;1652:3-35. PMID: 28791631
66. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
67. Pines et al. Oncogenic mutant forms of EGFR: lessons in signal transduction and targets for cancer therapy. *FEBS Lett.* 2010 Jun 18;584(12):2699-706. PMID: 20388509
68. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature*. 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552
69. da et al. EGFR mutations and lung cancer. *Annu Rev Pathol.* 2011;6:49-69. doi: 10.1146/annurev-pathol-011110-130206. PMID: 20887192
70. Arcila et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol. Cancer Ther.* 2013 Feb;12(2):220-9. PMID: 23371856
71. Kobayashi et al. EGFR Exon 18 Mutations in Lung Cancer: Molecular Predictors of Augmented Sensitivity to Afatinib or Neratinib as Compared with First- or Third-Generation TKIs. *Clin Cancer Res.* 2015 Dec 1;21(23):5305-13. doi: 10.1158/1078-0432.CCR-15-1046. Epub 2015 Jul 23. PMID: 26206867
72. Yasuda et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med.* 2013 Dec 18;5(216):216ra177. PMID: 24353160
73. Chiu et al. Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Treatment Response in Advanced Lung Adenocarcinomas with G719X/L861Q/S768I Mutations. *J Thorac Oncol.* 2015 May;10(5):793-9. PMID: 25668120
74. Karachaliou et al. KRAS mutations in lung cancer. *Clin Lung Cancer.* 2013 May;14(3):205-14. PMID: 23122493
75. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
76. Mitsudomi et al. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. *FEBS J.* 2010 Jan;277(2):301-8. PMID: 19922469
77. Gazdar. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene.* 2009 Aug;28 Suppl 1:S24-31. PMID: 19680293
78. Gan et al. The EGFRvIII variant in glioblastoma multiforme. *J Clin Neurosci.* 2009 Jun;16(6):748-54. PMID: 19324552
79. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/021743s025lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf)
80. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/206995s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/206995s004lbl.pdf)
81. Riely et al. Clinical course of patients with non-small cell lung cancer and epidermal growth factor receptor exon 19 and exon 21 mutations treated with gefitinib or erlotinib. *Clin Cancer Res.* 2006 Feb 1;12(3 Pt 1):839-44. PMID: 16467097
82. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/201292s017lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/201292s017lbl.pdf)
83. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/211288s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/211288s003lbl.pdf)
84. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 3.2025]
85. Naidoo et al. Epidermal growth factor receptor exon 20 insertions in advanced lung adenocarcinomas: Clinical outcomes and response to erlotinib. *Cancer.* 2015 Sep 15;121(18):3212-3220. PMID: 26096453
86. Vyse et al. Targeting EGFR exon 20 insertion mutations in non-small cell lung cancer. *Signal Transduct Target Ther.* 2019;4:5. PMID: 30854234
87. Yi et al. A comparison of epidermal growth factor receptor mutation testing methods in different tissue types in non-small cell lung cancer. *Int J Mol Med.* 2014 Aug;34(2):464-74. PMID: 24891042
88. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-granted-fast-track-designation-fda>
89. <https://investors.cullinanoncology.com/news-releases/news-release-details/fda-grants-breakthrough-therapy-designation-cullinan-oncologys>
90. <https://www.prnewswire.com/news-releases/fda-grants-breakthrough-therapy-designation-for-dizal-pharmaceuticals-dzd9008-in-patients-with-locally-advanced-or-metastatic-non-small-cell-lung-cancer-harboring-egfr-exon20-insertion-301469692.html>
91. Madic et al. EGFR C797S, EGFR T790M and EGFR sensitizing mutations in non-small cell lung cancer revealed by six-color crystal digital PCR. *Oncotarget.* 2018 Dec 21;9(100):37393-37406. PMID: 30647840

## References (continued)

92. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/208065s033lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/208065s033lbl.pdf)
93. Niederst et al. The Allelic Context of the C797S Mutation Acquired upon Treatment with Third-Generation EGFR Inhibitors Impacts Sensitivity to Subsequent Treatment Strategies. *Clin. Cancer Res.* 2015 Sep 1;21(17):3924-33. PMID: 25964297
94. Wang et al. Lung Adenocarcinoma Harboring EGFR T790M and In Trans C797S Responds to Combination Therapy of First- and Third-Generation EGFR TKIs and Shifts Allelic Configuration at Resistance. *J Thorac Oncol.* 2017 Nov;12(11):1723-1727. PMID: 28662863
95. <https://investors.blackdiamondtherapeutics.com//news-releases/news-release-details/black-diamond-therapeutics-announces-corporate-update-and>
96. Ciardiello et al. The role of anti-EGFR therapies in EGFR-TKI-resistant advanced non-small cell lung cancer. *Cancer Treat Rev.* 2024 Jan;122:102664. PMID: 38064878
97. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/761210s007lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/761210s007lbl.pdf)
98. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/219008s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/219008s000lbl.pdf)
99. <https://investors.erasca.com//news-releases/news-release-details/erasca-granted-fda-fast-track-designation-cns-penetrant-egfr>
100. <https://iis.aastocks.com/20231227/11015917-0.PDF>
101. <http://iis.aastocks.com/20230612/10770455-0.PDF>
102. <https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>
103. NCCN Guidelines® - NCCN-Pediatric Central Nervous System Cancers [Version 2.2025]
104. Buccoliero et al. Pediatric High Grade Glioma Classification Criteria and Molecular Features of a Case Series. *Genes (Basel).* 2022 Mar 31;13(4). PMID: 35456430
105. Louis et al. cIMPACT-NOW update 6: new entity and diagnostic principle recommendations of the cIMPACT-Utrecht meeting on future CNS tumor classification and grading. *Brain Pathol.* 2020 Jul;30(4):844-856. PMID: 32307792
106. Holmes et al. The vascular endothelial growth factor (VEGF) family: angiogenic factors in health and disease. *Genome Biol.* 2005;6(2):209. PMID: 15693956
107. Fontanella et al. Clinical advances in the development of novel VEGFR2 inhibitors. *Ann Transl Med.* 2014 Dec;2(12):123. PMID: 25568876
108. Su et al. The role of the VEGF-C/VEGFR-3 axis in cancer progression. *Br J Cancer.* 2007 Feb 26;96(4):541-5. PMID: 17164762
109. Wilhelm et al. BAY 43-9006 exhibits broad spectrum oral antitumor activity and targets the RAF/MEK/ERK pathway and receptor tyrosine kinases involved in tumor progression and angiogenesis. *Cancer Res.* 2004 Oct 1;64(19):7099-109. PMID: 15466206
110. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/021923s024lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/021923s024lbl.pdf)
111. Babina et al. Advances and challenges in targeting FGFR signalling in cancer. *Nat. Rev. Cancer.* 2017 May;17(5):318-332. PMID: 28303906
112. Ahmad et al. Mechanisms of FGFR-mediated carcinogenesis. *Biochim. Biophys. Acta.* 2012 Apr;1823(4):850-60. PMID: 22273505
113. Sarabipour et al. Mechanism of FGF receptor dimerization and activation. *Nat Commun.* 2016 Jan 4;7:10262. doi: 10.1038/ncomms10262. PMID: 26725515
114. Repana et al. Targeting FGF19/FGFR4 Pathway: A Novel Therapeutic Strategy for Hepatocellular Carcinoma. *Diseases.* 2015 Oct 28;3(4):294-305. PMID: 28943626
115. Lu et al. Fibroblast Growth Factor Receptor 4 (FGFR4) Selective Inhibitors as Hepatocellular Carcinoma Therapy: Advances and Prospects. *J. Med. Chem.* 2018 Nov 16. PMID: 30403487
116. Helsten et al. The FGFR Landscape in Cancer: Analysis of 4,853 Tumors by Next-Generation Sequencing. *Clin. Cancer Res.* 2016 Jan 1;22(1):259-67. PMID: 26373574
117. Cha et al. FGFR2 amplification is predictive of sensitivity to regorafenib in gastric and colorectal cancers in vitro. *Mol Oncol.* 2018 Jun;12(7):993-1003. PMID: 29573334
118. Chae et al. Inhibition of the fibroblast growth factor receptor (FGFR) pathway: the current landscape and barriers to clinical application. *Oncotarget.* 2017 Feb 28;8(9):16052-16074. PMID: 28030802
119. Porta et al. FGFR a promising druggable target in cancer: Molecular biology and new drugs. *Crit. Rev. Oncol. Hematol.* 2017 May;113:256-267. PMID: 28427515
120. Gozgit et al. Ponatinib (AP24534), a multitargeted pan-FGFR inhibitor with activity in multiple FGFR-amplified or mutated cancer models. *Mol. Cancer Ther.* 2012 Mar;11(3):690-9. PMID: 22238366
121. Yamamoto et al. Lenvatinib, an angiogenesis inhibitor targeting VEGFR/FGFR, shows broad antitumor activity in human tumor xenograft models associated with microvessel density and pericyte coverage. *Vasc Cell.* 2014 Sep 6;6:18. doi: 10.1186/2045-824X-6-18. eCollection 2014. PMID: 25197551

## References (continued)

122. Kim et al. Pazopanib, a novel multitargeted kinase inhibitor, shows potent in vitro antitumor activity in gastric cancer cell lines with FGFR2 amplification. *Mol. Cancer Ther.* 2014 Nov;13(11):2527-36. PMID: 25249557
123. Hibi et al. FGFR gene alterations in lung squamous cell carcinoma are potential targets for the multikinase inhibitor nintedanib. *Cancer Sci.* 2016 Nov;107(11):1667-1676. PMID: 27581340
124. Kim et al. First-in-Human Phase I Study of Fisogatinib (BLU-554) Validates Aberrant FGF19 Signaling as a Driver Event in Hepatocellular Carcinoma. *Cancer Discov.* 2019 Dec;9(12):1696-1707. PMID: 31575541