

Patient Name: 한한희
Gender: M
Sample ID: N25-232

Primary Tumor Site:
Collection Date: 2025.09.10

Sample Cancer Type: Lung Cancer

Table of Contents	Page	Report Highlights
Variant Details	2	3 Relevant Biomarkers
Biomarker Descriptions	3	1 Therapies Available
Alert Details	9	25 Clinical Trials
Relevant Therapy Summary	14	

Relevant Lung Cancer Findings

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	None detected	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	KRAS p.(G12V) c.35G>T	ROS1	None detected
MET	None detected		

Genomic Alteration	Finding
Tumor Mutational Burden	0.95 Mut/Mb measured

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	KRAS p.(G12V) c.35G>T KRAS proto-oncogene, GTPase Allele Frequency: 14.55% Locus: chr12:25398284 Transcript: NM_033360.4	None*	bevacizumab + chemotherapy [†]	21
IIC	CDKN2A deletion cyclin dependent kinase inhibitor 2A Locus: chr9:21968178	None*	None*	3
IIC	ATRX deletion ATRX, chromatin remodeler Locus: chrX:76763769	None*	None*	1

* Public data sources included in relevant therapies: FDA¹, NCCN, EMA², ESMO

* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Line of therapy: I: First-line therapy, II+: Other line of therapy

Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

Prevalent cancer biomarkers without relevant evidence based on included data sources
CUL4B deletion, UGT1A1 p.(G71R) c.211G>A, NQO1 p.(P187S) c.559C>T, ZRSR2 deletion, BCOR deletion, USP9X deletion, DDX3X deletion, KDM6A deletion, RBM10 deletion, KDM5C deletion, SMC1A deletion, AMER1 deletion, ZMYM3 deletion, STAG2 deletion, PHF6 deletion, Tumor Mutational Burden

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
KRAS	p.(G12V)	c.35G>T	COSM520	chr12:25398284	14.55%	NM_033360.4	missense
UGT1A1	p.(G71R)	c.211G>A	COSM4415616	chr2:234669144	48.80%	NM_000463.3	missense
NQO1	p.(P187S)	c.559C>T	.	chr16:69745145	48.67%	NM_000903.3	missense
SUGP1	p.(N289S)	c.866A>G	.	chr19:19413095	48.41%	NM_172231.4	missense

Copy Number Variations

Gene	Locus	Copy Number	CNV Ratio
CDKN2A	chr9:21968178	0.48	0.7
ATRX	chrX:76763769	0	0.53
CUL4B	chrX:119660593	0	0.58
ZRSR2	chrX:15808582	0	0.57
BCOR	chrX:39911340	0	0.57
USP9X	chrX:40982869	0	0.54
DDX3X	chrX:41193501	0	0.52
KDM6A	chrX:44732715	0	0.55
RBM10	chrX:47006798	0	0.54
KDM5C	chrX:53221892	0	0.56
SMC1A	chrX:53406966	0	0.53
AMER1	chrX:63409727	0	0.58
ZMYM3	chrX:70460753	0	0.53
STAG2	chrX:123156472	0	0.55
PHF6	chrX:133511628	0	0.52
EIF1AX	chrX:20148599	0	0.56
ARAF	chrX:47422311	0	0.55
AR	chrX:66766015	0	0.56

Biomarker Descriptions

KRAS p.(G12V) c.35G>T

KRAS proto-oncogene, GTPase

Background: The KRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes NRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival^{7,8,9}.

Alterations and prevalence: Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. KRAS mutations are observed in up to 10-20% of uterine cancer, 30-35% of lung adenocarcinoma and colorectal cancer, and about 60% of pancreatic cancer⁵. The majority of KRAS mutations consist of point mutations occurring at G12, G13, and Q61^{5,10,11}. Mutations at A59, K117, and A146 have also been observed but are less frequent^{6,12}.

Potential relevance: The FDA has approved the small molecule inhibitors, sotorasib¹³ (2021) and adagrasib¹⁴ (2022), for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC). Sotorasib and adagrasib are also useful in certain circumstances for KRAS G12C-mutated pancreatic adenocarcinoma¹⁵. The FDA has also granted breakthrough therapy designation (2022) to the KRAS G12C inhibitor, GDC-6036¹⁶, for KRAS G12C-mutated non-small cell lung cancer. The SHP2 inhibitor, BBP-398¹⁷ was granted fast track designation (2022) in combination with sotorasib for previously treated patients with KRAS G12C-mutated metastatic NSCLC. The RAF/MEK clamp, avutemetinib¹⁸ was also granted fast track designation (2024) in combination with sotorasib for KRAS G12C-mutated metastatic NSCLC who have received at least one prior systemic therapy and have not been previously treated with a KRAS G12C inhibitor. The KRAS G12C inhibitor, BBO-8520¹⁹, was granted fast track designation in 2025 for previously treated KRAS G12C-mutated patients with metastatic NSCLC. The KRAS G12C inhibitor, D3S-001²⁰, was granted fast track designation in 2024 for KRAS G12C-mutated patients with advanced unresectable or metastatic colorectal cancers. The PLK1 inhibitor, onvansertib²¹, was granted fast track designation (2020) in combination with bevacizumab and FOLFIRI for second-line treatment of patients with KRAS-mutated metastatic colorectal cancer (mCRC). The EGFR antagonists, cetuximab²² and panitumumab²³, are contraindicated for treatment of colorectal cancer patients with KRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)¹². Additionally, KRAS mutations are associated with poor prognosis in NSCLC²⁴.

CDKN2A deletion

cyclin dependent kinase inhibitor 2A

Background: CDKN2A encodes cyclin dependent kinase inhibitor 2A, a cell cycle regulator that controls G1/S progression¹. CDKN2A, also known as p16/INK4A, belongs to a family of INK4 cyclin-dependent kinase inhibitors, which also includes CDKN2B (p15/INK4B), CDKN2C (p18/INK4C), and CDKN2D (p19/INK4D)⁸⁶. The INK4 family regulates cell cycle progression by inhibiting CDK4 or CDK6, thereby preventing the phosphorylation of Rb^{87,88,89}. CDKN2A encodes two alternative transcript variants, namely p16 and p14ARF, both of which exhibit differential tumor suppressor functions⁹⁰. Specifically, the CDKN2A/p16 transcript inhibits cell cycle kinases CDK4 and CDK6, whereas the CDKN2A/p14ARF transcript stabilizes the tumor suppressor protein p53 to prevent its degradation^{1,90,91}. CDKN2A aberrations commonly co-occur with CDKN2B⁸⁶. Loss of CDKN2A/p16 results in downstream inactivation of the Rb and p53 pathways, leading to uncontrolled cell proliferation⁹². Germline mutations of CDKN2A are known to confer a predisposition to melanoma and pancreatic cancer^{93,94}.

Alterations and prevalence: Somatic alterations in CDKN2A often result in loss of function (LOF) which is attributed to copy number loss, truncating, or missense mutations⁹⁵. Somatic mutations in CDKN2A are observed in 20% of head and neck squamous cell carcinoma and pancreatic adenocarcinoma, 15% of lung squamous cell carcinoma, 13% of skin cutaneous melanoma, 8% of esophageal adenocarcinoma, 7% of bladder urothelial carcinoma, 6% of cholangiocarcinoma, 4% of lung adenocarcinoma and stomach adenocarcinoma, and 2% of liver hepatocellular carcinoma, uterine carcinosarcoma, and cervical squamous cell carcinoma^{5,6}. Biallelic deletion of CDKN2A is observed in 56% of glioblastoma multiforme, 45% of mesothelioma, 39% of esophageal adenocarcinoma, 32% of bladder urothelial carcinoma, 31% of skin cutaneous melanoma and head and neck squamous cell carcinoma, 28% of pancreatic adenocarcinoma, 27% of diffuse large B-cell lymphoma, 26% of lung squamous cell carcinoma, 17% of lung adenocarcinoma and cholangiocarcinoma, 15% of sarcoma, 11% of stomach adenocarcinoma and of brain lower grade glioma, 7% of adrenocortical carcinoma, 6% of liver hepatocellular carcinoma, 4% of breast invasive carcinoma, kidney renal papillary cell carcinoma and thymoma, 3% of ovarian serous cystadenocarcinoma and kidney renal clear cell carcinoma, and 2% of uterine carcinosarcoma and kidney chromophobe^{5,6}. Alterations in CDKN2A are also observed in pediatric cancers⁶. Biallelic deletion of CDKN2A is observed in 68% of T-lymphoblastic leukemia/lymphoma, 40% of B-lymphoblastic leukemia/lymphoma, 25% of glioma, 19% of bone cancer, and 6% of embryonal tumors⁶. Somatic mutations in CDKN2A are observed in less than 1.5% of bone cancer (5 in 327 cases), B-lymphoblastic leukemia/lymphoma (3 in 252 cases), and leukemia (1 in 354 cases)⁶.

Potential relevance: Loss of CDKN2A can be useful in the diagnosis of mesothelioma, and mutations in CDKN2A are ancillary diagnostic markers of malignant peripheral nerve sheath tumors^{38,96,97}. Additionally, deletion of CDKN2B is a molecular marker used in staging Grade 4 pediatric IDH-mutant astrocytoma⁹⁸. Currently, no therapies are approved for CDKN2A aberrations. However, CDKN2A

Biomarker Descriptions (continued)

LOF leading to CDK4/6 activation may confer sensitivity to CDK inhibitors such as palbociclib and abemaciclib^{99,100,101}. Alternatively, CDKN2A expression and Rb inactivation demonstrate resistance to palbociclib in cases of glioblastoma multiforme¹⁰². CDKN2A (p16) expression is associated with a favorable prognosis for progression-free survival (PFS) and overall survival (OS) in p16/HPV positive head and neck cancer^{103,104,105,106}.

ATRX deletion

ATRX, chromatin remodeler

Background: The ATRX gene encodes the ATRX chromatin remodeler and ATPase/helicase domain protein, which belongs to SWI/SNF family of chromatin remodeling proteins¹. The SWI/SNF proteins are a group of DNA translocases that use ATP hydrolysis to remodel chromatin structure and maintain genomic integrity by controlling transcriptional regulation, DNA repair, and chromosome stability through the regulation of telomere length^{52,53,54,55}. ATRX is a tumor suppressor that interacts with the MRE11-RAD50-NBN (MRN) complex, which is involved in double-stranded DNA (dsDNA) break repair^{56,57,58}.

Alterations and prevalence: Somatic mutations of ATRX are observed in 38% of brain lower grade glioma, 15% of uterine corpus endometrial carcinoma, 14% of sarcoma, 9% of glioblastoma multiforme and skin cutaneous melanoma, 7% of colorectal adenocarcinoma, 6% of lung adenocarcinoma, stomach adenocarcinoma, and cervical squamous cell carcinoma, 5% of bladder urothelial carcinoma and lung squamous cell carcinoma, 4% of adrenocortical carcinoma, head and neck squamous cell carcinoma and uterine carcinosarcoma, and 2% of diffuse large B-cell lymphoma, ovarian serous cystadenocarcinoma, breast invasive carcinoma, pheochromocytoma and paraganglioma, kidney renal clear cell carcinoma, pancreatic adenocarcinoma, liver hepatocellular carcinoma and kidney chromophobe^{5,6}. Biallelic deletion of ATRX is observed in 7% of sarcoma, 3% of kidney chromophobe, and 2% of brain lower grade glioma^{5,6}. Although alterations of ATRX in pediatric populations are rare, somatic mutations are observed in 6% of gliomas, 4% of bone cancer, 3% of soft tissue sarcoma, and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 252 cases), embryonal tumor (3 in 332 cases), and leukemia (2 in 354 cases)⁶. Biallelic deletion of ATRX is observed in 1% of peripheral nervous system tumors (1 in 91 cases) in and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 731 cases)⁶.

Potential relevance: Currently, no therapies are approved for ATRX aberrations. Loss of ATRX protein expression correlates with the presence of ATRX mutations^{59,60}. ATRX deficiency along with IDH mutation and TP53 mutation is diagnostic of astrocytoma IDH-mutant as defined by the World Health Organization (WHO)^{61,62}.

CUL4B deletion

cullin 4B

Background: The CUL4B gene encodes cullin 4B, a member of the cullin family, which includes CUL1, CUL2, CUL3, CUL4a, CUL5, CUL7, and Parc^{1,2}. CUL4B belongs to the CUL4 subfamily which also includes CUL4A³. CUL4A and CUL4B share greater than 80% sequence identity and functional redundancy^{3,4}. Cullin proteins share a conserved cullin homology domain and act as molecular scaffolds for RING E3 ubiquitin ligases to assemble into cullin-RING ligase complexes (CRLs)². CUL4B is part of the CRL4 complex which is responsible for ubiquitination and degradation of a variety of substrates where substrate specificity is dependent on the substrate recognition component of the CRL4 complex⁴. CRL4 substrates include oncoproteins, tumor suppressors, nucleotide excision repair proteins, cell cycle promoters, histone methylation proteins, and tumor-related signaling molecules, thereby impacting various processes critical to tumor development and progression and supporting a complex role of CUL4B in oncogenesis^{3,4}.

Alterations and prevalence: Somatic mutations in CUL4B are observed in 9% of uterine corpus endometrial carcinoma, 5% of skin cutaneous melanoma, and 2% of bladder urothelial carcinoma, cervical squamous cell carcinoma, colorectal adenocarcinoma, uterine carcinosarcoma, brain lower grade glioma, and lung squamous cell carcinoma^{5,6}. Amplification of CUL4B is observed in 2% of diffuse large B-cell lymphoma^{5,6}. Biallelic loss of CUL4B is observed in 1% sarcoma and testicular germ cell tumors^{5,6}.

Potential relevance: Currently, no therapies are approved for CUL4B aberrations.

UGT1A1 p.(G71R) c.211G>A

UDP glucuronosyltransferase family 1 member A1

Background: The UGT1A1 gene encodes UDP glucuronosyltransferase family 1 member A1, a member of the UDP-glucuronosyltransferase 1A (UGT1A) subfamily of the UGT protein superfamily^{1,113}. UGTs are microsomal membrane-bound enzymes that catalyze the glucuronidation of endogenous and xenobiotic compounds and transform the lipophilic molecules into excretable, hydrophilic metabolites^{113,114}. UGTs play an important role in drug metabolism, detoxification, and metabolite homeostasis. Differential expression of UGTs can promote cancer development, disease progression, as well as drug resistance¹¹⁵. Specifically, elevated expression of UGT1As are associated with resistance to many anti-cancer drugs due to drug inactivation and lower active drug concentrations. However, reduced expression and downregulation of UGT1As are implicated in bladder and hepatocellular tumorigenesis and progression due to toxin accumulation^{115,116,117,118}. Furthermore, UGT1A1 polymorphisms, such as UGT1A1*28,

Biomarker Descriptions (continued)

UGT1A1*93, and UGT1A1*6, confer an increased risk of severe toxicity to irinotecan-based chemotherapy treatment of solid tumors, due to reduced glucuronidation of the irinotecan metabolite, SN-38¹¹⁹.

Alterations and prevalence: Biallelic deletion of UGT1A1 has been observed in 6% of sarcoma, 3% of brain lower grade glioma and uveal melanoma, and 2% of thymoma, cervical squamous cell carcinoma, bladder urothelial carcinoma, head and neck squamous cell carcinoma, and esophageal adenocarcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for UGT1A1 aberrations.

ZRSR2 deletion

zinc finger CCCH-type, RNA binding motif and serine/arginine rich 2

Background: The ZRSR2 gene encodes the zinc finger CCCH-type, RNA binding motif and serine/arginine-rich 2 protein, a component of the spliceosome. Specifically, ZRSR2 encodes a splicing factor that is involved in the recognition of the 3' intron splice site⁷¹. ZRSR2 interacts with components of the pre-spliceosome assembly including SRSF2 and U2AF2/U2AF1 heterodimer^{71,72}. Mutations in ZRSR2 can lead to deregulated global and alternative mRNA splicing, nuclear-cytoplasm export, and unspliced mRNA degradation while concurrently altering the expression of multiple genes^{71,73}.

Alterations and prevalence: ZRSR2 alterations including nonsense and frameshift mutations are observed in 5-10% of myelodysplastic syndromes (MDS) and 4% of uterine cancer. ZRSR2 deletions are observed in 4% of diffuse large B-cell lymphoma (DLBCL), 3% of head and neck and esophageal cancers^{6,30}.

Potential relevance: Mutation of ZRSR2 is associated with poor prognosis in myelodysplastic syndromes as well as poor/adverse risk in acute myeloid leukemia (AML)^{30,40,41}.

BCOR deletion

BCL6 corepressor

Background: The BCOR gene encodes the B-cell CLL/lymphoma 6 (BCL6) co-repressor protein, which potentiates transcriptional repression by BCL6^{25,26}. BCOR also associates with class I and II histone deacetylases (HDACs), suggesting an alternate mechanism for BCOR-mediated transcriptional repression independent of BCL6²⁶. Genetic alterations in BCOR result in protein dysfunction, which suggests BCOR functions as a tumor suppressor gene^{27,28,29}.

Alterations and prevalence: Genetic alterations in BCOR include missense, nonsense, and frameshift mutations that result in loss of function and have been observed in up to 5% of myelodysplastic syndromes (MDS), 5-10% of chronic myelomonocytic leukemia (CMML), and 1-5% of acute myeloid leukemia (AML)^{5,30,31,32}. Higher mutational frequencies are reported in some solid tumors, including up to 15% of uterine cancer and 5-10% of colorectal cancer, stomach cancer, cholangiocarcinoma, and melanoma^{5,6}. Although less common, BCOR fusions and internal tandem duplications (ITDs) have been reported in certain rare cancer types^{33,34,35}. Specifically, BCOR::CCNB3 rearrangements define a particular subset of sarcomas with Ewing sarcoma-like morphology known as BCOR::CCNB3 sarcomas (BCS)^{36,37}. Alterations in BCOR are also observed in pediatric cancers^{5,6}. Somatic mutations are observed in 13% of soft tissue sarcoma, 4% of glioma, 3% of retinoblastoma, 2% of bone cancer, 1% of B-lymphoblastic leukemia/lymphoma (3 in 252 cases), and less than 1% of embryonal tumors (3 in 332 cases), leukemia (2 in 311 cases), and Wilms tumor (2 in 710 cases)^{5,6}. Other alterations have been reported in clear cell carcinoma of the kidney, a rare pediatric renal malignant tumor, with one study reporting the presence of BCOR ITDs in more than 90% of cases³³.

Potential relevance: BCOR rearrangement, including inv(X)(p11.4p11.22) resulting in BCOR::CCNB3 fusion, is diagnostic of sarcoma with BCOR genetic alterations, a subset of undifferentiated round cell sarcomas^{38,39}. Additionally, translocation t(x;22)(p11;q13) resulting in ZC3H7B::BCOR fusion is a useful ancillary diagnostic marker of high-grade endometrial stromal sarcoma³⁸. Somatic mutation in BCOR is one of the possible molecular abnormality requirements for the diagnosis of myelodysplasia-related AML (AML-MR) and is associated with poor prognosis in AML and MDS^{30,31,40,41,42}. In FLT3-ITD negative AML patients under 65 with intermediate cytogenetic prognosis, mutations in BCOR confer inferior overall survival (OS) as well as relapse-free survival (RFS) compared to those without BCOR abnormalities (OS = 13.6% vs. 55%; RFS = 14.3% vs. 44.5%)³². Additionally, BCOR ITDs and BCOR::EP300 fusion are molecular alterations of significance in pediatric gliomas^{43,44}.

USP9X deletion

ubiquitin specific peptidase 9 X-linked

Background: The USP9X gene encodes the ubiquitin specific peptidase 9 X-linked protein¹. USP9X is a deubiquitinating enzyme (DUB) and a member of the ubiquitin-specific protease (USP) subclass of cysteine proteases⁶³. DUBs are responsible for protein deubiquitination, thereby counter-regulating post-transcriptional ubiquitin modification of proteins within the cell^{63,64}. USP9X has many

Biomarker Descriptions (continued)

substrates and is commonly upregulated in several solid tumor types, supporting an oncogenic role for USP9X⁶⁴. Conversely, in some cancer types, USP9X has been observed to function as a tumor suppressor, suggesting its exact role in cancer may be dependent on its substrates⁶⁴. In breast cancer, USP9X has been shown to stabilize BRCA1 by inhibiting its ubiquitination, thereby influencing the regulation of homologous recombination and repair⁶⁴.

Alterations and prevalence: Somatic mutations are observed in 16% of uterine corpus endometrial carcinoma, 11% of skin cutaneous melanoma, 7% of colorectal adenocarcinoma, 6% of cholangiocarcinoma, 5% of stomach adenocarcinoma, lung squamous cell carcinoma, diffuse large B-cell lymphoma (DLBCL), and head and neck squamous cell carcinoma^{5,6}. Biallelic deletions are observed in 4% of esophageal adenocarcinoma, 3% of head and neck squamous cell carcinoma, 2% of mesothelioma, uterine carcinosarcoma, and lung squamous cell carcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for USP9X aberrations.

DDX3X deletion

DEAD-box helicase 3, X-linked

Background: The DDX3X gene encodes DEAD-box helicase 3 X-linked, a member of the DEAD-box protein family, which is part of the RNA helicase superfamily II^{1,120}. DEAD-box helicases contain twelve conserved motifs including a "DEAD" domain which is characterized by a conserved amino acid sequence of Asp-Glu-Ala-Asp (DEAD)^{120,121,122,123}. In DEAD-box proteins, the DEAD domain interacts with β - and γ -phosphates of ATP through Mg²⁺ and is required for ATP hydrolysis¹²⁰. DDX3X is involved in several processes including the unwinding of double-stranded RNA, splicing of pre-mRNA, RNA export, transcription, and translation^{124,125,126,127,128,129,130,131}. Deregulation of DDX3X has been shown to impact cancer progression by modulating proliferation, metastasis, and drug resistance¹²⁴.

Alterations and prevalence: Somatic mutations in DDX3X are observed in 9% of skin cutaneous melanoma and uterine corpus endometrial carcinoma, 7% of diffuse large B-cell lymphoma, 4% of cervical squamous cell carcinoma, bladder urothelial carcinoma, and stomach adenocarcinoma, and 2% of lung squamous cell carcinoma and head and neck squamous cell carcinoma^{5,6}. Biallelic loss of DDX3X is observed in 4% of esophageal adenocarcinoma, 3% of head and neck squamous cell carcinoma, and 2% of mesothelioma and lung squamous cell carcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for DDX3X aberrations.

KDM6A deletion

lysine demethylase 6A

Background: The KDM6A gene encodes the lysine demethylase 6A protein¹. KDM6A is a histone demethylase that belongs to the KDM6 family of histone H3 lysine demethylases that also includes KDM6B and KDM6C⁷⁴. Methylation of histone lysine and arginine residues functions to regulate transcription and the DNA damage response, specifically in the recruitment of DNA repair proteins and transcriptional repression⁵⁰. KDM6A removes methylation of di- and trimethylated histone 3 lysine 27 (H3K27)^{49,74}. KDM6A also interacts with various transcription factors as well as KMT2C, KMT2D, and CBP/p300 chromatin-modifying enzymes, and the SWI/SNF chromatin-remodeling complex to facilitate transcriptional regulation⁷⁴. Mutations in KDM6A lead to activation of the histone methyltransferase, EZH2, resulting in transcriptional repression⁷⁴. KDM6A is believed to function as a tumor suppressor by antagonizing EZH2-mediated transcriptional repression and promoting transcriptional regulation^{74,75}.

Alterations and prevalence: Somatic mutations in KDM6A are observed in 26% of bladder urothelial carcinoma, 7% of uterine corpus endometrial carcinoma, 5% of skin cutaneous melanoma, lung squamous cell carcinoma, and 4% of esophageal adenocarcinoma, kidney renal papillary cell carcinoma, pancreatic adenocarcinoma, cervical squamous cell carcinoma, and head and neck squamous cell carcinoma^{5,6}. Biallelic loss of KDM6A is observed in 8% of esophageal adenocarcinoma, 4% of lung squamous cell carcinoma, 3% of head and neck squamous cell carcinoma, bladder urothelial carcinoma, and pancreatic adenocarcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for KDM6A aberrations. Pre-clinical data suggest that KDM6A loss of function or inactivating mutations may respond to EZH2 inhibitors⁷⁵.

RBM10 deletion

RNA binding motif protein 10

Background: RBM10 encodes RNA binding motif protein 10, a member of the RNA binding proteins (RBP) family^{1,45}. RBM10 regulates RNA splicing and post-transcriptional modification of mRNA^{45,46}. RBM10 is suggested to function as a tumor suppressor by promoting apoptosis and inhibiting cellular proliferation through regulation of the MDM2 and p53 feedback loops, as well as influencing BAX

Biomarker Descriptions (continued)

expression⁴⁵. RBM10 has been observed to promote transformation and proliferation in lung cancer, supporting an oncogenic role for RBM10^{47,48}.

Alterations and prevalence: Somatic mutations in RBM10 are observed in 7% of lung adenocarcinoma, 6% of uterine corpus endometrial carcinoma, 4% of bladder urothelial carcinoma, 3% of colorectal adenocarcinoma and skin cutaneous melanoma, and 2% of diffuse large B-cell lymphoma, pancreatic adenocarcinoma, adrenocortical carcinoma, cervical squamous cell carcinoma, esophageal adenocarcinoma, stomach adenocarcinoma, and kidney chromophobe^{5,6}. Biallelic loss of RBM10 is observed in 3% of esophageal adenocarcinoma and 2% of head and neck squamous cell carcinoma^{5,6}. Amplification of RBM10 is observed in 5% of ovarian serous cystadenocarcinoma, 4% of uterine carcinosarcoma, and 2% of sarcoma, uterine corpus endometrial carcinoma, adrenocortical carcinoma, and diffuse large B-cell lymphoma^{5,6}.

Potential relevance: Currently, no therapies are approved for RBM10 aberrations.

KDM5C deletion

lysine demethylase 5C

Background: The KDM5C gene encodes the lysine demethylase 5C protein, a histone demethylase, also known as JARID1C^{1,49}. Methylation of histone lysine and arginine residues functions to regulate transcription and DNA damage response⁵⁰. KDM5C removes methylation of di- and trimethylated histone H3 lysine 4 (H3K4) and is involved in the repression of transcription in response to DNA damage^{49,50}. KDM5C alterations result in aberrant H3K4 trimethylation at active replication origins which can lead to stalled DNA replication⁵¹.

Alterations and prevalence: Somatic mutations in KDM5C are observed in 9% of uterine corpus endometrial carcinoma, 5% of kidney renal clear cell carcinoma, stomach adenocarcinoma, skin cutaneous melanoma, 4% of lung adenocarcinoma and uterine carcinosarcoma^{5,6}. Biallelic loss of KDM5C is observed in 3% of esophageal adenocarcinoma and 2% of head and neck squamous cell carcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for KDM5C aberrations.

SMC1A deletion

structural maintenance of chromosomes 1A

Background: SMC1A encodes the structural maintenance of chromosomes 1A and belongs to structural maintenance of chromosomes (SMCs) family, which consists of SMC1A, SMC1B, SMC2, SMC3, SMC4, SMC5, and SMC6^{1,82,83}. As a part of the cohesion-core complex, SMC1A plays a crucial role in chromosome segregation during mitosis and meiosis^{82,84}. SMC1A also plays a role in cell cycle regulation, DNA damage repair, gene transcription regulation, and genomic organization⁸². SMC1A aberrations, including overexpression, have been observed in several cancer types and have been proposed to promote tumor formation and epithelial to mesenchymal transition^{83,85}.

Alterations and prevalence: Somatic mutations in SMC1A are observed in 11% of uterine corpus endometrial carcinoma, 5% of skin cutaneous melanoma and acute myeloid leukemia, 4% of colorectal adenocarcinoma and bladder urothelial carcinoma, 3% cervical squamous cell carcinoma and glioblastoma multiforme, 2% diffuse large B-Cell lymphoma, adrenocortical carcinoma, stomach adenocarcinoma, uterine carcinosarcoma, ovarian serous cystadenocarcinoma and lung adenocarcinoma^{5,6}. Amplification of SMC1A is found in 4% of diffuse large B-Cell lymphoma, 3% of sarcoma, and 2% of ovarian serous cystadenocarcinoma, adrenocortical carcinoma, and uterine carcinosarcoma^{5,6}. Biallelic loss of SMC1A is found in 3% of esophageal adenocarcinoma and 2% of head and neck squamous cell carcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for SMC1A aberrations.

AMER1 deletion

APC membrane recruitment protein 1

Background: The AMER1 gene encodes APC membrane recruitment protein 1¹. AMER1 works in complex with CTNNB1, APC, AXIN1, and AXIN2 to regulate the WNT pathway^{1,107}. The WNT signaling pathway is responsible for regulating several key components during embryogenesis and has been observed to be involved in tumorigenesis^{108,109}. Consequently, the WNT signaling pathway is a target for therapeutic response in various cancer types¹⁰⁹. The AMER1 gene is located on the X chromosome and is commonly inactivated in Wilms tumor, a pediatric kidney cancer¹¹⁰. AMER1 has also been observed to influence cell proliferation, tumorigenesis, migration, invasion, and cell cycle arrest¹⁰⁷.

Biomarker Descriptions (continued)

Alterations and prevalence: Somatic mutations of AMER1 are observed in 13% of colorectal adenocarcinoma, 10% of uterine corpus endometrial carcinoma, 8% of skin cutaneous melanoma, 7% of lung adenocarcinoma, 4% of stomach adenocarcinoma, and uterine carcinosarcoma, 3% of lung squamous cell carcinoma, cervical squamous cell carcinoma, bladder urothelial carcinoma, and 2% of diffuse large B-cell lymphoma, liver hepatocellular carcinoma, head and neck squamous cell carcinoma, and breast invasive carcinoma^{5,6}. Biallelic deletion of AMER1 is observed in 2% of esophageal adenocarcinoma, diffuse large b-cell lymphoma, uterine carcinosarcoma, lung squamous cell carcinoma, and pancreatic adenocarcinoma, and 1% of stomach adenocarcinoma, sarcoma, liver hepatocellular carcinoma, colorectal adenocarcinoma, head and neck squamous cell carcinoma, uterine corpus endometrial carcinoma, and ovarian serous cystadenocarcinoma^{5,6}.

Potential relevance: Currently, no therapies are approved for AMER1 aberrations.

ZMYM3 deletion

zinc finger MYM-type containing 3

Background: The ZMYM3 gene encodes the zinc finger MYM-type containing 3 protein¹. While the function is not fully understood, ZMYM3 is capable of binding histones and DNA, and may facilitate the repair of double-strand breaks (DSBs)¹¹¹.

Alterations and prevalence: Somatic mutations in ZMYM3 are observed in 12% of uterine corpus endometrial carcinoma, 5% of skin cutaneous melanoma, 4% of colorectal adenocarcinoma, 3% of lung adenocarcinoma, lung squamous cell carcinoma, cervical squamous cell carcinoma, esophageal adenocarcinoma, and bladder urothelial carcinoma^{5,6}. In prostate cancer, ZMYM3 mutations have been observed to be enriched in African American men compared to white men with one study demonstrating occurrence in 11.7% vs. 2.7% of patients, respectively¹¹². Biallelic deletion of ZMYM3 is observed in 3% of cholangiocarcinoma and 2% of sarcoma and kidney chromophobe^{5,6}.

Potential relevance: Currently, no therapies are approved for ZMYM3 aberrations.

STAG2 deletion

stromal antigen 2

Background: The STAG2 gene encodes the stromal antigen 2 protein, one of the core proteins in the cohesin complex, which regulates the separation of sister chromatids during cell division^{65,66}. Components of the cohesion complex include SMC1A, SMC3, and RAD21, which bind to STAG1/STAG2 paralogs^{67,68}. Inactivating mutations in STAG2 contribute to X-linked neurodevelopmental disorders, aneuploidy, and chromosomal instability in cancer^{67,69}.

Alterations and prevalence: Somatic mutations in STAG2 include nonsense, frameshift, splice site variants³⁰. Somatic mutations in STAG2 are observed in various solid tumors including 14% of bladder cancer, 10% of uterine cancer, 3% of stomach cancer, and 4% of lung adenocarcinoma⁶. In addition, mutations in STAG2 are observed in 5-10% of myelodysplastic syndrome (MDS), 3% of acute myeloid leukemia, and 2% of diffuse large B-cell lymphoma^{6,30}.

Potential relevance: Mutations in STAG2 are associated with poor prognosis and adverse risk in MDS and Acute Myeloid Leukemia^{30,40,41}. Truncating mutations in STAG2 lead to a loss of function in bladder cancer and are often identified as an early event associated with low grade and stage tumors⁷⁰.

PHF6 deletion

PHD finger protein 6

Background: The PHF6 gene encodes the plant homeodomain (PHD) finger protein 6 which contains four nuclear localization signals and two imperfect PHD zinc finger domains. PHF6 is a tumor suppressor that interacts with the nucleosome remodeling deacetylase (NuRD) complex, which regulates nucleosome positioning and transcription of genes involved in development and cell-cycle progression^{76,77}.

Alterations and prevalence: The majority of PHF6 aberrations are nonsense, frameshift (70%), or missense (30%) mutations, which result in complete loss of protein expression^{76,78,79,80}. Truncating or missense mutations in PHF6 are observed in 38% of adult and 16% of pediatric T-cell acute lymphoblastic leukemia (T-ALL), 20-25% of mixed phenotype acute leukemias (MPAL), and 3% of AML, and 2.6% of hepatocellular carcinoma (HCC)^{78,80}. Missense mutations recurrently involve codon C215 and the second zinc finger domain of PHF6⁷⁸. PHF6 mutations are frequently observed in hematologic malignancies from male patients^{76,78}.

Potential relevance: Somatic mutations in PHF6 are associated with reduced overall survival in AML patients treated with high-dose induction chemotherapy⁸¹.

Alerts Informed By Public Data Sources

Current FDA Information

Contraindicated Not recommended Resistance Breakthrough Fast Track

FDA information is current as of 2025-05-14. For the most up-to-date information, search www.fda.gov.

KRAS p.(G12V) c.35G>T

cetuximab

Cancer type: Colorectal Cancer Label as of: 2021-09-24 Variant class: KRAS G12 mutation

Indications and usage:

Erbixux® is an epidermal growth factor receptor (EGFR) antagonist indicated for treatment of:

Head and Neck Cancer

- Locally or regionally advanced squamous cell carcinoma of the head and neck in combination with radiation therapy.
- Recurrent locoregional disease or metastatic squamous cell carcinoma of the head and neck in combination with platinum-based therapy with fluorouracil.
- Recurrent or metastatic squamous cell carcinoma of the head and neck progressing after platinum-based therapy.

Colorectal Cancer

K-Ras wild-type, EGFR-expressing, metastatic colorectal cancer as determined by FDA-approved test

- in combination with FOLFIRI for first-line treatment,
- in combination with irinotecan in patients who are refractory to irinotecan-based chemotherapy,
- as a single agent in patients who have failed oxaliplatin- and irinotecan-based chemotherapy or who are intolerant to irinotecan.

Limitations of Use: Erbixux® is not indicated for treatment of Ras-mutant colorectal cancer or when the results of the Ras mutation tests are unknown.

BRAF V600E Mutation-Positive Metastatic Colorectal Cancer (CRC)

- in combination with encorafenib, for the treatment of adult patients with metastatic colorectal cancer (CRC) with a BRAF V600E mutation, as detected by an FDA-approved test, after prior therapy.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf

KRAS p.(G12V) c.35G>T (continued)**🚫 panitumumab****Cancer type:** Colorectal Cancer**Label as of:** 2025-01-16**Variant class:** KRAS G12 mutation**Indications and usage:**

VECTIBIX® is an epidermal growth factor receptor (EGFR) antagonist indicated for the treatment of:

Adult patients with wild-type RAS (defined as wild-type in both KRAS and NRAS as determined by an FDA-approved test) Metastatic Colorectal Cancer (mCRC)*:

- In combination with FOLFOX for first-line treatment.
- As monotherapy following disease progression after prior treatment with fluoropyrimidine, oxaliplatin, and irinotecan-containing chemotherapy.

KRAS G12C-mutated Metastatic Colorectal Cancer (mCRC)*

- In combination with sotorasib, for the treatment of adult patients with KRAS G12C-mutated mCRC, as determined by an FDA-approved test, who have received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.

*Limitations of Use: VECTIBIX® is not indicated for the treatment of patients with RAS-mutant mCRC unless used in combination with sotorasib in KRAS G12C-mutated mCRC. VECTIBIX® is not indicated for the treatment of patients with mCRC for whom RAS mutation status is unknown.

Reference:

https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125147s213lbl.pdf

Current NCCN Information**🚫** Contraindicated**⊖** Not recommended**🛡️** Resistance**🚀** Breakthrough**🏠** Fast Track

NCCN information is current as of 2025-05-01. To view the most recent and complete version of the guideline, go online to NCCN.org.

For NCCN International Adaptations & Translations, search www.nccn.org/global/what-we-do/international-adaptations.

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

KRAS p.(G12V) c.35G>T**🚫 cetuximab****Cancer type:** Colon Cancer**Variant class:** KRAS G12 mutation**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

Reference: NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2025]

KRAS p.(G12V) c.35G>T (continued)**❌ cetuximab**

Cancer type: Rectal Cancer

Variant class: KRAS G12 mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exons 2, 3, and 4) or NRAS mutation (exons 2, 3, and 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 2.2025]

❌ panitumumab

Cancer type: Colon Cancer

Variant class: KRAS G12 mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

Reference: NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2025]

❌ panitumumab

Cancer type: Rectal Cancer

Variant class: KRAS G12 mutation

Summary:

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exons 2, 3, and 4) or NRAS mutation (exons 2, 3, and 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 2.2025]

Current EMA Information**❌** Contraindicated**⊖** Not recommended**🛡** Resistance**🚀** Breakthrough**A** Fast TrackEMA information is current as of 2025-05-14. For the most up-to-date information, search www.ema.europa.eu.**KRAS p.(G12V) c.35G>T****❌ cetuximab, cetuximab + oxaliplatin**

Cancer type: Colorectal Cancer

Label as of: 2025-01-16

Variant class: KRAS G12 mutation

Reference:https://www.ema.europa.eu/en/documents/product-information/erbitux-epar-product-information_en.pdf**❌ panitumumab + oxaliplatin**

Cancer type: Colorectal Cancer

Label as of: 2025-05-07

Variant class: KRAS G12 mutation

Reference:https://www.ema.europa.eu/en/documents/product-information/vectibix-epar-product-information_en.pdf

Current ESMO Information

Contraindicated Not recommended Resistance Breakthrough Fast Track

ESMO information is current as of 2025-05-01. For the most up-to-date information, search www.esmo.org.

KRAS p.(G12V) c.35G>T

cetuximab

Cancer type: Colorectal Cancer Variant class: KRAS G12 mutation

Summary:
ESMO Clinical Practice Guidelines include the following supporting statement:

- "The presence of RAS mutations is associated with resistance to anti-EGFR mAbs and knowing the expanded RAS mutational status is mandatory for use of both cetuximab and panitumumab, avoiding anti-EGFR mAb treatment when a RAS mutation is confirmed".
- "RAS testing is mandatory before treatment with anti-EGFR mAbs and can be carried out on either the primary tumor or other metastatic sites [III, A]".

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2022.10.003> (published)]

panitumumab

Cancer type: Colorectal Cancer Variant class: KRAS G12 mutation

Summary:
ESMO Clinical Practice Guidelines include the following supporting statement:

- "The presence of RAS mutations is associated with resistance to anti-EGFR mAbs and knowing the expanded RAS mutational status is mandatory for use of both cetuximab and panitumumab, avoiding anti-EGFR mAb treatment when a RAS mutation is confirmed".
- "RAS testing is mandatory before treatment with anti-EGFR mAbs and can be carried out on either the primary tumor or other metastatic sites [III, A]".

Reference: ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2022.10.003> (published)]

Genes Assayed

Genes Assayed for the Detection of DNA Sequence Variants

ABL1, ABL2, ACVR1, AKT1, AKT2, AKT3, ALK, AR, ARAF, ATP1A1, AURKA, AURKB, AURKC, AXL, BCL2, BCL2L12, BCL6, BCR, BMP5, BRAF, BTK, CACNA1D, CARD11, CBL, CCND1, CCND2, CCND3, CCNE1, CD79B, CDK4, CDK6, CHD4, CSF1R, CTNNB1, CUL1, CYSLTR2, DDR2, DGCR8, DROSHA, E2F1, EGFR, EIF1AX, EPAS1, ERBB2, ERBB3, ERBB4, ESR1, EZH2, FAM135B, FGF7, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FOXL2, FOXO1, GATA2, GLI1, GNA11, GNAQ, GNAS, HIF1A, HRAS, IDH1, IDH2, IKBKB, IL6ST, IL7R, IRF4, IRS4, KCNJ5, KDR, KIT, KLF4, KLF5, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAPK1, MAX, MDM4, MECOM, MED12, MEF2B, MET, MITF, MPL, MTOR, MYC, MYCN, MYD88, MYO10, NFE2L2, NRAS, NSD2, NT5C2, NTRK1, NTRK2, NTRK3, NUP93, PAX5, PCBP1, PDGFRA, PDGFRB, PIK3C2B, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PIK3R2, PIM1, PLCG1, PPP2R1A, PPP6C, PRKACA, PTPN11, PTPRD, PXDNL, RAC1, RAF1, RARA, RET, RGS7, RHEB, RHOA, RICTOR, RIT1, ROS1, RPL10, SETBP1, SF3B1, SIX1, SIX2, SLC01B3, SMC1A, SMO, SNCAIP, SOS1, SOX2, SPOP, SRC, SRSF2, STAT3, STAT5B, STAT6, TAF1, TERT, TGFB1, TOP1, TOP2A, TPMT, TRRAP, TSHR, U2AF1, USP8, WAS, XPO1, ZNF217, ZNF429

Genes Assayed (continued)

Genes Assayed for the Detection of Copy Number Variations

ABCB1, ABL1, ABL2, ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AKT1, AKT2, AKT3, ALK, AMER1, APC, AR, ARAF, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AURKA, AURKC, AXIN1, AXIN2, AXL, B2M, BAP1, BARD1, BCL2, BCL2L12, BCL6, BCOR, BLM, BMPR2, BRAF, BRCA1, BRCA2, BRIP1, CARD11, CASP8, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDK4, CDK6, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHD4, CHEK1, CHEK2, CIC, CREBBP, CSMD3, CTCF, CTLA4, CTNND2, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, DAXX, DDR1, DDR2, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, EGFR, EIF1AX, ELF3, EMSY, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERBB2, ERBB3, ERBB4, ERCC2, ERCC4, ERRFI1, ESR1, ETV6, EZH2, FAM135B, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAT1, FBXW7, FGF19, FGF23, FGF3, FGF4, FGF9, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FUBP1, FYN, GATA2, GATA3, GLI3, GNA13, GNAS, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, IDH2, IGF1R, IKBKB, IL7R, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KDR, KEAP1, KIT, KLF5, KMT2A, KMT2B, KMT2C, KMT2D, KRAS, LARP4B, LATS1, LATS2, MAGOH, MAP2K1, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK1, MAPK8, MAX, MCL1, MDM2, MDM4, MECOM, MEF2B, MEN1, MET, MGA, MITF, MLH1, MLH3, MPL, MRE11, MSH2, MSH3, MSH6, MTAP, MTOR, MUTYH, MYC, MYCL, MYCN, MYD88, NBN, NCOR1, NF1, NF2, NFE2L2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, NRAS, NTRK1, NTRK3, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PCBP1, PDCD1, PDCD1LG2, PDGFRA, PDGFRB, PDIA3, PGD, PHF6, PIK3C2B, PIK3CA, PIK3CB, PIK3R1, PIK3R2, PIM1, PLCG1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R1A, PPP2R2A, PPP6C, PRDM1, PRDM9, PRKACA, PRKAR1A, PTCH1, PTEN, PTPN11, PTPRT, PXDN, RAC1, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RAF1, RARA, RASA1, RASA2, RB1, RBM10, RECQL4, RET, RHEB, RICTOR, RIT1, RNASEH2A, RNASEH2B, RNF43, ROS1, RPA1, RPS6KB1, RPTOR, RUNX1, SDHA, SDHB, SDHD, SETBP1, SETD2, SF3B1, SLC01B3, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SMC1A, SMO, SOX9, SPEN, SPOP, SRC, STAG2, STAT3, STAT6, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TERT, TET2, TGFB2, TNFAIP3, TNFRSF14, TOP1, TP53, TP63, TPMT, TPP2, TSC1, TSC2, U2AF1, USP8, USP9X, VHL, WT1, XPO1, XRCC2, XRCC3, YAP1, YES1, ZFH3, ZMYM3, ZNF217, ZNF429, ZRSR2

Genes Assayed for the Detection of Fusions

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSPO2, RSPO3, TERT

Genes Assayed with Full Exon Coverage

ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AMER1, APC, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AXIN1, AXIN2, B2M, BAP1, BARD1, BCOR, BLM, BMPR2, BRCA1, BRCA2, BRIP1, CALR, CASP8, CBFB, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHEK1, CHEK2, CIC, CIITA, CREBBP, CSMD3, CTCF, CTLA4, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, CYP2D6, DAXX, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, ELF3, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERCC2, ERCC4, ERCC5, ERRFI1, ETV6, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAS, FAT1, FBXW7, FUBP1, GATA3, GNA13, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, ID3, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KEAP1, KLHL13, KMT2A, KMT2B, KMT2C, KMT2D, LARP4B, LATS1, LATS2, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK8, MEN1, MGA, MLH1, MLH3, MRE11, MSH2, MSH3, MSH6, MTAP, MTUS2, MUTYH, NBN, NCOR1, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PDCD1, PDCD1LG2, PDIA3, PGD, PHF6, PIK3R1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R2A, PRDM1, PRDM9, PRKAR1A, PSMB10, PSMB8, PSMB9, PTCH1, PTEN, PTPRT, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RASA1, RASA2, RB1, RBM10, RECQL4, RNASEH2A, RNASEH2B, RNASEH2C, RNF43, RPA1, RPL22, RPL5, RUNX1, RUNX1T1, SDHA, SDHB, SDHC, SDHD, SETD2, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SOCS1, SOX9, SPEN, STAG2, STAT1, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TET2, TGFB2, TMEM132D, TNFAIP3, TNFRSF14, TP53, TP63, TPP2, TSC1, TSC2, UGT1A1, USP9X, VHL, WT1, XRCC2, XRCC3, ZBTB20, ZFH3, ZMYM3, ZRSR2

Relevant Therapy Summary

 In this cancer type
  In other cancer type
  In this cancer type and other cancer types
  No evidence

KRAS p.(G12V) c.35G>T

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
bevacizumab + CAPOX	×	×	×	○	×
bevacizumab + FOLFIRI	×	×	×	○	×
bevacizumab + FOLFOX	×	×	×	○	×
bevacizumab + FOLFOXIRI	×	×	×	○	×
RMC-6236	×	×	×	×	● (III)
daratumumab, TG-01 (Targovax), QS-21 Stimulon, nivolumab	×	×	×	×	● (II)
regorafenib	×	×	×	×	● (II)
afatinib, selumetinib	×	×	×	×	● (I/II)
anti-KRAS G12V mTCR, chemotherapy, aldesleukin	×	×	×	×	● (I/II)
ERAS-0015	×	×	×	×	● (I/II)
IMM-1-104	×	×	×	×	● (I/II)
YL-15293	×	×	×	×	● (I/II)
zotatifin	×	×	×	×	● (I/II)
AFNT-211	×	×	×	×	● (I)
HMPL-415	×	×	×	×	● (I)
JAB-3312	×	×	×	×	● (I)
JSI-1187	×	×	×	×	● (I)
KRAS peptide vaccine, poly-ICLC, nivolumab, ipilimumab	×	×	×	×	● (I)
KRAS TCR, aldesleukin, SLATE 001, chemotherapy	×	×	×	×	● (I)
KRAS-EphA-2-CAR-DC, anti-PD-1, ipilimumab	×	×	×	×	● (I)
Nest-1	×	×	×	×	● (I)
NW-301V	×	×	×	×	● (I)
ZEN-3694, binimetinib	×	×	×	×	● (I)

CDKN2A deletion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
palbociclib	×	×	×	×	● (II)
palbociclib, abemaciclib	×	×	×	×	● (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

CDKN2A deletion (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
AMG 193	×	×	×	×	<div></div> (I/II)

ATRX deletion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
pamiparib, tislelizumab	×	×	×	×	<div></div> (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Thermo Fisher Scientific's Ion Torrent OncoPrint Reporter software was used in generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on OncoPrint Reporter (6.1.1 data version 2025.06(006)). The data presented here are from a curated knowledge base of publicly available information, but may not be exhaustive. FDA information was sourced from www.fda.gov and is current as of 2025-05-14. NCCN information was sourced from www.nccn.org and is current as of 2025-05-01. EMA information was sourced from www.ema.europa.eu and is current as of 2025-05-14. ESMO information was sourced from www.esmo.org and is current as of 2025-05-01. Clinical Trials information is current as of 2025-05-01. For the most up-to-date information regarding a particular trial, search www.clinicaltrials.gov by NCT ID or search local clinical trials authority website by local identifier listed in 'Other identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials listed in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Sarikas et al. The cullin protein family. *Genome Biol.* 2011;12(4):220. PMID: 21554755
3. Sang et al. The role and mechanism of CRL4 E3 ubiquitin ligase in cancer and its potential therapy implications. *Oncotarget.* 2015 Dec 15;6(40):42590-602. PMID: 26460955
4. Cheng et al. The emerging role for Cullin 4 family of E3 ligases in tumorigenesis. *Biochim Biophys Acta Rev Cancer.* 2019 Jan;1871(1):138-159. PMID: 30602127
5. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
6. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
7. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer.* 2011 Oct 13;11(11):761-74. PMID: 21993244
8. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
9. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
10. Román et al. KRAS oncogene in non-small cell lung cancer: clinical perspectives on the treatment of an old target. *Mol Cancer.* 2018 Feb 19;17(1):33. doi: 10.1186/s12943-018-0789-x. PMID: 29455666
11. Dinu et al. Prognostic significance of KRAS gene mutations in colorectal cancer--preliminary study. *J Med Life.* 2014 Oct-Dec;7(4):581-7. PMID: 25713627
12. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
13. https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/214665Orig1s009correctedlbl.pdf
14. https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/216340s005lbl.pdf
15. NCCN Guidelines® - NCCN-Pancreatic Adenocarcinoma [Version 2.2025]
16. <https://assets.cwp.roche.com/f/126832/x/5738a7538b/irp230202.pdf>
17. <https://bridgebio.com/news/bridgebio-pharma-announces-first-lung-cancer-patient-dosed-in-phase-1-2-trial-and-us-fda-fast-track-designation-for-shp2-inhibitor-bbp-398-in-combination-with-amgens-lumakras-sotorasib/>
18. <https://investor.verastem.com/news-releases/news-release-details/verastem-oncology-granted-fast-track-designation-combination>
19. <https://www.businesswire.com/news/home/20250109170439/en/>
20. <https://www.d3bio.com/press-releases/d3-bios-d3s-001-receives-u-s-fda-fast-track-designation-for-the-treatment-of-colorectal-cancer-with-kras-g12c-mutation>
21. https://cardiffoncology.com/wp-content/uploads/2021/07/Cardiff_Oncology_Investor_Presentation_July_2021.pdf
22. https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf
23. https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125147s213lbl.pdf
24. Slebos et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. *N. Engl. J. Med.* 1990 Aug 30;323(9):561-5. PMID: 2199829
25. Gearhart et al. Polycomb group and SCF ubiquitin ligases are found in a novel BCOR complex that is recruited to BCL6 targets. *Mol. Cell. Biol.* 2006 Sep;26(18):6880-9. PMID: 16943429
26. Huynh et al. BCOR, a novel corepressor involved in BCL-6 repression. *Genes Dev.* 2000 Jul 15;14(14):1810-23. PMID: 10898795
27. Kelly et al. Bcor loss perturbs myeloid differentiation and promotes leukaemogenesis. *Nat Commun.* 2019 Mar 22;10(1):1347. PMID: 30902969
28. Cao et al. BCOR regulates myeloid cell proliferation and differentiation. *Leukemia.* 2016 May;30(5):1155-65. PMID: 26847029
29. Yamamoto et al. Clarifying the impact of polycomb complex component disruption in human cancers. *Mol. Cancer Res.* 2014 Apr;12(4):479-84. PMID: 24515802
30. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 2.2025]
31. Damm et al. BCOR and BCORL1 mutations in myelodysplastic syndromes and related disorders. *Blood.* 2013 Oct 31;122(18):3169-77. PMID: 24047651
32. Terada et al. Usefulness of BCOR gene mutation as a prognostic factor in acute myeloid leukemia with intermediate cytogenetic prognosis. *Genes Chromosomes Cancer.* 2018 Aug;57(8):401-408. PMID: 29663558

References (continued)

33. Wong et al. Clear cell sarcomas of the kidney are characterised by BCOR gene abnormalities, including exon 15 internal tandem duplications and BCOR-CCNB3 gene fusion. *Histopathology*. 2018 Jan;72(2):320-329. PMID: 28833375
34. Cramer et al. Successful Treatment of Recurrent Primitive Myxoid Mesenchymal Tumor of Infancy With BCOR Internal Tandem Duplication. *J Natl Compr Canc Netw*. 2017 Jul;15(7):868-871. PMID: 28687574
35. Peters et al. BCOR-CCNB3 fusions are frequent in undifferentiated sarcomas of male children. *Mod. Pathol*. 2015 Apr;28(4):575-86. PMID: 25360585
36. Puls et al. BCOR-CCNB3 (Ewing-like) sarcoma: a clinicopathologic analysis of 10 cases, in comparison with conventional Ewing sarcoma. *Am. J. Surg. Pathol*. 2014 Oct;38(10):1307-18. PMID: 24805859
37. Kao et al. BCOR-CCNB3 Fusion Positive Sarcomas: A Clinicopathologic and Molecular Analysis of 36 Cases With Comparison to Morphologic Spectrum and Clinical Behavior of Other Round Cell Sarcomas. *Am. J. Surg. Pathol*. 2018 May;42(5):604-615. PMID: 29300189
38. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 5.2024]
39. NCCN Guidelines® - NCCN-Bone Cancer [Version 2.2025]
40. NCCN Guidelines® - NCCN-Acute Myeloid Leukemia [Version 2.2025]
41. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood*. 2022 Sep 22;140(12):1345-1377. PMID: 35797463
42. Khoury et al. The 5th edition of the World Health Organization Classification of Haematolymphoid Tumours: Myeloid and Histiocytic/Dendritic Neoplasms. *Leukemia*. 2022 Jul;36(7):1703-1719. PMID: 35732831
43. Torre et al. Recurrent EP300-BCOR Fusions in Pediatric Gliomas With Distinct Clinicopathologic Features. *J Neuropathol Exp Neurol*. 2019 Apr 1;78(4):305-314. PMID: 30816933
44. Wang et al. Clinical, pathological, and molecular features of central nervous system tumors with BCOR internal tandem duplication. *Pathol Res Pract*. 2024 Jul;259:155367. PMID: 38797130
45. Cao et al. RBM10 Regulates Tumor Apoptosis, Proliferation, and Metastasis. *Front Oncol*. 2021;11:603932. PMID: 33718153
46. Zhang et al. RNA binding motif protein 10 suppresses lung cancer progression by controlling alternative splicing of eukaryotic translation initiation factor 4H. *EBioMedicine*. 2020 Nov;61:103067. PMID: 33130397
47. Sun et al. Functional role of RBM10 in lung adenocarcinoma proliferation. *Int J Oncol*. 2019 Feb;54(2):467-478. PMID: 30483773
48. Loiselle et al. RBM10 promotes transformation-associated processes in small cell lung cancer and is directly regulated by RBM5. *PLoS One*. 2017;12(6):e0180258. PMID: 28662214
49. Iwase et al. The X-linked mental retardation gene SMCX/JARID1C defines a family of histone H3 lysine 4 demethylases. *Cell*. 2007 Mar 23;128(6):1077-88. PMID: 17320160
50. Gong et al. Histone methylation and the DNA damage response. *Mutat Res*. 2017 Sep 23;780:37-47. PMID: 31395347
51. Rondinelli et al. H3K4me3 demethylation by the histone demethylase KDM5C/JARID1C promotes DNA replication origin firing. *Nucleic Acids Res*. 2015 Mar 11;43(5):2560-74. PMID: 25712104
52. Ryan et al. Snf2-family proteins: chromatin remodellers for any occasion. *Curr Opin Chem Biol*. 2011 Oct;15(5):649-56. PMID: 21862382
53. Heyer et al. Rad54: the Swiss Army knife of homologous recombination?. *Nucleic Acids Res*. 2006;34(15):4115-25. PMID: 16935872
54. Matsuda et al. Mutations in the RAD54 recombination gene in primary cancers. *Oncogene*. 1999 Jun 3;18(22):3427-30. PMID: 10362365
55. Abedalthagafi et al. The alternative lengthening of telomere phenotype is significantly associated with loss of ATRX expression in high-grade pediatric and adult astrocytomas: a multi-institutional study of 214 astrocytomas. *Mod. Pathol*. 2013 Nov;26(11):1425-32. PMID: 23765250
56. Clynes et al. ATRX dysfunction induces replication defects in primary mouse cells. *PLoS ONE*. 2014;9(3):e92915. PMID: 24651726
57. Tang et al. A novel transcription regulatory complex containing death domain-associated protein and the ATR-X syndrome protein. *J. Biol. Chem*. 2004 May 7;279(19):20369-77. PMID: 14990586
58. Xue et al. The ATRX syndrome protein forms a chromatin-remodeling complex with Daxx and localizes in promyelocytic leukemia nuclear bodies. *Proc. Natl. Acad. Sci. U.S.A.* 2003 Sep 16;100(19):10635-40. PMID: 12953102
59. Pisapia. The Updated World Health Organization Glioma Classification: Cellular and Molecular Origins of Adult Infiltrating Gliomas. *Arch. Pathol. Lab. Med*. 2017 Dec;141(12):1633-1645. PMID: 29189064
60. Jiao et al. Frequent ATRX, CIC, FUBP1 and IDH1 mutations refine the classification of malignant gliomas. *Oncotarget*. 2012 Jul;3(7):709-22. PMID: 22869205

References (continued)

61. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol.* 2021 Aug 2;23(8):1231-1251. PMID: 34185076
62. NCCN Guidelines® - NCCN-Central Nervous System Cancers [Version 5.2024]
63. Dufner et al. Ubiquitin-specific protease 8 (USP8/UBPy): a prototypic multidomain deubiquitinating enzyme with pleiotropic functions. *Biochem Soc Trans.* 2019 Dec 20;47(6):1867-1879. PMID: 31845722
64. Lu et al. USP9X stabilizes BRCA1 and confers resistance to DNA-damaging agents in human cancer cells. *Cancer Med.* 2019 Nov;8(15):6730-6740. PMID: 31512408
65. Mehta et al. Cohesin: functions beyond sister chromatid cohesion. *FEBS Lett.* 2013 Aug 2;587(15):2299-312. PMID: 23831059
66. Aquila et al. The role of STAG2 in bladder cancer. *Pharmacol. Res.* 2018 May;131:143-149. PMID: 29501732
67. Mullegama et al. De novo loss-of-function variants in STAG2 are associated with developmental delay, microcephaly, and congenital anomalies. *Am. J. Med. Genet. A.* 2017 May;173(5):1319-1327. PMID: 28296084
68. van et al. Synthetic lethality between the cohesin subunits STAG1 and STAG2 in diverse cancer contexts. *Elife.* 2017 Jul 10;6. PMID: 28691904
69. Solomon et al. Mutational inactivation of STAG2 causes aneuploidy in human cancer. *Science.* 2011 Aug 19;333(6045):1039-43. PMID: 21852505
70. Solomon et al. Frequent truncating mutations of STAG2 in bladder cancer. *Nat. Genet.* 2013 Dec;45(12):1428-30. PMID: 24121789
71. Madan et al. Aberrant splicing of U12-type introns is the hallmark of ZRSR2 mutant myelodysplastic syndrome. *Nat Commun.* 2015 Jan 14;6:6042. doi: 10.1038/ncomms7042. PMID: 25586593
72. Tronchère et al. A protein related to splicing factor U2AF35 that interacts with U2AF65 and SR proteins in splicing of pre-mRNA. *Nature.* 1997 Jul 24;388(6640):397-400. PMID: 9237760
73. Chesnais et al. Spliceosome mutations in myelodysplastic syndromes and chronic myelomonocytic leukemia. *Oncotarget.* 2012 Nov;3(11):1284-93. PMID: 23327988
74. Tran et al. Lysine Demethylase KDM6A in Differentiation, Development, and Cancer. *Mol Cell Biol.* 2020 Sep 28;40(20). PMID: 32817139
75. Ler et al. Loss of tumor suppressor KDM6A amplifies PRC2-regulated transcriptional repression in bladder cancer and can be targeted through inhibition of EZH2. *Sci Transl Med.* 2017 Feb 22;9(378). PMID: 28228601
76. Wendorff et al. Phf6 Loss Enhances HSC Self-Renewal Driving Tumor Initiation and Leukemia Stem Cell Activity in T-ALL. *Cancer Discov.* 2019 Mar;9(3):436-451. PMID: 30567843
77. Lower et al. Mutations in PHF6 are associated with Börjeson-Forssman-Lehmann syndrome. *Nat. Genet.* 2002 Dec;32(4):661-5. PMID: 12415272
78. Van et al. PHF6 mutations in T-cell acute lymphoblastic leukemia. *Nat. Genet.* 2010 Apr;42(4):338-42. PMID: 20228800
79. Van et al. PHF6 mutations in adult acute myeloid leukemia. *Leukemia.* 2011 Jan;25(1):130-4. PMID: 21030981
80. Yoo et al. Somatic mutation of PHF6 gene in T-cell acute lymphoblastic leukemia, acute myelogenous leukemia and hepatocellular carcinoma. *Acta Oncol.* 2012 Jan;51(1):107-11. PMID: 21736506
81. Patel et al. Prognostic relevance of integrated genetic profiling in acute myeloid leukemia. *N. Engl. J. Med.* 2012 Mar 22;366(12):1079-89. PMID: 22417203
82. Musio. The multiple facets of the SMC1A gene. *Gene.* 2020 Jun 15;743:144612. PMID: 32222533
83. Nie et al. Clinical Significance and Integrative Analysis of the SMC Family in Hepatocellular Carcinoma. *Front Med (Lausanne).* 2021;8:727965. PMID: 34527684
84. Yatskevich et al. Organization of Chromosomal DNA by SMC Complexes. *Annu Rev Genet.* 2019 Dec 3;53:445-482. PMID: 31577909
85. Yadav et al. SMC1A is associated with radioresistance in prostate cancer and acts by regulating epithelial-mesenchymal transition and cancer stem-like properties. *Mol Carcinog.* 2019 Jan;58(1):113-125. PMID: 30242889
86. Xia et al. Dominant role of CDKN2B/p15INK4B of 9p21.3 tumor suppressor hub in inhibition of cell-cycle and glycolysis. *Nat Commun.* 2021 Apr 6;12(1):2047. PMID: 33824349
87. Scruggs et al. Loss of CDKN2B Promotes Fibrosis via Increased Fibroblast Differentiation Rather Than Proliferation. *Am. J. Respir. Cell Mol. Biol.* 2018 Aug;59(2):200-214. PMID: 29420051
88. Roussel. The INK4 family of cell cycle inhibitors in cancer. *Oncogene.* 1999 Sep 20;18(38):5311-7. PMID: 10498883
89. Aytac et al. Rb independent inhibition of cell growth by p15(INK4B). *Biochem. Biophys. Res. Commun.* 1999 Aug 27;262(2):534-8. PMID: 10462509
90. Hill et al. The genetics of melanoma: recent advances. *Annu Rev Genomics Hum Genet.* 2013;14:257-79. PMID: 23875803

References (continued)

91. Kim et al. The regulation of INK4/ARF in cancer and aging. *Cell*. 2006 Oct 20;127(2):265-75. PMID: 17055429
92. Sekulic et al. Malignant melanoma in the 21st century: the emerging molecular landscape. *Mayo Clin. Proc.* 2008 Jul;83(7):825-46. PMID: 18613999
93. Orlow et al. CDKN2A germline mutations in individuals with cutaneous malignant melanoma. *J. Invest. Dermatol.* 2007 May;127(5):1234-43. PMID: 17218939
94. Bartsch et al. CDKN2A germline mutations in familial pancreatic cancer. *Ann. Surg.* 2002 Dec;236(6):730-7. PMID: 12454511
95. Adib et al. CDKN2A Alterations and Response to Immunotherapy in Solid Tumors. *Clin Cancer Res.* 2021 Jul 15;27(14):4025-4035. PMID: 34074656
96. NCCN Guidelines® - NCCN-Mesothelioma: Peritoneal [Version 2.2025]
97. NCCN Guidelines® - NCCN-Mesothelioma: Pleural [Version 2.2025]
98. Louis et al. cIMPACT-NOW update 6: new entity and diagnostic principle recommendations of the cIMPACT-Utrecht meeting on future CNS tumor classification and grading. *Brain Pathol.* 2020 Jul;30(4):844-856. PMID: 32307792
99. Longwen et al. Frequent genetic aberrations in the cell cycle related genes in mucosal melanoma indicate the potential for targeted therapy. *J Transl Med.* 2019 Jul 29;17(1):245. PMID: 31358010
100. Logan et al. PD-0332991, a potent and selective inhibitor of cyclin-dependent kinase 4/6, demonstrates inhibition of proliferation in renal cell carcinoma at nanomolar concentrations and molecular markers predict for sensitivity. *Anticancer Res.* 2013 Aug;33(8):2997-3004. PMID: 23898052
101. von et al. Preclinical Characterization of Novel Chordoma Cell Systems and Their Targeting by Pharmacological Inhibitors of the CDK4/6 Cell-Cycle Pathway. *Cancer Res.* 2015 Sep 15;75(18):3823-31. PMID: 26183925
102. Cen et al. p16-Cdk4-Rb axis controls sensitivity to a cyclin-dependent kinase inhibitor PD0332991 in glioblastoma xenograft cells. *Neuro-oncology.* 2012 Jul;14(7):870-81. PMID: 22711607
103. Vitzthum et al. The role of p16 as a biomarker in nonoropharyngeal head and neck cancer. *Oncotarget.* 2018 Sep 7;9(70):33247-33248. PMID: 30279955
104. Chung et al. p16 protein expression and human papillomavirus status as prognostic biomarkers of nonoropharyngeal head and neck squamous cell carcinoma. *J. Clin. Oncol.* 2014 Dec 10;32(35):3930-8. PMID: 25267748
105. Bryant et al. Prognostic Role of p16 in Nonoropharyngeal Head and Neck Cancer. *J. Natl. Cancer Inst.* 2018 Dec 1;110(12):1393-1399. PMID: 29878161
106. Stephen et al. Significance of p16 in Site-specific HPV Positive and HPV Negative Head and Neck Squamous Cell Carcinoma. *Cancer Clin Oncol.* 2013;2(1):51-61. PMID: 23935769
107. Liu et al. Aging (Albany NY). 2020 May 4;12(9):8372-8396. PMID: 32365332
108. Komiya et al. Wnt signal transduction pathways. *Organogenesis.* 2008 Apr;4(2):68-75. PMID: 19279717
109. Zhang et al. *J Hematol Oncol.* 2020 Dec 4;13(1):165. PMID: 33276800
110. Rivera et al. An X chromosome gene, WTX, is commonly inactivated in Wilms tumor. *Science.* 2007 Feb 2;315(5812):642-5. PMID: 17204608
111. Leung et al. ZMYM3 regulates BRCA1 localization at damaged chromatin to promote DNA repair. *Genes Dev.* 2017 Feb 1;31(3):260-274. PMID: 28242625
112. Liu et al. Distinct Genomic Alterations in Prostate Tumors Derived from African American Men. *Mol Cancer Res.* 2020 Dec;18(12):1815-1824. PMID: 33115829
113. Ouzzine et al. The UDP-glucuronosyltransferases of the blood-brain barrier: their role in drug metabolism and detoxication. *Front Cell Neurosci.* 2014;8:349. PMID: 25389387
114. Nagar et al. Uridine diphosphoglucuronosyltransferase pharmacogenetics and cancer. *Oncogene.* 2006 Mar 13;25(11):1659-72. PMID: 16550166
115. Allain et al. Emerging roles for UDP-glucuronosyltransferases in drug resistance and cancer progression. *Br J Cancer.* 2020 Apr;122(9):1277-1287. PMID: 32047295
116. Izumi et al. Expression of UDP-glucuronosyltransferase 1A in bladder cancer: association with prognosis and regulation by estrogen. *Mol Carcinog.* 2014 Apr;53(4):314-24. PMID: 23143693
117. Sundararaghavan et al. Glucuronidation and UGT isozymes in bladder: new targets for the treatment of uroepithelial carcinomas?. *Oncotarget.* 2017 Jan 10;8(2):3640-3648. PMID: 27690298
118. Lu et al. Drug-Metabolizing Activity, Protein and Gene Expression of UDP-Glucuronosyltransferases Are Significantly Altered in Hepatocellular Carcinoma Patients. *PLoS One.* 2015;10(5):e0127524. PMID: 26010150
119. Karas et al. *JCO Oncol Pract.* 2021 Dec 3;OP2100624. PMID: 34860573

References (continued)

120. Rocak et al. DEAD-box proteins: the driving forces behind RNA metabolism. *Nat Rev Mol Cell Biol.* 2004 Mar;5(3):232-41. PMID: 14991003
121. Fuller-Pace. The DEAD box proteins DDX5 (p68) and DDX17 (p72): multi-tasking transcriptional regulators. *Biochim Biophys Acta.* 2013 Aug;1829(8):756-63. PMID: 23523990
122. Ali. DEAD-box RNA helicases: The driving forces behind RNA metabolism at the crossroad of viral replication and antiviral innate immunity. *Virus Res.* 2021 Apr 15;296:198352. PMID: 33640359
123. Linder et al. Looking back on the birth of DEAD-box RNA helicases. *Biochim Biophys Acta.* 2013 Aug;1829(8):750-5. PMID: 23542735
124. Lin. DDX3X Multifunctionally Modulates Tumor Progression and Serves as a Prognostic Indicator to Predict Cancer Outcomes. *Int J Mol Sci.* 2019 Dec 31;21(1). PMID: 31906196
125. Song et al. The mechanism of RNA duplex recognition and unwinding by DEAD-box helicase DDX3X. *Nat Commun.* 2019 Jul 12;10(1):3085. PMID: 31300642
126. Zhou et al. Comprehensive proteomic analysis of the human spliceosome. *Nature.* 2002 Sep 12;419(6903):182-5. PMID: 12226669
127. Yedavalli et al. Requirement of DDX3 DEAD box RNA helicase for HIV-1 Rev-RRE export function. *Cell.* 2004 Oct 29;119(3):381-92. PMID: 15507209
128. Chao et al. DDX3, a DEAD box RNA helicase with tumor growth-suppressive property and transcriptional regulation activity of the p21waf1/cip1 promoter, is a candidate tumor suppressor. *Cancer Res.* 2006 Jul 1;66(13):6579-88. PMID: 16818630
129. Chuang et al. Requirement of the DEAD-Box protein ded1p for messenger RNA translation. *Science.* 1997 Mar 7;275(5305):1468-71. PMID: 9045610
130. Shih et al. Candidate tumor suppressor DDX3 RNA helicase specifically represses cap-dependent translation by acting as an eIF4E inhibitory protein. *Oncogene.* 2008 Jan 24;27(5):700-14. PMID: 17667941
131. Lee et al. Human DDX3 functions in translation and interacts with the translation initiation factor eIF3. *Nucleic Acids Res.* 2008 Aug;36(14):4708-18. PMID: 18628297