

Patient Name: 김민경
Gender: F
Sample ID: N25-251

Primary Tumor Site: Esophagus
Collection Date: 2022.07.11.

Sample Cancer Type: Esophageal Cancer

Table of Contents	Page	Report Highlights
Variant Details	2	1 Relevant Biomarkers
Biomarker Descriptions	2	0 Therapies Available
Relevant Therapy Summary	6	1 Clinical Trials

Relevant Esophageal Cancer Findings

Gene	Finding
BRAF	None detected
ERBB2	None detected
NTRK1	None detected
NTRK2	None detected
NTRK3	None detected
RET	None detected

Genomic Alteration	Finding
Tumor Mutational Burden	2.91 Mut/Mb measured

Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	FANCA p.(Q676Hfs*5) c.2026_2027dup Fanconi anemia complementation group A Allele Frequency: 46.54% Locus: chr16:89838209 Transcript: NM_000135.4	None*	None*	1

* Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO

* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Line of therapy: I: First-line therapy, II+: Other line of therapy

Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

Prevalent cancer biomarkers without relevant evidence based on included data sources

BRIP1 c.918+1G>A, CDKN2B p.(P9Afs*8) c.24_25insG, Microsatellite stable, FUBP1 p.(R344*) c.1030C>T, HLA-B deletion, NQO1 p.(P187S) c.559C>T, Tumor Mutational Burden

Variant Details

DNA Sequence Variants

Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
FANCA	p.(Q676Hfs*5)	c.2026_2027dup	.	chr16:89838209	46.54%	NM_000135.4	frameshift Insertion
BRIP1	p.(?)	c.918+1G>A	.	chr17:59885827	3.25%	NM_032043.3	unknown
CDKN2B	p.(P9Afs*8)	c.24_25insG	.	chr9:22008928	6.02%	NM_004936.4	frameshift Insertion
FUBP1	p.(R344*)	c.1030C>T	.	chr1:78429758	3.28%	NM_003902.5	nonsense
NQO1	p.(P187S)	c.559C>T	.	chr16:69745145	41.33%	NM_000903.3	missense
SETD2	p.(R1740W)	c.5218C>T	.	chr3:47129662	3.38%	NM_014159.7	missense
MSH3	p.(A57_A65del)	c.162_204delTGCAGC GGCCGCAGCGCCGC AGCGCCCCAGCGCC CCCAGCTinsCGCAGC GCCCCAGCG	.	chr5:79950708	100.00%	NM_002439.5	nonframeshift Block Substitution
HLA-B	p.([T118I;L119I])	c.353_355delCCCinsT CA	.	chr6:31324208	100.00%	NM_005514.8	missense, missense
CREBBP	p.(Q301H)	c.903G>T	.	chr16:3860676	47.01%	NM_004380.3	missense
USP9X	p.(S1253L)	c.3758C>T	.	chrX:41047318	2.94%	NM_001039590.3	missense
GCNA	p.(R486H)	c.1457G>A	.	chrX:70825570	2.54%	NM_052957.4	missense

Copy Number Variations

Gene	Locus	Copy Number	CNV Ratio
HLA-B	chr6:31322252	0	0.46

Biomarker Descriptions

FANCA p.(Q676Hfs*5) c.2026_2027dup

Fanconi anemia complementation group A

Background: The FANCA gene encodes the FA complementation group A protein, a member of the Fanconi Anemia (FA) family, which also includes FANCB, FANCC, FANCD1 (BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCI (BRIP1), FANCL, FANCM, and FANCN (PALB2)¹. FA genes are tumor suppressors that are responsible for the maintenance of replication fork stability, DNA damage repair through the removal of interstrand cross-links (ICL), and subsequent initiation of the homologous recombination repair (HRR) pathway^{43,44}. In response to DNA damage, FANCA, FANCB, FANCC, FANCE, FANCF, FANCG, FANCL, and FANCM assemble to form the FA core complex which is responsible for the monoubiquitination of the FANCI-FANCD2 (ID2) complex⁴³. Monoubiquitination of the ID2 complex promotes co-localization with BRCA1/2, which is critical in BRCA mediated DNA repair^{45,46}. Loss of function mutations in the FA family and HRR pathway, including FANCA, can result in the BRCAness phenotype, characterized by a defect in the HRR pathway, mimicking BRCA1 or BRCA2 loss^{13,47}. Germline mutations in FA genes lead to Fanconi Anemia, a condition characterized by chromosomal instability and congenital abnormalities, including bone marrow failure and cancer predisposition^{48,49}. Of those diagnosed with FA, mutations in FANCA are the most common and confer predisposition to myelodysplastic syndrome, acute myeloid leukemia, and solid tumors^{44,49,50,51,52}.

Alterations and prevalence: Somatic mutations in FANCA are observed in 4-8% of uterine, colorectal, and bladder cancers and about 6% of melanoma⁸. Biallelic loss is also reported in 2-5% of uveal melanoma, invasive breast carcinoma, ovarian cancer, and prostate cancer⁸.

Biomarker Descriptions (continued)

Potential relevance: The PARP inhibitor, talazoparib⁵³ in combination with enzalutamide is approved (2023) for metastatic castration-resistant prostate cancer (mCRPC) with mutations in HRR genes that includes FANCA. Consistent with other genes that contribute to the BRCAness phenotype, mutations in FANCA are shown to confer enhanced sensitivity in vitro to DNA damaging agents, including cisplatin, as well as PARP inhibitors such as olaparib^{54,55}. FANCA copy number loss along with reduced expression has also been associated with genetic instability in sporadic acute myeloid leukemia (AML)⁵².

BRIP1 c.918+1G>A

BRCA1 interacting protein C-terminal helicase 1

Background: The BRIP1 gene encodes the BRCA1 interacting protein C-terminal helicase 1 and is a member of the RecQ DEAH helicase family that plays a role in homologous recombination repair (HRR) of double-stranded breaks (DSBs) in DNA¹⁰. BRIP1 interacts directly with BRCA1 through the BRCT domain and controls BRCA1-dependent DNA repair and the DNA damage-induced G2-M checkpoint control¹¹. BRIP1 is a tumor suppressor gene. Loss of function mutations in BRIP1 are implicated in the BRCAness phenotype, characterized by a defect in HRR, mimicking BRCA1 or BRCA2 loss^{12,13}. Germline aberrations in BRIP1 are associated with inherited disorders such as Fanconi anemia (FA)¹⁴. Specifically, BRIP1 was shown to be biallelically inactivated in FA patients and is also considered a high-risk gene for familial late-onset ovarian cancer^{14,15}. BRIP1 germline mutations confer ~ 10% cumulative risk of ovarian cancer and are associated with an increased risk of colorectal cancer^{10,16}.

Alterations and prevalence: Somatic mutations in BRIP1 are observed in up to 8% of uterine corpus endometrial carcinoma, 5% of skin cutaneous melanoma, and 4% of bladder urothelial carcinoma^{8,9}.

Potential relevance: The PARP inhibitor, olaparib¹⁷ is approved (2020) for metastatic castration-resistant prostate cancer (mCRPC) with deleterious or suspected deleterious, germline or somatic mutations in HRR genes that includes BRIP1. Consistent with other genes associated with the BRCAness phenotype, BRIP1 mutations may aid in selecting patients likely to respond to PARP inhibitors or platinum therapy^{12,18}. In 2022, the FDA granted fast track designation to the small molecule inhibitor, pidnarulex¹⁹, for BRCA1/2, PALB2, or other homologous recombination deficiency (HRD) mutations in breast and ovarian cancers.

CDKN2B p.(P9Afs*8) c.24_25insG

cyclin dependent kinase inhibitor 2B

Background: CDKN2B encodes cyclin dependent kinase inhibitor 2B, a cell cycle regulator that controls G1/S progression^{1,56}. CDKN2B, also known as p15/INK4B, belongs to a family of INK4 cyclin-dependent kinase inhibitors, which also includes CDKN2A (p16/INK4A), CDKN2C (p18/INK4C), and CDKN2D (p19/INK4D)⁵⁶. The INK4 family regulates cell cycle progression by inhibiting CDK4 or CDK6, thereby preventing the phosphorylation of Rb^{57,58,59}. CDKN2B is a tumor suppressor and aberrations in this gene commonly co-occur with CDKN2A⁵⁶. Germline mutations in CDKN2B are linked to pancreatic cancer predisposition and familial renal cell carcinoma^{1,60,61}.

Alterations and prevalence: CDKN2B copy number loss is a frequently occurring somatic aberration that is observed in 55% of glioblastoma multiforme, 43% of mesothelioma, 35% of esophageal adenocarcinoma, 31% of bladder urothelial carcinoma, 29% of skin cutaneous melanoma, 28% of head and neck squamous cell carcinoma, 27% of pancreatic adenocarcinoma, 26% of lung squamous cell carcinoma, 25% of diffuse large B-cell lymphoma, 16% of lung adenocarcinoma, 15% of sarcoma, 14% of cholangiocarcinoma, 11% of stomach adenocarcinoma and brain lower grade glioma, 5% of liver hepatocellular carcinoma, 4% of adrenocortical carcinoma, breast invasive carcinoma, thymoma, and kidney renal papillary cell carcinoma, 3% of kidney renal clear cell carcinoma and ovarian serous cystadenocarcinoma, and 2% of uterine carcinosarcoma and kidney chromophobe^{8,9}. Somatic mutations in CDKN2B are observed in 2% of uterine carcinosarcoma^{8,9}. CDKN2B copy number loss is also observed in pediatric cancers, including 64% of childhood T-lymphoblastic leukemia/lymphoma, 37% of pediatric B-lymphoblastic leukemia/lymphoma, 25% of pediatric gliomas, 14% of pediatric bone cancers, 6% of embryonal tumors, and 2% of peripheral nervous system cancers^{8,9}. Somatic mutations in CDKN2B are observed in less than 1% of bone cancer (1 in 327 cases)^{8,9}.

Potential relevance: Currently, no therapies are approved for CDKN2B aberrations. Homozygous deletion of CDKN2B is a molecular marker used in staging grade 4 pediatric IDH-mutant astrocytoma⁶².

Microsatellite stable

Background: Microsatellites are short tandem repeats (STR) of 1 to 6 bases of DNA between 5 to 50 repeat units in length. There are approximately 0.5 million STRs that occupy 3% of the human genome²⁰. Microsatellite instability (MSI) is defined as a change in the length of a microsatellite in a tumor as compared to normal tissue^{21,22}. MSI is closely tied to the status of the mismatch repair (MMR) genes. In humans, the core MMR genes include MLH1, MSH2, MSH6, and PMS2²³. Mutations and loss of expression in MMR genes, known as defective MMR (dMMR), lead to MSI. In contrast, when MMR genes lack alterations, they are referred to as MMR proficient (pMMR). Consensus criteria were first described in 1998 and defined MSI-high (MSI-H) as instability in two or more of the following five markers: BAT25, BAT26, D5S346, D2S123, and D17S250²⁴. Tumors with instability in one of the five markers were defined as MSI-low (MSI-L) whereas, those with instability in zero markers were defined as MS-stable (MSS)²⁴. Tumors classified as MSI-L are

Biomarker Descriptions (continued)

often phenotypically indistinguishable from MSS tumors and tend to be grouped with MSS^{25,26,27,28,29}. MSI-H is a hallmark of Lynch syndrome (LS), also known as hereditary non-polyposis colorectal cancer, which is caused by germline mutations in the MMR genes²². LS is associated with an increased risk of developing colorectal cancer, as well as other cancers, including endometrial and stomach cancer^{21,22,26,30}.

Alterations and prevalence: The MSI-H phenotype is observed in 30% of uterine corpus endometrial carcinoma, 20% of stomach adenocarcinoma, 15-20% of colon adenocarcinoma, and 5-10% of rectal adenocarcinoma^{21,22,31,32}. MSI-H is also observed in 5% of adrenal cortical carcinoma and at lower frequencies in other cancers such as esophageal, liver, and ovarian cancers^{31,32}.

Potential relevance: Anti-PD-1 immune checkpoint inhibitors including pembrolizumab³³ (2014) and nivolumab³⁴ (2015) are approved for patients with MSI-H or dMMR colorectal cancer who have progressed following chemotherapy. Pembrolizumab³³ is also approved as a single agent, for the treatment of patients with advanced endometrial carcinoma that is MSI-H or dMMR with disease progression on prior therapy who are not candidates for surgery or radiation. Importantly, pembrolizumab is approved for the treatment of MSI-H or dMMR solid tumors that have progressed following treatment, with no alternative option and is the first anti-PD-1 inhibitor to be approved with a tumor agnostic indication³³. Dostarlimab³⁵ (2021) is also approved for dMMR recurrent or advanced endometrial carcinoma or solid tumors that have progressed on prior treatment and is recommended as a subsequent therapy option in dMMR/MSI-H advanced or metastatic colon or rectal cancer^{27,36}. The cytotoxic T-lymphocyte antigen 4 (CTLA-4) blocking antibody, ipilimumab³⁷ (2011), is approved alone or in combination with nivolumab in MSI-H or dMMR colorectal cancer that has progressed following treatment with chemotherapy. MSI-H may confer a favorable prognosis in colorectal cancer although outcomes vary depending on stage and tumor location^{27,38,39}. Specifically, MSI-H is a strong prognostic indicator of better overall survival (OS) and relapse free survival (RFS) in stage II as compared to stage III colorectal cancer patients³⁹. The majority of patients with tumors classified as either MSS or pMMR do not benefit from treatment with single-agent immune checkpoint inhibitors as compared to those with MSI-H tumors^{40,41}. However, checkpoint blockade with the addition of chemotherapy or targeted therapies have demonstrated response in MSS or pMMR cancers^{40,41}.

FUBP1 p.(R344*) c.1030C>T

far upstream element binding protein 1

Background: The FUBP1 gene encodes the far upstream element binding protein 1, a DNA/RNA binding protein implicated in a variety of cellular functions^{1,42}. Specifically, FUBP1 is observed to bind single-stranded DNA (ssDNA) and RNA resulting in the regulation of transcription, translation, and splicing⁴². FUBP1 activates the transcription of targets including the oncogene MYC which functions in cell cycle regulation, metabolism, and apoptosis⁴². FUBP1 is also observed to repress the transcription of targets including the tumor suppressors CDKN1A, CDKN2B, and CDKN1B, which function in cell cycle regulation⁴².

Alterations and prevalence: Somatic mutations in FUBP1 are observed in 9% of brain lower grade glioma, 6% of uterine corpus endometrial carcinoma, 4% of skin cutaneous melanoma, and 3% of colorectal adenocarcinoma^{8,9}. Mutations typically result in inactivation of FUBP1 through alteration of splicing sites, introduction of stop codons, or out-of-frame insertions or deletions⁴². Biallelic loss of FUBP1 is observed in 3% of pheochromocytoma and paraganglioma^{8,9}. Co-deletion of 1p and 19q is frequently observed in oligodendrogliomas, which results in the monoallelic loss of FUBP1 and CIC on 19q⁴².

Potential relevance: Currently, no therapies are approved for FUBP1 aberrations.

HLA-B deletion

major histocompatibility complex, class I, B

Background: The HLA-B gene encodes the major histocompatibility complex, class I, B¹. MHC (major histocompatibility complex) class I molecules are located on the cell surface of nucleated cells and present antigens from within the cell for recognition by cytotoxic T cells². MHC class I molecules are heterodimers composed of two polypeptide chains, α and B2M³. The classical MHC class I genes include HLA-A, HLA-B, and HLA-C and encode the α polypeptide chains, which present short polypeptide chains, of 7 to 11 amino acids, to the immune system to distinguish self from non-self^{4,5,6}. Downregulation of MHC class I promotes tumor evasion of the immune system, suggesting a tumor suppressor role for HLA-B⁷.

Alterations and prevalence: Somatic mutations in HLA-B are observed in 10% of diffuse large B-cell lymphoma (DLBCL), 5% of cervical squamous cell carcinoma and stomach adenocarcinoma, 4% of head and neck squamous cell carcinoma and colorectal adenocarcinoma, 3% of uterine cancer, and 2% of esophageal adenocarcinoma and skin cutaneous melanoma^{8,9}. Biallelic loss of HLA-B is observed in 5% of DLBCL^{8,9}.

Potential relevance: Currently, no therapies are approved for HLA-B aberrations.

Genes Assayed

Genes Assayed for the Detection of DNA Sequence Variants

ABL1, ABL2, ACVR1, AKT1, AKT2, AKT3, ALK, AR, ARAF, ATP1A1, AURKA, AURKB, AURKC, AXL, BCL2, BCL2L12, BCL6, BCR, BMP5, BRAF, BTK, CACNA1D, CARD11, CBL, CCND1, CCND2, CCND3, CCNE1, CD79B, CDK4, CDK6, CHD4, CSF1R, CTNNB1, CUL1, CYSLTR2, DDR2, DGCR8, DROSHA, E2F1, EGFR, EIF1AX, EPAS1, ERBB2, ERBB3, ERBB4, ESR1, EZH2, FAM135B, FGF7, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FOXL2, FOXO1, GATA2, GLI1, GNA11, GNAQ, GNAS, HIF1A, HRAS, IDH1, IDH2, IKBKB, IL6ST, IL7R, IRF4, IRS4, KCNJ5, KDR, KIT, KLF4, KLF5, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAPK1, MAX, MDM4, MECOM, MED12, MEF2B, MET, MITF, MPL, MTOR, MYC, MYCN, MYD88, MYO1D, NFE2L2, NRAS, NSD2, NT5C2, NTRK1, NTRK2, NTRK3, NUP93, PAX5, PCBP1, PDGFRA, PDGFRB, PIK3C2B, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PIK3R2, PIM1, PLCG1, PPP2R1A, PPP6C, PRKACA, PTPN11, PTPRD, PXDNL, RAC1, RAF1, RARA, RET, RGS7, RHEB, RHOA, RICTOR, RIT1, ROS1, RPL10, SETBP1, SF3B1, SIX1, SIX2, SLC01B3, SMC1A, SMO, SNCAIP, SOS1, SOX2, SPOP, SRC, SRSF2, STAT3, STAT5B, STAT6, TAF1, TERT, TGFB1, TOP1, TOP2A, TPMT, TRRAP, TSHR, U2AF1, USP8, WAS, XPO1, ZNF217, ZNF429

Genes Assayed for the Detection of Copy Number Variations

ABCB1, ABL1, ABL2, ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AKT1, AKT2, AKT3, ALK, AMER1, APC, AR, ARAF, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AURKA, AURKC, AXIN1, AXIN2, AXL, B2M, BAP1, BARD1, BCL2, BCL2L12, BCL6, BCOR, BLM, BMPR2, BRAF, BRCA1, BRCA2, BRIP1, CARD11, CASP8, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDK4, CDK6, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHD4, CHEK1, CHEK2, CIC, CREBBP, CSMD3, CTCF, CTLA4, CTNND2, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, DAXX, DDR1, DDR2, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, EGFR, EIF1AX, ELF3, EMSY, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERBB2, ERBB3, ERBB4, ERCC2, ERCC4, ERFF1, ESR1, ETV6, EZH2, FAM135B, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAT1, FBXW7, FGF19, FGF23, FGF3, FGF4, FGF9, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FUBP1, FYN, GATA2, GATA3, GLI3, GNA13, GNAS, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, IDH2, IGF1R, IKBKB, IL7R, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KDR, KEAP1, KIT, KLF5, KMT2A, KMT2B, KMT2C, KMT2D, KRAS, LARP4B, LATS1, LATS2, MAGOH, MAP2K1, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK1, MAPK8, MAX, MCL1, MDM2, MDM4, MECOM, MEF2B, MEN1, MET, MGA, MITF, MLH1, MLH3, MPL, MRE11, MSH2, MSH3, MSH6, MTAP, MTOR, MUTYH, MYC, MYCL, MYCN, MYD88, NBN, NCOR1, NF1, NF2, NFE2L2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, NRAS, NTRK1, NTRK3, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PCBP1, PDCD1, PDCD1LG2, PDGFRA, PDGFRB, PDIA3, PGD, PHF6, PIK3C2B, PIK3CA, PIK3CB, PIK3R1, PIK3R2, PIM1, PLCG1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R1A, PPP2R2A, PPP6C, PRDM1, PRDM9, PRKACA, PRKAR1A, PTCH1, PTEN, PTPN11, PTPRT, PXDNL, RAC1, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RAF1, RARA, RASA1, RASA2, RB1, RBM10, RECQL4, RET, RHEB, RICTOR, RIT1, RNASEH2A, RNASEH2B, RNF43, ROS1, RPA1, RPS6KB1, RPTOR, RUNX1, SDHA, SDHB, SDHD, SETBP1, SETD2, SF3B1, SLC01B3, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SMC1A, SMO, SOX9, SPEN, SPOP, SRC, STAG2, STAT3, STAT6, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TERT, TET2, TGFB2, TNFAIP3, TNFRSF14, TOP1, TP53, TP63, TPMT, TPP2, TSC1, TSC2, U2AF1, USP8, USP9X, VHL, WT1, XPO1, XRCC2, XRCC3, YAP1, YES1, ZFH3, ZMYM3, ZNF217, ZNF429, ZRSR2

Genes Assayed for the Detection of Fusions

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSP02, RSP03, TERT

Genes Assayed with Full Exon Coverage

ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AMER1, APC, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AXIN1, AXIN2, B2M, BAP1, BARD1, BCOR, BLM, BMPR2, BRCA1, BRCA2, BRIP1, CALR, CASP8, CBFB, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHEK1, CHEK2, CIC, CIITA, CREBBP, CSMD3, CTCF, CTLA4, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, CYP2D6, DAXX, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, ELF3, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERCC2, ERCC4, ERCC5, ERFF1, ETV6, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAS, FAT1, FBXW7, FUBP1, GATA3, GNA13, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, ID3, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KEAP1, KLHL13, KMT2A, KMT2B, KMT2C, KMT2D, LARP4B, LATS1, LATS2, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK8, MEN1, MGA, MLH1, MLH3, MRE11, MSH2, MSH3, MSH6, MTAP, MTUS2, MUTYH, NBN, NCOR1, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PDCD1, PDCD1LG2, PDIA3, PGD, PHF6, PIK3R1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R2A, PRDM1, PRDM9, PRKAR1A, PSMB10, PSMB8, PSMB9, PTCH1, PTEN, PTPRT, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RASA1, RASA2, RB1, RBM10, RECQL4, RNASEH2A, RNASEH2B, RNASEH2C, RNF43, RPA1, RPL22, RPL5, RUNX1, RUNX1T1, SDHA, SDHB, SDHC, SDHD, SETD2, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SOCS1, SOX9, SPEN, STAG2, STAT1, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TET2, TGFB2, TMEM132D, TNFAIP3, TNFRSF14, TP53, TP63, TPP2, TSC1, TSC2, UGT1A1, USP9X, VHL, WT1, XRCC2, XRCC3, ZBTB20, ZFH3, ZMYM3, ZRSR2

Relevant Therapy Summary

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

FANCA p.(Q676Hfs*5) c.2026_2027dup

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
talazoparib	×	×	×	×	<div></div> (II)

* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

HRR Details

Gene/Genomic Alteration	Finding
LOH percentage	0.0%
Not Detected	Not Applicable

Homologous recombination repair (HRR) genes were defined from published evidence in relevant therapies, clinical guidelines, as well as clinical trials, and include - BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, and RAD54L.

Thermo Fisher Scientific's Ion Torrent Oncomine Reporter software was used in generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on Oncomine Reporter (6.1.1 data version 2025.06(006)). The data presented here are from a curated knowledge base of publicly available information, but may not be exhaustive. FDA information was sourced from www.fda.gov and is current as of 2025-05-14. NCCN information was sourced from www.nccn.org and is current as of 2025-05-01. EMA information was sourced from www.ema.europa.eu and is current as of 2025-05-14. ESMO information was sourced from www.esmo.org and is current as of 2025-05-01. Clinical Trials information is current as of 2025-05-01. For the most up-to-date information regarding a particular trial, search www.clinicaltrials.gov by NCT ID or search local clinical trials authority website by local identifier listed in 'Other identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials listed in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Hulpke et al. The MHC I loading complex: a multitasking machinery in adaptive immunity. *Trends Biochem Sci.* PMID: 23849087
3. Adams et al. The adaptable major histocompatibility complex (MHC) fold: structure and function of nonclassical and MHC class I-like molecules. *Annu Rev Immunol.* 2013;31:529-61. PMID: 23298204
4. Rossjohn et al. T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol.* 2015;33:169-200. PMID: 25493333
5. Parham. MHC class I molecules and KIRs in human history, health and survival. *Nat Rev Immunol.* 2005 Mar;5(3):201-14. PMID: 15719024
6. Sidney et al. HLA class I supertypes: a revised and updated classification. *BMC Immunol.* 2008 Jan 22;9:1. PMID: 18211710
7. Cornel et al. MHC Class I Downregulation in Cancer: Underlying Mechanisms and Potential Targets for Cancer Immunotherapy. *Cancers (Basel).* 2020 Jul 2;12(7). PMID: 32630675
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
10. Ali et al. BRIP-1 germline mutation and its role in colon cancer: presentation of two case reports and review of literature. *BMC Med Genet.* 2019 May 7;20(1):75. PMID: 31064327
11. Daino et al. Loss of the BRCA1-interacting helicase BRIP1 results in abnormal mammary acinar morphogenesis. *PLoS ONE.* 2013;8(9):e74013. PMID: 24040146
12. Lim et al. Evaluation of the methods to identify patients who may benefit from PARP inhibitor use. *Endocr. Relat. Cancer.* 2016 Jun;23(6):R267-85. PMID: 27226207
13. Lord et al. BRCAness revisited. *Nat. Rev. Cancer.* 2016 Feb;16(2):110-20. PMID: 26775620
14. Taniguchi et al. Molecular pathogenesis of Fanconi anemia: recent progress. *Blood.* 2006 Jun 1;107(11):4223-33. PMID: 16493006
15. Weber-Lassalle et al. BRIP1 loss-of-function mutations confer high risk for familial ovarian cancer, but not familial breast cancer. *Breast Cancer Res.* 2018 Jan 24;20(1):7. PMID: 29368626
16. Ramus et al. Germline Mutations in the BRIP1, BARD1, PALB2, and NBN Genes in Women With Ovarian Cancer. *J. Natl. Cancer Inst.* 2015 Nov;107(11). PMID: 26315354
17. https://www.accessdata.fda.gov/drugsatfda_docs/label/2023/208558s028lbl.pdf
18. Pennington et al. Germline and somatic mutations in homologous recombination genes predict platinum response and survival in ovarian, fallopian tube, and peritoneal carcinomas. *Clin. Cancer Res.* 2014 Feb 1;20(3):764-75. PMID: 24240112
19. <https://www.senhwabio.com/en/news/20220125>
20. Lander et al. Initial sequencing and analysis of the human genome. *Nature.* 2001 Feb 15;409(6822):860-921. PMID: 11237011
21. Baudrin et al. Molecular and Computational Methods for the Detection of Microsatellite Instability in Cancer. *Front Oncol.* 2018 Dec 12;8:621. doi: 10.3389/fonc.2018.00621. eCollection 2018. PMID: 30631754
22. Nojadeh et al. Microsatellite instability in colorectal cancer. *EXCLI J.* 2018;17:159-168. PMID: 29743854
23. Saeed et al. Microsatellites in Pursuit of Microbial Genome Evolution. *Front Microbiol.* 2016 Jan 5;6:1462. doi: 10.3389/fmicb.2015.01462. eCollection 2015. PMID: 26779133
24. Boland et al. A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res.* 1998 Nov 15;58(22):5248-57. PMID: 9823339
25. Halford et al. Low-level microsatellite instability occurs in most colorectal cancers and is a nonrandomly distributed quantitative trait. *Cancer Res.* 2002 Jan 1;62(1):53-7. PMID: 11782358
26. Imai et al. Carcinogenesis and microsatellite instability: the interrelationship between genetics and epigenetics. *Carcinogenesis.* 2008 Apr;29(4):673-80. PMID: 17942460
27. NCCN Guidelines® - NCCN-Colon Cancer [Version 3.2025]
28. Pawlik et al. Colorectal carcinogenesis: MSI-H versus MSI-L. *Dis. Markers.* 2004;20(4-5):199-206. PMID: 15528785
29. Lee et al. Low-Level Microsatellite Instability as a Potential Prognostic Factor in Sporadic Colorectal Cancer. *Medicine (Baltimore).* 2015 Dec;94(50):e2260. PMID: 26683947
30. Latham et al. Microsatellite Instability Is Associated With the Presence of Lynch Syndrome Pan-Cancer. *J. Clin. Oncol.* 2019 Feb 1;37(4):286-295. PMID: 30376427

References (continued)

31. Cortes-Ciriano et al. A molecular portrait of microsatellite instability across multiple cancers. *Nat Commun.* 2017 Jun 6;8:15180. doi: 10.1038/ncomms15180. PMID: 28585546
32. Bonneville et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precis Oncol.* 2017;2017. PMID: 29850653
33. https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125514s174lbl.pdf
34. https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125554s129lbl.pdf
35. https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/761174s009lbl.pdf
36. NCCN Guidelines® - NCCN-Rectal Cancer [Version 2.2025]
37. https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125377s133lbl.pdf
38. Ribic et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N. Engl. J. Med.* 2003 Jul 17;349(3):247-57. PMID: 12867608
39. Klingbiel et al. Prognosis of stage II and III colon cancer treated with adjuvant 5-fluorouracil or FOLFIRI in relation to microsatellite status: results of the PETACC-3 trial. *Ann. Oncol.* 2015 Jan;26(1):126-32. PMID: 25361982
40. Hermel et al. The Emerging Role of Checkpoint Inhibition in Microsatellite Stable Colorectal Cancer. *J Pers Med.* 2019 Jan 16;9(1). PMID: 30654522
41. Ciardiello et al. Immunotherapy of colorectal cancer: Challenges for therapeutic efficacy. *Cancer Treat. Rev.* 2019 Jun;76:22-32. PMID: 31079031
42. Debaize et al. The master regulator FUBP1: its emerging role in normal cell function and malignant development. *Cell Mol Life Sci.* 2019 Jan;76(2):259-281. PMID: 30343319
43. Niraj et al. The Fanconi Anemia Pathway in Cancer. *Annu Rev Cancer Biol.* 2019 Mar;3:457-478. PMID: 30882047
44. Rodríguez et al. Fanconi anemia pathway. *Curr Biol.* 2017 Sep 25;27(18):R986-R988. PMID: 28950089
45. Garcia-Higuera et al. Interaction of the Fanconi anemia proteins and BRCA1 in a common pathway. *Mol. Cell.* 2001 Feb;7(2):249-62. PMID: 11239454
46. Hussain et al. Direct interaction of FANCD2 with BRCA2 in DNA damage response pathways. *Hum. Mol. Genet.* 2004 Jun 15;13(12):1241-8. PMID: 15115758
47. Byrum et al. Defining and Modulating 'BRCAness'. *Trends Cell Biol.* 2019 Sep;29(9):740-751. PMID: 31362850
48. Michl et al. Interplay between Fanconi anemia and homologous recombination pathways in genome integrity. *EMBO J.* 2016 May 2;35(9):909-23. PMID: 27037238
49. Abbasi et al. A rare FANCA gene variation as a breast cancer susceptibility allele in an Iranian population. *Mol Med Rep.* 2017 Jun;15(6):3983-3988. PMID: 28440412
50. Levran et al. Sequence variation in the Fanconi anemia gene FAA. *Proc. Natl. Acad. Sci. U.S.A.* 1997 Nov 25;94(24):13051-6. PMID: 9371798
51. Antonio et al. A comprehensive strategy for the subtyping of patients with Fanconi anaemia: conclusions from the Spanish Fanconi Anemia Research Network. *J. Med. Genet.* 2007 Apr;44(4):241-9. PMID: 17105750
52. Tischkowitz et al. Deletion and reduced expression of the Fanconi anemia FANCA gene in sporadic acute myeloid leukemia. *Leukemia.* 2004 Mar;18(3):420-5. PMID: 14749703
53. https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/217439s000lbl.pdf
54. McCabe et al. Deficiency in the repair of DNA damage by homologous recombination and sensitivity to poly(ADP-ribose) polymerase inhibition. *Cancer Res.* 2006 Aug 15;66(16):8109-15. PMID: 16912188
55. Wilkes et al. A germline FANCA alteration that is associated with increased sensitivity to DNA damaging agents. *Cold Spring Harb Mol Case Stud.* 2017 Sep;3(5). PMID: 28864460
56. Xia et al. Dominant role of CDKN2B/p15INK4B of 9p21.3 tumor suppressor hub in inhibition of cell-cycle and glycolysis. *Nat Commun.* 2021 Apr 6;12(1):2047. PMID: 33824349
57. Scruggs et al. Loss of CDKN2B Promotes Fibrosis via Increased Fibroblast Differentiation Rather Than Proliferation. *Am. J. Respir. Cell Mol. Biol.* 2018 Aug;59(2):200-214. PMID: 29420051
58. Roussel. The INK4 family of cell cycle inhibitors in cancer. *Oncogene.* 1999 Sep 20;18(38):5311-7. PMID: 10498883
59. Aytac et al. Rb independent inhibition of cell growth by p15(INK4B). *Biochem. Biophys. Res. Commun.* 1999 Aug 27;262(2):534-8. PMID: 10462509
60. Jafri et al. Germline Mutations in the CDKN2B Tumor Suppressor Gene Predispose to Renal Cell Carcinoma. *Cancer Discov.* 2015 Jul;5(7):723-9. PMID: 25873077

References (continued)

61. Tu et al. CDKN2B deletion is essential for pancreatic cancer development instead of unmeaningful co-deletion due to juxtaposition to CDKN2A. *Oncogene*. 2018 Jan 4;37(1):128-138. PMID: 28892048
62. Louis et al. cIMPACT-NOW update 6: new entity and diagnostic principle recommendations of the cIMPACT-Utrecht meeting on future CNS tumor classification and grading. *Brain Pathol*. 2020 Jul;30(4):844-856. PMID: 32307792