

**Patient Name:** 하영자  
**Gender:** F  
**Sample ID:** N25-286

**Primary Tumor Site:** Lung  
**Collection Date:** 2025.10.22

## Sample Cancer Type: Lung Cancer

Table of Contents	Page
Variant Details	2
Biomarker Descriptions	2
Alert Details	6
Relevant Therapy Summary	8

**Report Highlights**  
2 Relevant Biomarkers  
18 Therapies Available  
201 Clinical Trials

## Relevant Lung Cancer Findings

Gene	Finding	Gene	Finding
ALK	None detected	NTRK1	None detected
BRAF	None detected	NTRK2	None detected
EGFR	<b>EGFR p.(L858R) c.2573T&gt;G</b>	NTRK3	None detected
ERBB2	None detected	RET	None detected
KRAS	None detected	ROS1	None detected
MET	None detected		

Genomic Alteration	Finding
Tumor Mutational Burden	<b>3.85 Mut/Mb measured</b>

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>EGFR p.(L858R) c.2573T&gt;G</b> epidermal growth factor receptor Allele Frequency: 20.21% Locus: chr7:55259515 Transcript: NM_005228.5	<b>afatinib</b> <sup>1, 2 / I, II+</sup> <b>amivantamab + lazertinib</b> <sup>1, 2 / I, II+</sup> <b>bevacizumab† + erlotinib</b> <sup>2 / I, II+</sup> <b>dacomitinib</b> <sup>1, 2 / I, II+</sup> <b>erlotinib</b> <sup>2 / I, II+</sup> <b>erlotinib + ramucirumab</b> <sup>1, 2 / I, II+</sup> <b>gefitinib</b> <sup>1, 2 / I, II+</sup> <b>osimertinib</b> <sup>1, 2 / I, II+</sup> <b>osimertinib + chemotherapy</b> <sup>1, 2 / I</sup> <b>amivantamab + chemotherapy</b> <sup>1, 2 / II+</sup> <b>datopotamab deruxtecan-dlnk</b> <sup>1 / II+</sup> <b>BAT1706 + erlotinib</b> <sup>2</sup> gefitinib + chemotherapy <sup>I</sup> atezolizumab + bevacizumab + chemotherapy <sup>II+</sup>	None*	201

\* Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

\* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

† Includes biosimilars/generics

Line of therapy: I: First-line therapy, II+: Other line of therapy

Tier Reference: Li et al. *Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists.* J Mol Diagn. 2017 Jan;19(1):4-23.

## Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	TP53 p.(G266R) c.796G>A	None*	None*	6
	tumor protein p53			
	Allele Frequency: 19.70%			
	Locus: chr17:7577142			
	Transcript: NM_000546.6			

\* Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO  
\* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO  
† Includes biosimilars/generics  
Line of therapy: I: First-line therapy, II+: Other line of therapy  
Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

 Alerts informed by public data sources:  Contraindicated,  Resistance,  Breakthrough,  Fast Track

EGFR p.(L858R) c.2573T>G  izarontamab brengitecan<sup>1</sup>, patritumab deruxtecan<sup>1</sup>  
 DB-1310<sup>1</sup>

Public data sources included in alerts: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources  
Microsatellite stable, HLA-A p.(L180\*) c.539T>A, Tumor Mutational Burden

## Variant Details

DNA Sequence Variants							
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
EGFR	p.(L858R)	c.2573T>G	COSM6224	chr7:55259515	20.21%	NM_005228.5	missense
TP53	p.(G266R)	c.796G>A	COSM10794	chr17:7577142	19.70%	NM_000546.6	missense
HLA-A	p.(L180*)	c.539T>A	.	chr6:29911240	100.00%	NM_001242758.1	nonsense
KMT2D	p.(V1062L)	c.3184G>C	.	chr12:49444187	40.17%	NM_003482.4	missense
SOX9	p.(A118V)	c.353C>T	.	chr17:70117885	7.05%	NM_000346.4	missense

## Biomarker Descriptions

### EGFR p.(L858R) c.2573T>G

epidermal growth factor receptor

Background: The EGFR gene encodes the epidermal growth factor receptor (EGFR), a member of the ERBB/human epidermal growth factor receptor (HER) tyrosine kinase family<sup>1</sup>. In addition to EGFR/ERBB1/HER1, other members of the ERBB/HER family include ERBB2/HER2, ERBB3/HER3, and ERBB4/HER4<sup>55</sup>. EGFR ligand-induced dimerization results in kinase activation and leads to stimulation of oncogenic signaling pathways, including the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways<sup>56</sup>. Activation of these pathways promotes cell proliferation, differentiation, and survival<sup>57,58</sup>.

Alterations and prevalence: Recurrent somatic mutations in the tyrosine kinase domain (TKD) of EGFR are observed in approximately 10-20% of lung adenocarcinoma, and at higher frequencies in never-smoker, female, and Asian populations<sup>8,9,59,60</sup>. The most common mutations occur near the ATP-binding pocket of the TKD and include short in-frame deletions in exon 19 (EGFR exon 19 deletion) and the L858R amino acid substitution in exon 21<sup>61</sup>. These mutations constitutively activate EGFR resulting in downstream signaling, and represent 80% of the EGFR mutations observed in lung cancer<sup>61</sup>. A second group of less prevalent activating mutations includes E709K, G719X, S768I, L861Q, and short in-frame insertion mutations in exon 20<sup>62,63,64,65</sup>. EGFR activating mutations in lung cancer tend to be mutually exclusive to KRAS activating mutations<sup>66</sup>. In contrast, a different set of recurrent activating EGFR mutations in

## Biomarker Descriptions (continued)

the extracellular domain includes R108K, A289V and G598V and are primarily observed in glioblastoma<sup>61,67</sup>. Amplification of EGFR is observed in several cancer types including 44% of glioblastoma multiforme, 12% of esophageal adenocarcinoma, 10% of head and neck squamous cell carcinoma, 8% of brain lower grade glioma, 6% of lung squamous cell carcinoma, 5% of bladder urothelial carcinoma, lung adenocarcinoma, and stomach adenocarcinoma, 3% of cholangiocarcinoma, and 2% of cervical squamous cell carcinoma, sarcoma, and breast invasive carcinoma<sup>8,9,15,60,67</sup>. Deletion of exons 2-7, encoding the extracellular domain of EGFR (EGFRvIII), results in overexpression of a ligand-independent constitutively active protein and is observed in approximately 30% of glioblastoma<sup>68,69,70</sup>. Alterations in EGFR are rare in pediatric cancers<sup>8,9</sup>. Somatic mutations are observed in 2% of bone cancer and glioma, 1% of leukemia (4 in 354 cases), and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 252 cases), peripheral nervous system cancers (1 in 1158 cases), and embryonal tumors (3 in 332 cases)<sup>8,9</sup>. Amplification of EGFR is observed in 2% of bone cancer and less than 1% of Wilms tumor (1 in 136 cases), B-lymphoblastic leukemia/lymphoma (2 in 731 cases), and leukemia (1 in 250 cases)<sup>8,9</sup>.

**Potential relevance:** Approved first-generation EGFR tyrosine kinase inhibitors (TKIs) include erlotinib<sup>71</sup> (2004) and gefitinib<sup>72</sup> (2015), which block the activation of downstream signaling by reversible interaction with the ATP-binding site. Although initially approved for advanced lung cancer, the discovery that drug sensitivity was associated with exon 19 and exon 21 activating mutations allowed first-generation TKIs to become subsequently approved for front-line therapy in lung cancer tumors containing exon 19 or exon 21 activating mutations<sup>73</sup>. Second-generation TKIs afatinib<sup>74</sup> (2013) and dacomitinib<sup>75</sup> (2018) bind EGFR and other ERBB/HER gene family members irreversibly and were subsequently approved. First- and second-generation TKIs afatinib, dacomitinib, erlotinib, and gefitinib are recommended for the treatment NSCLC harboring EGFR exon 19 insertions, exon 19 deletions, point mutations L861Q, L858R, S768I, and codon 719 mutations, whereas most EGFR exon 20 insertions, except p.A763\_Y764insFQEA, confer resistance to the same therapies<sup>76,77,78,79</sup>. In 2025, the FDA approved the irreversible EGFR inhibitor, sunvozertinib<sup>80</sup>, for the treatment of locally advanced or metastatic non-small cell lung cancer in adult patients with EGFR exon 20 insertion mutations whose disease has progressed on or after platinum-based chemotherapy. In 2022, the FDA granted breakthrough therapy designation to the irreversible EGFR inhibitor, CLN-081 (TPC-064)<sup>81</sup> for locally advanced or metastatic non-small cell lung cancer harboring EGFR exon 20 insertion mutations. In lung cancer containing EGFR exon 19 or 21 activating mutations, treatment with TKIs is eventually associated with the emergence of drug resistance<sup>82</sup>. The primary resistance mutation that emerges following treatment with first-generation TKI is T790M, accounting for 50-60% of resistant cases<sup>61</sup>. Third generation TKIs were developed to maintain sensitivity in the presence of T790M<sup>82</sup>. Osimertinib<sup>83</sup> (2015) is an irreversible inhibitor indicated for metastatic EGFR T790M positive lung cancer and for the first-line treatment of metastatic NSCLC containing EGFR exon 19 deletions or exon 21 L858R mutations. Like first-generation TKIs, treatment with osimertinib is associated with acquired resistance, specifically the C797S mutation, which occurs in 22-44% of cases<sup>82</sup>. The T790M and C797S mutations may be each selected following sequential treatment with a first-generation TKI followed by a third-generation TKI or vice versa<sup>84</sup>. T790M and C797S can occur in either cis or trans allelic orientation<sup>84</sup>. If C797S is observed following progression after treatment with a third-generation TKI in the first-line setting, sensitivity may be retained to first-generation TKIs<sup>84</sup>. If C797S co-occurs in trans with T790M following sequential treatment with first- and third-generation TKIs, patients may exhibit sensitivity to combination first- and third-generation TKIs, but resistance to third-generation TKIs alone<sup>84,85</sup>. However, C797S occurring in cis conformation with T790M, confers resistance to first- and third-generation TKIs<sup>84</sup>. Fourth-generation TKIs are in development to overcome acquired resistance mutations after osimertinib treatment, including BDTX-1535<sup>86</sup> (2024), a CNS-penetrating small molecule inhibitor, that received fast track designation from the FDA for the treatment of patients with EGFR C797S-positive NSCLC who have disease progression on or after a third-generation EGFR TKI. EGFR-targeting antibodies including cetuximab (2004), panitumumab (2006), and necitumumab (2016) are under investigation in combination with EGFR-targeting TKIs for efficacy against EGFR mutations<sup>87</sup>. The bispecific antibody, amivantamab<sup>88</sup> (2021), targeting EGFR and MET was approved for NSCLC tumors harboring EGFR exon 20 insertion mutations. A small molecule kinase inhibitor, lazertinib<sup>89</sup> (2024), was approved in combination with amivantamab as a first-line treatment for adult patients with locally advanced or metastatic NSCLC with EGFR exon 19 deletions or exon 21 L858R mutations. HLX-42<sup>90</sup>, an anti-EGFR-antibody-drug conjugate (ADC) consisting of an anti-EGFR monoclonal antibody conjugated with a novel high potency DNA topoisomerase I (topo I) inhibitor, also received fast track designation (2024) for the treatment of patients with advanced or metastatic EGFR-mutated non-small cell lung cancer whose disease has progressed on a third-generation EGFR tyrosine kinase inhibitor. CPO301<sup>91</sup> (2023) received a fast track designation from the FDA for the treatment of EGFR mutations in patients with metastatic NSCLC who are relapsed/refractory or ineligible for EGFR targeting therapy such as 3rd-generation EGFR inhibitors, including osimertinib. The Oncoprex immunogene therapy quaratusugene ozeplasmid<sup>92</sup> (2020), in combination with osimertinib, received fast track designation from the FDA for NSCLC tumors harboring EGFR mutations that progressed on osimertinib alone. Amplification and mutations of EGFR commonly occur in H3-wild type IDH-wild type diffuse pediatric high-grade glioma<sup>93,94,95</sup>.

### TP53 p.(G266R) c.796G>A

#### *tumor protein p53*

**Background:** The TP53 gene encodes the tumor suppressor protein p53, which binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair<sup>1</sup>. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis<sup>10</sup>. Alterations in TP53 are required for oncogenesis as they result in loss of protein function and gain of transforming potential<sup>11</sup>. Germline mutations in TP53 are

## Biomarker Descriptions (continued)

the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers<sup>12,13</sup>.

**Alterations and prevalence:** TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)<sup>8,9,14,15,16,17</sup>. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common, including substitutions at codons R158, R175, Y220, R248, R273, and R282<sup>8,9</sup>. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes<sup>18,19,20,21</sup>. Alterations in TP53 are also observed in pediatric cancers<sup>8,9</sup>. Somatic mutations are observed in 53% of non-Hodgkin lymphoma, 24% of soft tissue sarcoma, 19% of glioma, 13% of bone cancer, 9% of B-lymphoblastic leukemia/lymphoma, 4% of embryonal tumors, 3% of Wilms tumor and leukemia, 2% of T-lymphoblastic leukemia/lymphoma, and less than 1% of peripheral nervous system cancers (5 in 1158 cases)<sup>8,9</sup>. Biallelic loss of TP53 is observed in 10% of bone cancer, 2% of Wilms tumor, and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 731 cases) and leukemia (1 in 250 cases)<sup>8,9</sup>.

**Potential relevance:** The small molecule p53 reactivator, PC14586<sup>22</sup> (2020), received a fast track designation by the FDA for advanced tumors harboring a TP53 Y220C mutation. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation<sup>23,24</sup>. TP53 mutation is a diagnostic marker of SHH-activated, TP53-mutant medulloblastoma<sup>25</sup>. TP53 mutations confer poor prognosis and poor risk in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), and chronic lymphocytic leukemia (CLL), and acute lymphoblastic leukemia (ALL)<sup>26,27,28,29,30</sup>. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant<sup>31</sup>. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system<sup>32</sup>.

### Microsatellite stable

**Background:** Microsatellites are short tandem repeats (STR) of 1 to 6 bases of DNA between 5 to 50 repeat units in length. There are approximately 0.5 million STRs that occupy 3% of the human genome<sup>33</sup>. Microsatellite instability (MSI) is defined as a change in the length of a microsatellite in a tumor as compared to normal tissue<sup>34,35</sup>. MSI is closely tied to the status of the mismatch repair (MMR) genes. In humans, the core MMR genes include MLH1, MSH2, MSH6, and PMS2<sup>36</sup>. Mutations and loss of expression in MMR genes, known as defective MMR (dMMR), lead to MSI. In contrast, when MMR genes lack alterations, they are referred to as MMR proficient (pMMR). Consensus criteria were first described in 1998 and defined MSI-high (MSI-H) as instability in two or more of the following five markers: BAT25, BAT26, D5S346, D2S123, and D17S250<sup>37</sup>. Tumors with instability in one of the five markers were defined as MSI-low (MSI-L) whereas, those with instability in zero markers were defined as MS-stable (MSS)<sup>37</sup>. Tumors classified as MSI-L are often phenotypically indistinguishable from MSS tumors and tend to be grouped with MSS<sup>38,39,40,41,42</sup>. MSI-H is a hallmark of Lynch syndrome (LS), also known as hereditary non-polyposis colorectal cancer, which is caused by germline mutations in the MMR genes<sup>35</sup>. LS is associated with an increased risk of developing colorectal cancer, as well as other cancers, including endometrial and stomach cancer<sup>34,35,39,43</sup>.

**Alterations and prevalence:** The MSI-H phenotype is observed in 30% of uterine corpus endothelial carcinoma, 20% of stomach adenocarcinoma, 15-20% of colon adenocarcinoma, and 5-10% of rectal adenocarcinoma<sup>34,35,44,45</sup>. MSI-H is also observed in 5% of adrenal cortical carcinoma and at lower frequencies in other cancers such as esophageal, liver, and ovarian cancers<sup>44,45</sup>.

**Potential relevance:** Anti-PD-1 immune checkpoint inhibitors including pembrolizumab<sup>46</sup> (2014) and nivolumab<sup>47</sup> (2015) are approved for patients with MSI-H or dMMR colorectal cancer who have progressed following chemotherapy. Pembrolizumab<sup>46</sup> is also approved as a single agent, for the treatment of patients with advanced endometrial carcinoma that is MSI-H or dMMR with disease progression on prior therapy who are not candidates for surgery or radiation. Importantly, pembrolizumab is approved for the treatment of MSI-H or dMMR solid tumors that have progressed following treatment, with no alternative option and is the first anti-PD-1 inhibitor to be approved with a tumor agnostic indication<sup>46</sup>. Dostarlimab<sup>48</sup> (2021) is also approved for dMMR recurrent or advanced endometrial carcinoma or solid tumors that have progressed on prior treatment and is recommended as a subsequent therapy option in dMMR/MSI-H advanced or metastatic colon or rectal cancer<sup>40,49</sup>. The cytotoxic T-lymphocyte antigen 4 (CTLA-4) blocking antibody, ipilimumab<sup>50</sup> (2011), is approved alone or in combination with nivolumab in MSI-H or dMMR colorectal cancer that has progressed following treatment with chemotherapy. MSI-H may confer a favorable prognosis in colorectal cancer although outcomes vary depending on stage and tumor location<sup>40,51,52</sup>. Specifically, MSI-H is a strong prognostic indicator of better overall survival (OS) and relapse free survival (RFS) in stage II as compared to stage III colorectal cancer patients<sup>52</sup>. The majority of patients with tumors classified as either MSS or pMMR do not benefit from treatment with single-agent immune checkpoint inhibitors as compared to those with MSI-H tumors<sup>53,54</sup>. However, checkpoint blockade with the addition of chemotherapy or targeted therapies have demonstrated response in MSS or pMMR cancers<sup>53,54</sup>.

## Biomarker Descriptions (continued)

### HLA-A p.(L180\*) c.539T>A

*major histocompatibility complex, class I, A*

**Background:** The HLA-A gene encodes the major histocompatibility complex, class I, A<sup>1</sup>. MHC (major histocompatibility complex) class I molecules are located on the cell surface of nucleated cells and present antigens from within the cell for recognition by cytotoxic T cells<sup>2</sup>. MHC class I molecules are heterodimers composed of two polypeptide chains,  $\alpha$  and B2M<sup>3</sup>. The classical MHC class I genes include HLA-A, HLA-B, and HLA-C and encode the  $\alpha$  polypeptide chains, which present short polypeptide chains, of 7 to 11 amino acids, to the immune system to distinguish self from non-self<sup>4,5,6</sup>. Downregulation of MHC class I promotes tumor evasion of the immune system, suggesting a tumor suppressor role for HLA-A<sup>7</sup>.

**Alterations and prevalence:** Somatic mutations in HLA-A are observed in 7% of diffuse large B-cell lymphoma (DLBCL), 4% of cervical squamous cell carcinoma and head and neck squamous cell carcinoma, 3% of colorectal adenocarcinoma, and 2% of uterine corpus endometrial carcinoma and stomach adenocarcinoma<sup>8,9</sup>. Biallelic loss of HLA-A is observed in 4% of DLBCL<sup>8,9</sup>.

**Potential relevance:** Currently, no therapies are approved for HLA-A aberrations.

## Alerts Informed By Public Data Sources

### Current FDA Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

FDA information is current as of 2025-09-17. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

### EGFR p.(L858R) c.2573T>G

#### izationaltamab brengitecan

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR L858R mutation

**Supporting Statement:**

The FDA has granted Breakthrough designation to EGFR/HER3 targeting bispecific antibody-drug conjugate (ADC), izationaltamab brengitecan, for the treatment of patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) harboring EGFR exon 19 deletions or exon 21 L858R substitution mutations who experienced disease progression on or after treatment with an EGFR TKI and platinum-based chemotherapy.

**Reference:**

<https://www.onclive.com/view/fda-grants-breakthrough-therapy-designation-to-izationaltamab-brengitecan-in-egfr-nsclc>

#### patritumab deruxtecan

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR L858R mutation or EGFRi sensitizing mutation

**Supporting Statement:**

The FDA has granted Breakthrough Therapy designation to a potential first-in-class HER3 directed antibody-drug conjugate, patritumab deruxtecan, for metastatic or locally advanced, EGFR-mutant non-small cell lung cancer.

**Reference:**

<https://www.cancernetwork.com/view/fda-grants-breakthrough-therapy-status-to-patritumab-deruxtecan-for-egfr-metastatic-nsclc>

#### DB-1310

**Cancer type:** Non-Small Cell Lung Cancer

**Variant class:** EGFR L858R mutation

**Supporting Statement:**

The FDA has granted Fast Track designation to the HER3-targeting antibody-drug conjugate, DB-1310, for the treatment of adult patients with advanced, unresectable or metastatic non-squamous non-small cell lung cancer with EGFR exon 19 deletion or L858R mutation and who have progressed after treatment with a third-generation EGFR tyrosine kinase inhibitor and platinum-based chemotherapy.

**Reference:**

<https://www.targetedonc.com/view/novel-her3-adc-receives-fda-fast-track-for-refractory-nsclc>

## Genes Assayed

### Genes Assayed for the Detection of DNA Sequence Variants

ABL1, ABL2, ACVR1, AKT1, AKT2, AKT3, ALK, AR, ARAF, ATP1A1, AURKA, AURKB, AURKC, AXL, BCL2, BCL2L12, BCL6, BCR, BMP5, BRAF, BTK, CACNA1D, CARD11, CBL, CCND1, CCND2, CCND3, CCNE1, CD79B, CDK4, CDK6, CHD4, CSF1R, CTNNB1, CUL1, CYSLTR2, DDR2, DGCR8, DROSHA, E2F1, EGFR, EIF1AX, EPAS1, ERBB2, ERBB3, ERBB4, ESR1, EZH2, FAM135B, FGF7, FGFR1, FGFR2, FGFR3,

## Genes Assayed (continued)

### Genes Assayed for the Detection of DNA Sequence Variants (continued)

FGFR4, FLT3, FLT4, FOXA1, FOXL2, FOXO1, GATA2, GLI1, GNA11, GNAQ, GNAS, HIF1A, HRAS, IDH1, IDH2, IKBKB, IL6ST, IL7R, IRF4, IRS4, KCNJ5, KDR, KIT, KLF4, KLF5, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAPK1, MAX, MDM4, MECOM, MED12, MEF2B, MET, MITF, MPL, MTOR, MYC, MYCN, MYD88, MYO1, NFE2L2, NRAS, NSD2, NT5C2, NTRK1, NTRK2, NTRK3, NUP93, PAX5, PCBP1, PDGFRA, PDGFRB, PIK3C2B, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PIK3R2, PIM1, PLCG1, PPP2R1A, PPP6C, PRKACA, PTPN11, PTPRD, PXDNL, RAC1, RAF1, RARA, RET, RGS7, RHEB, RHOA, RICTOR, RIT1, ROS1, RPL10, SETBP1, SF3B1, SIX1, SIX2, SLC01B3, SMC1A, SMO, SNCAIP, SOS1, SOX2, SPOP, SRC, SRSF2, STAT3, STAT5B, STAT6, TAF1, TERT, TGFB1, TOP1, TOP2A, TPMT, TRRAP, TSHR, U2AF1, USP8, WAS, XPO1, ZNF217, ZNF429

### Genes Assayed for the Detection of Copy Number Variations

ABCB1, ABL1, ABL2, ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AKT1, AKT2, AKT3, ALK, AMER1, APC, AR, ARAF, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AURKA, AURKC, AXIN1, AXIN2, AXL, B2M, BAP1, BARD1, BCL2, BCL2L12, BCL6, BCOR, BLM, BMPR2, BRAF, BRCA1, BRCA2, BRIP1, CARD11, CASP8, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDK4, CDK6, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHD4, CHEK1, CHEK2, CIC, CREBBP, CSMD3, CTCF, CTLA4, CTNND2, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, DAXX, DDR1, DDR2, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, EGFR, EIF1AX, ELF3, EMSY, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERBB2, ERBB3, ERBB4, ERCC2, ERCC4, ERFF1, ESR1, ETV6, EZH2, FAM135B, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAT1, FBXW7, FGF19, FGF23, FGF3, FGF4, FGF9, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FUBP1, FYN, GATA2, GATA3, GLI3, GNA13, GNAS, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, IDH2, IGF1R, IKBKB, IL7R, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KDR, KEAP1, KIT, KLF5, KMT2A, KMT2B, KMT2C, KMT2D, KRAS, LARP4B, LATS1, LATS2, MAGOH, MAP2K1, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK1, MAPK8, MAX, MCL1, MDM2, MDM4, MECOM, MEF2B, MEN1, MET, MGA, MITF, MLH1, MLH3, MPL, MRE11, MSH2, MSH3, MSH6, MTAP, MTOR, MUTYH, MYC, MYCL, MYCN, MYD88, NBN, NCOR1, NF1, NF2, NFE2L2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, NRAS, NTRK1, NTRK3, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PCBP1, PDCD1, PDCD1LG2, PDGFRA, PDGFRB, PDIA3, PGD, PHF6, PIK3C2B, PIK3CA, PIK3CB, PIK3R1, PIK3R2, PIM1, PLCG1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R1A, PPP2R2A, PPP6C, PRDM1, PRDM9, PRKACA, PRKAR1A, PTCH1, PTEN, PTPN11, PTPRT, PXDNL, RAC1, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RAF1, RARA, RASA1, RASA2, RB1, RBM10, RECQL4, RET, RHEB, RICTOR, RIT1, RNASEH2A, RNASEH2B, RNF43, ROS1, RPA1, RPS6KB1, RPTOR, RUNX1, SDHA, SDHB, SDHD, SETBP1, SETD2, SF3B1, SLC01B3, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SMC1A, SMO, SOX9, SPEN, SPOP, SRC, STAG2, STAT3, STAT6, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TERT, TET2, TGFB2, TNFAIP3, TNFRSF14, TOP1, TP53, TP63, TPMT, TPP2, TSC1, TSC2, U2AF1, USP8, USP9X, VHL, WT1, XPO1, XRCC2, XRCC3, YAP1, YES1, ZFH3, ZMYM3, ZNF217, ZNF429, ZRSR2

### Genes Assayed for the Detection of Fusions

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, REL, RET, ROS1, RSPO2, RSPO3, TERT

### Genes Assayed with Full Exon Coverage

ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AMER1, APC, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AXIN1, AXIN2, B2M, BAP1, BARD1, BCOR, BLM, BMPR2, BRCA1, BRCA2, BRIP1, CALR, CASP8, CBFB, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHEK1, CHEK2, CIC, CIITA, CREBBP, CSMD3, CTCF, CTLA4, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, CYP2D6, DAXX, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, ELF3, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERCC2, ERCC4, ERCC5, ERFF1, ETV6, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAS, FAT1, FBXW7, FUBP1, GATA3, GNA13, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, ID3, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KEAP1, KLHL13, KMT2A, KMT2B, KMT2C, KMT2D, LARP4B, LATS1, LATS2, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK8, MEN1, MGA, MLH1, MLH3, MRE11, MSH2, MSH3, MSH6, MTAP, MTUS2, MUTYH, NBN, NCOR1, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PDCD1, PDCD1LG2, PDIA3, PGD, PHF6, PIK3R1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R2A, PRDM1, PRDM9, PRKAR1A, PSMB10, PSMB8, PSMB9, PTCH1, PTEN, PTPRT, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RASA1, RASA2, RB1, RBM10, RECQL4, RNASEH2A, RNASEH2B, RNASEH2C, RNF43, RPA1, RPL22, RPL5, RUNX1, RUNX1T1, SDHA, SDHB, SDHC, SDHD, SETD2, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SOCS1, SOX9, SPEN, STAG2, STAT1, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TET2, TGFB2, TMEM132D, TNFAIP3, TNFRSF14, TP53, TP63, TPP2, TSC1, TSC2, UGT1A1, USP9X, VHL, WT1, XRCC2, XRCC3, ZBTB20, ZFH3, ZMYM3, ZRSR2

## Relevant Therapy Summary

● In this cancer type

○ In other cancer type

● In this cancer type and other cancer types

✕ No evidence

### EGFR p.(L858R) c.2573T>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
osimertinib	●	●	●	●	● (III)
afatinib	●	●	●	●	● (II)
dacomitinib	●	●	●	●	● (II)
gefitinib	●	●	●	●	● (II)
erlotinib + ramucirumab	●	●	●	●	✕
amivantamab + carboplatin + pemetrexed	●	●	●	✕	✕
amivantamab + lazertinib	●	●	●	✕	✕
datopotamab deruxtecan-dlnk	●	●	✕	✕	✕
osimertinib + chemotherapy + pemetrexed	●	✕	●	✕	✕
bevacizumab + erlotinib	✕	●	●	●	✕
erlotinib	✕	●	●	●	✕
osimertinib + carboplatin + pemetrexed	✕	●	✕	✕	✕
osimertinib + cisplatin + pemetrexed	✕	●	✕	✕	✕
BAT1706 + erlotinib	✕	✕	●	✕	✕
bevacizumab (Allergan) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Biocon) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Celltrion) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Mabxience) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Pfizer) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Samsung Bioepis) + erlotinib	✕	✕	●	✕	✕
bevacizumab (Stada) + erlotinib	✕	✕	●	✕	✕
atezolizumab + bevacizumab + carboplatin + paclitaxel	✕	✕	✕	●	✕
gefitinib + carboplatin + pemetrexed	✕	✕	✕	●	✕
adebrelimab, bevacizumab, chemotherapy	✕	✕	✕	✕	● (IV)
afatinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (IV)
befotertinib	✕	✕	✕	✕	● (IV)
bevacizumab, almonertinib, chemotherapy	✕	✕	✕	✕	● (IV)
catequentinib, toripalimab	✕	✕	✕	✕	● (IV)
EGFR tyrosine kinase inhibitor	✕	✕	✕	✕	● (IV)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

 In this cancer type    
  In other cancer type    
  In this cancer type and other cancer types    
  No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
furmonertinib, chemotherapy	×	×	×	×	● (IV)
gefitinib, chemotherapy	×	×	×	×	● (IV)
gefitinib, endostatin	×	×	×	×	● (IV)
natural product, gefitinib, erlotinib, icotinib hydrochloride, osimertinib, almonertinib, furmonertinib	×	×	×	×	● (IV)
almonertinib, apatinib	×	×	×	×	● (III)
almonertinib, catequentinib	×	×	×	×	● (III)
almonertinib, chemotherapy	×	×	×	×	● (III)
almonertinib, radiation therapy	×	×	×	×	● (III)
befotertinib, icotinib hydrochloride	×	×	×	×	● (III)
bevacizumab, osimertinib	×	×	×	×	● (III)
CK-101, gefitinib	×	×	×	×	● (III)
datopotamab deruxtecan-dlnk, osimertinib	×	×	×	×	● (III)
furmonertinib	×	×	×	×	● (III)
furmonertinib, osimertinib, chemotherapy	×	×	×	×	● (III)
gefitinib, afatinib, erlotinib, metformin hydrochloride	×	×	×	×	● (III)
glumetinib, osimertinib	×	×	×	×	● (III)
icotinib hydrochloride, catequentinib	×	×	×	×	● (III)
icotinib hydrochloride, chemotherapy	×	×	×	×	● (III)
icotinib hydrochloride, radiation therapy	×	×	×	×	● (III)
izalontamab brengitecan	×	×	×	×	● (III)
izalontamab brengitecan, osimertinib	×	×	×	×	● (III)
JMT-101, osimertinib	×	×	×	×	● (III)
osimertinib, bevacizumab	×	×	×	×	● (III)
osimertinib, chemotherapy	×	×	×	×	● (III)
osimertinib, datopotamab deruxtecan-dlnk	×	×	×	×	● (III)
sacituzumab tirumotecan	×	×	×	×	● (III)
sacituzumab tirumotecan, osimertinib	×	×	×	×	● (III)
savolitinib, osimertinib	×	×	×	×	● (III)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

 In this cancer type    
  In other cancer type    
  In this cancer type and other cancer types    
  No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
SH-1028	×	×	×	×	● (III)
TY-9591, osimertinib	×	×	×	×	● (III)
PM-1080, almonertinib	×	×	×	×	● (II/III)
SCTB-14, chemotherapy	×	×	×	×	● (II/III)
ABSK-043, furmonertinib	×	×	×	×	● (II)
afatinib, chemotherapy	×	×	×	×	● (II)
almonertinib	×	×	×	×	● (II)
almonertinib, adabrelimab, chemotherapy	×	×	×	×	● (II)
almonertinib, bevacizumab	×	×	×	×	● (II)
almonertinib, chemoradiation therapy	×	×	×	×	● (II)
almonertinib, dacomitinib	×	×	×	×	● (II)
amivantamab, chemotherapy	×	×	×	×	● (II)
amivantamab, lazertinib, chemotherapy	×	×	×	×	● (II)
atezolizumab, bevacizumab, tiragolumab	×	×	×	×	● (II)
befotertinib, bevacizumab, chemotherapy	×	×	×	×	● (II)
bevacizumab, afatinib	×	×	×	×	● (II)
bevacizumab, furmonertinib	×	×	×	×	● (II)
cadonilimab, chemotherapy, catequentinib	×	×	×	×	● (II)
camrelizumab, apatinib	×	×	×	×	● (II)
capmatinib, osimertinib, ramucirumab	×	×	×	×	● (II)
catequentinib, almonertinib	×	×	×	×	● (II)
catequentinib, chemotherapy	×	×	×	×	● (II)
chemotherapy, atezolizumab, bevacizumab	×	×	×	×	● (II)
dacomitinib, osimertinib	×	×	×	×	● (II)
EGFR tyrosine kinase inhibitor, osimertinib, chemotherapy	×	×	×	×	● (II)
EGFR tyrosine kinase inhibitor, radiation therapy	×	×	×	×	● (II)
erlotinib, chemotherapy	×	×	×	×	● (II)
erlotinib, OBI-833	×	×	×	×	● (II)
furmonertinib, bevacizumab	×	×	×	×	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
furmonertinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
furmonertinib, catequentinib	✕	✕	✕	✕	● (II)
furmonertinib, chemotherapy, bevacizumab	✕	✕	✕	✕	● (II)
furmonertinib, icotinib hydrochloride	✕	✕	✕	✕	● (II)
gefitinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
gefitinib, icotinib hydrochloride	✕	✕	✕	✕	● (II)
gefitinib, thalidomide	✕	✕	✕	✕	● (II)
icotinib hydrochloride	✕	✕	✕	✕	● (II)
icotinib hydrochloride, autologous RAK cell	✕	✕	✕	✕	● (II)
icotinib hydrochloride, osimertinib	✕	✕	✕	✕	● (II)
ivonescimab, chemotherapy	✕	✕	✕	✕	● (II)
izalontamab brengitecan, almonertinib	✕	✕	✕	✕	● (II)
JS-207, chemotherapy	✕	✕	✕	✕	● (II)
lazertinib	✕	✕	✕	✕	● (II)
lazertinib, bevacizumab	✕	✕	✕	✕	● (II)
lazertinib, chemotherapy	✕	✕	✕	✕	● (II)
osimertinib, bevacizumab, chemotherapy	✕	✕	✕	✕	● (II)
osimertinib, radiation therapy	✕	✕	✕	✕	● (II)
PLB-1004, bozitinib, osimertinib	✕	✕	✕	✕	● (II)
ramucirumab, erlotinib	✕	✕	✕	✕	● (II)
sunvozertinib	✕	✕	✕	✕	● (II)
sunvozertinib, catequentinib	✕	✕	✕	✕	● (II)
sunvozertinib, golidocitinib	✕	✕	✕	✕	● (II)
tislelizumab, chemotherapy, bevacizumab	✕	✕	✕	✕	● (II)
toripalimab	✕	✕	✕	✕	● (II)
toripalimab, bevacizumab, Clostridium butyricum, chemotherapy	✕	✕	✕	✕	● (II)
toripalimab, chemotherapy	✕	✕	✕	✕	● (II)
TY-9591, chemotherapy	✕	✕	✕	✕	● (II)
vabametkib, lazertinib	✕	✕	✕	✕	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

## Relevant Therapy Summary (continued)

● In this cancer type    
 ○ In other cancer type    
 ● In this cancer type and other cancer types    
 ✕ No evidence

### EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
YL-202	✕	✕	✕	✕	● (II)
zorifertinib, pirotinib	✕	✕	✕	✕	● (II)
AP-L1898	✕	✕	✕	✕	● (I/II)
BH-30643	✕	✕	✕	✕	● (I/II)
bozitinib, osimertinib	✕	✕	✕	✕	● (I/II)
BPI-361175	✕	✕	✕	✕	● (I/II)
chemotherapy, DZD-6008	✕	✕	✕	✕	● (I/II)
dacomitinib, catequentinib	✕	✕	✕	✕	● (I/II)
DAJH-1050766	✕	✕	✕	✕	● (I/II)
DB-1310, osimertinib	✕	✕	✕	✕	● (I/II)
dositinib	✕	✕	✕	✕	● (I/II)
FWD-1509	✕	✕	✕	✕	● (I/II)
H-002	✕	✕	✕	✕	● (I/II)
ifebemtiniib, furmonertinib	✕	✕	✕	✕	● (I/II)
MRTX0902	✕	✕	✕	✕	● (I/II)
necitumumab, osimertinib	✕	✕	✕	✕	● (I/II)
quaratusugene ozeplasmid, osimertinib	✕	✕	✕	✕	● (I/II)
RC-108, furmonertinib, toripalimab	✕	✕	✕	✕	● (I/II)
sotiburafusp alfa, chemotherapy	✕	✕	✕	✕	● (I/II)
sotiburafusp alfa, HB-0030	✕	✕	✕	✕	● (I/II)
sunvozertinib, chemotherapy	✕	✕	✕	✕	● (I/II)
TRX-221	✕	✕	✕	✕	● (I/II)
WSD-0922	✕	✕	✕	✕	● (I/II)
almonertinib, midazolam	✕	✕	✕	✕	● (I)
ASKC-202	✕	✕	✕	✕	● (I)
AZD-9592	✕	✕	✕	✕	● (I)
BG-60366	✕	✕	✕	✕	● (I)
BPI-1178, osimertinib	✕	✕	✕	✕	● (I)
catequentinib, gefitinib, metformin hydrochloride	✕	✕	✕	✕	● (I)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

Relevant Therapy Summary (continued)

In this cancer type

In other cancer type

In this cancer type and other cancer types

No evidence

EGFR p.(L858R) c.2573T>G (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
DZD-6008	×	×	×	×	● (I)
EGFR tyrosine kinase inhibitor, catequentinib	×	×	×	×	● (I)
genolimzumab, fruquintinib	×	×	×	×	● (I)
IBI-318, lenvatinib	×	×	×	×	● (I)
KQB-198, osimertinib	×	×	×	×	● (I)
LAVA-1223	×	×	×	×	● (I)
MRX-2843, osimertinib	×	×	×	×	● (I)
osimertinib, carotuximab	×	×	×	×	● (I)
osimertinib, Minnelide	×	×	×	×	● (I)
osimertinib, tegatrabetan	×	×	×	×	● (I)
patritumab deruxtecan	×	×	×	×	● (I)
repotrectinib, osimertinib	×	×	×	×	● (I)
VIC-1911, osimertinib	×	×	×	×	● (I)
WTS-004	×	×	×	×	● (I)
YH-013	×	×	×	×	● (I)
zipalertinib, chemotherapy, glumetinib, pimitespib, quemliclustat	×	×	×	×	● (I)

TP53 p.(G266R) c.796G>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
almonertinib, catequentinib	×	×	×	×	● (III)
osimertinib, chemotherapy	×	×	×	×	● (III)
osimertinib, bevacizumab, chemotherapy	×	×	×	×	● (II)
sunvozertinib, catequentinib	×	×	×	×	● (II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.

HRR Details

Gene/Genomic Alteration	Finding
Not Detected	Not Applicable

Homologous recombination repair (HRR) genes were defined from published evidence in relevant therapies, clinical guidelines, as well as clinical trials, and include - BRCA1, BRCA2, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, and RAD54L.

Thermo Fisher Scientific's Ion Torrent OncoPrint Reporter software was used in generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on OncoPrint Reporter (6.1.1 data version 2025.10(006)). The data presented here are from a curated knowledge base of publicly available information, but may not be exhaustive. FDA information was sourced from [www.fda.gov](http://www.fda.gov) and is current as of 2025-09-17. NCCN information was sourced from [www.nccn.org](http://www.nccn.org) and is current as of 2025-09-02. EMA information was sourced from [www.ema.europa.eu](http://www.ema.europa.eu) and is current as of 2025-09-17. ESMO information was sourced from [www.esmo.org](http://www.esmo.org) and is current as of 2025-09-02. Clinical Trials information is current as of 2025-09-02. For the most up-to-date information regarding a particular trial, search [www.clinicaltrials.gov](http://www.clinicaltrials.gov) by NCT ID or search local clinical trials authority website by local identifier listed in 'Other identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials listed in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.

## References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Hulpke et al. The MHC I loading complex: a multitasking machinery in adaptive immunity. *Trends Biochem Sci.* PMID: 23849087
3. Adams et al. The adaptable major histocompatibility complex (MHC) fold: structure and function of nonclassical and MHC class I-like molecules. *Annu Rev Immunol.* 2013;31:529-61. PMID: 23298204
4. Rossjohn et al. T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol.* 2015;33:169-200. PMID: 25493333
5. Parham. MHC class I molecules and KIRs in human history, health and survival. *Nat Rev Immunol.* 2005 Mar;5(3):201-14. PMID: 15719024
6. Sidney et al. HLA class I supertypes: a revised and updated classification. *BMC Immunol.* 2008 Jan 22;9:1. PMID: 18211710
7. Cornel et al. MHC Class I Downregulation in Cancer: Underlying Mechanisms and Potential Targets for Cancer Immunotherapy. *Cancers (Basel).* 2020 Jul 2;12(7). PMID: 32630675
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
10. Nag et al. The MDM2-p53 pathway revisited. *J Biomed Res.* 2013 Jul;27(4):254-71. PMID: 23885265
11. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell.* 2014 Mar 17;25(3):304-17. PMID: 24651012
12. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol.* 2010 Jan;2(1):a001008. PMID: 20182602
13. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med.* 2017 Apr 3;7(4). PMID: 28270529
14. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature.* 2012 Sep 27;489(7417):519-25. PMID: 22960745
15. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature.* 2015 Jan 29;517(7536):576-82. PMID: 25631445
16. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet.* 2016 Jun;48(6):607-16. PMID: 27158780
17. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature.* 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061
18. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat.* 2002 Jun;19(6):607-14. PMID: 12007217
19. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer.* 2011 Apr;2(4):466-74. PMID: 21779514
20. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene.* 2007 Apr 2;26(15):2157-65. PMID: 17401424
21. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat.* 2014 Jun;35(6):766-78. PMID: 24729566
22. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
23. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol.* 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534
24. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci.* 2017 Nov;74(22):4171-4187. PMID: 28643165
25. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol.* 2021 Aug 2;23(8):1231-1251. PMID: 34185076
26. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
27. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 2.2025]
28. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2025]
29. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 3.2025]

## References (continued)

30. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 2.2025]
31. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 3.2025]
32. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med.* 2020 Aug 3. PMID: 32747829
33. Lander et al. Initial sequencing and analysis of the human genome. *Nature.* 2001 Feb 15;409(6822):860-921. PMID: 11237011
34. Baudrin et al. Molecular and Computational Methods for the Detection of Microsatellite Instability in Cancer. *Front Oncol.* 2018 Dec 12;8:621. doi: 10.3389/fonc.2018.00621. eCollection 2018. PMID: 30631754
35. Nojadeh et al. Microsatellite instability in colorectal cancer. *EXCLI J.* 2018;17:159-168. PMID: 29743854
36. Saeed et al. Microsatellites in Pursuit of Microbial Genome Evolution. *Front Microbiol.* 2016 Jan 5;6:1462. doi: 10.3389/fmicb.2015.01462. eCollection 2015. PMID: 26779133
37. Boland et al. A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res.* 1998 Nov 15;58(22):5248-57. PMID: 9823339
38. Halford et al. Low-level microsatellite instability occurs in most colorectal cancers and is a nonrandomly distributed quantitative trait. *Cancer Res.* 2002 Jan 1;62(1):53-7. PMID: 11782358
39. Imai et al. Carcinogenesis and microsatellite instability: the interrelationship between genetics and epigenetics. *Carcinogenesis.* 2008 Apr;29(4):673-80. PMID: 17942460
40. NCCN Guidelines® - NCCN-Colon Cancer [Version 4.2025]
41. Pawlik et al. Colorectal carcinogenesis: MSI-H versus MSI-L. *Dis. Markers.* 2004;20(4-5):199-206. PMID: 15528785
42. Lee et al. Low-Level Microsatellite Instability as a Potential Prognostic Factor in Sporadic Colorectal Cancer. *Medicine (Baltimore).* 2015 Dec;94(50):e2260. PMID: 26683947
43. Latham et al. Microsatellite Instability Is Associated With the Presence of Lynch Syndrome Pan-Cancer. *J. Clin. Oncol.* 2019 Feb 1;37(4):286-295. PMID: 30376427
44. Cortes-Ciriano et al. A molecular portrait of microsatellite instability across multiple cancers. *Nat Commun.* 2017 Jun 6;8:15180. doi: 10.1038/ncomms15180. PMID: 28585546
45. Bonneville et al. Landscape of Microsatellite Instability Across 39 Cancer Types. *JCO Precis Oncol.* 2017;2017. PMID: 29850653
46. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125514s178lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125514s178lbl.pdf)
47. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125554s131lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125554s131lbl.pdf)
48. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/761174s009lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/761174s009lbl.pdf)
49. NCCN Guidelines® - NCCN-Rectal Cancer [Version 3.2025]
50. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125377s136lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125377s136lbl.pdf)
51. Ribic et al. Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based adjuvant chemotherapy for colon cancer. *N. Engl. J. Med.* 2003 Jul 17;349(3):247-57. PMID: 12867608
52. Klingbiel et al. Prognosis of stage II and III colon cancer treated with adjuvant 5-fluorouracil or FOLFIRI in relation to microsatellite status: results of the PETACC-3 trial. *Ann. Oncol.* 2015 Jan;26(1):126-32. PMID: 25361982
53. Hermel et al. The Emerging Role of Checkpoint Inhibition in Microsatellite Stable Colorectal Cancer. *J Pers Med.* 2019 Jan 16;9(1). PMID: 30654522
54. Ciardiello et al. Immunotherapy of colorectal cancer: Challenges for therapeutic efficacy. *Cancer Treat. Rev.* 2019 Jun;76:22-32. PMID: 31079031
55. King et al. Amplification of a novel v-erbB-related gene in a human mammary carcinoma. *Science.* 1985 Sep 6;229(4717):974-6. PMID: 2992089
56. Liu et al. EGFR-TKIs resistance via EGFR-independent signaling pathways. *Mol Cancer.* 2018 Feb 19;17(1):53. PMID: 29455669
57. Zhixiang. ErbB Receptors and Cancer. *Methods Mol. Biol.* 2017;1652:3-35. PMID: 28791631
58. Gutierrez et al. HER2: biology, detection, and clinical implications. *Arch. Pathol. Lab. Med.* 2011 Jan;135(1):55-62. PMID: 21204711
59. Pines et al. Oncogenic mutant forms of EGFR: lessons in signal transduction and targets for cancer therapy. *FEBS Lett.* 2010 Jun 18;584(12):2699-706. PMID: 20388509
60. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. *Nature.* 2014 Jul 31;511(7511):543-50. doi: 10.1038/nature13385. Epub 2014 Jul 9. PMID: 25079552

## References (continued)

61. da et al. EGFR mutations and lung cancer. *Annu Rev Pathol.* 2011;6:49-69. doi: 10.1146/annurev-pathol-011110-130206. PMID: 20887192
62. Arcila et al. EGFR exon 20 insertion mutations in lung adenocarcinomas: prevalence, molecular heterogeneity, and clinicopathologic characteristics. *Mol. Cancer Ther.* 2013 Feb;12(2):220-9. PMID: 23371856
63. Kobayashi et al. EGFR Exon 18 Mutations in Lung Cancer: Molecular Predictors of Augmented Sensitivity to Afatinib or Neratinib as Compared with First- or Third-Generation TKIs. *Clin Cancer Res.* 2015 Dec 1;21(23):5305-13. doi: 10.1158/1078-0432.CCR-15-1046. Epub 2015 Jul 23. PMID: 26206867
64. Yasuda et al. Structural, biochemical, and clinical characterization of epidermal growth factor receptor (EGFR) exon 20 insertion mutations in lung cancer. *Sci Transl Med.* 2013 Dec 18;5(216):216ra177. PMID: 24353160
65. Chiu et al. Epidermal Growth Factor Receptor Tyrosine Kinase Inhibitor Treatment Response in Advanced Lung Adenocarcinomas with G719X/L861Q/S768I Mutations. *J Thorac Oncol.* 2015 May;10(5):793-9. PMID: 25668120
66. Karachaliou et al. KRAS mutations in lung cancer. *Clin Lung Cancer.* 2013 May;14(3):205-14. PMID: 23122493
67. Brennan et al. The somatic genomic landscape of glioblastoma. *Cell.* 2013 Oct 10;155(2):462-77. PMID: 24120142
68. Mitsudomi et al. Epidermal growth factor receptor in relation to tumor development: EGFR gene and cancer. *FEBS J.* 2010 Jan;277(2):301-8. PMID: 19922469
69. Gazdar. Activating and resistance mutations of EGFR in non-small-cell lung cancer: role in clinical response to EGFR tyrosine kinase inhibitors. *Oncogene.* 2009 Aug;28 Suppl 1:S24-31. PMID: 19680293
70. Gan et al. The EGFRvIII variant in glioblastoma multiforme. *J Clin Neurosci.* 2009 Jun;16(6):748-54. PMID: 19324552
71. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2016/021743s025lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2016/021743s025lbl.pdf)
72. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/206995s004lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/206995s004lbl.pdf)
73. Riely et al. Clinical course of patients with non-small cell lung cancer and epidermal growth factor receptor exon 19 and exon 21 mutations treated with gefitinib or erlotinib. *Clin Cancer Res.* 2006 Feb 1;12(3 Pt 1):839-44. PMID: 16467097
74. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2022/201292s017lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2022/201292s017lbl.pdf)
75. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/211288s003lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/211288s003lbl.pdf)
76. NCCN Guidelines® - NCCN-Non-Small Cell Lung Cancer [Version 8.2025]
77. Naidoo et al. Epidermal growth factor receptor exon 20 insertions in advanced lung adenocarcinomas: Clinical outcomes and response to erlotinib. *Cancer.* 2015 Sep 15;121(18):3212-3220. PMID: 26096453
78. Vyse et al. Targeting EGFR exon 20 insertion mutations in non-small cell lung cancer. *Signal Transduct Target Ther.* 2019;4:5. PMID: 30854234
79. Yi et al. A comparison of epidermal growth factor receptor mutation testing methods in different tissue types in non-small cell lung cancer. *Int J Mol Med.* 2014 Aug;34(2):464-74. PMID: 24891042
80. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/219839s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/219839s000lbl.pdf)
81. <https://investors.cullinanoncology.com/news-releases/news-release-details/fda-grants-breakthrough-therapy-designation-cullinan-oncologys>
82. Madic et al. EGFR C797S, EGFR T790M and EGFR sensitizing mutations in non-small cell lung cancer revealed by six-color crystal digital PCR. *Oncotarget.* 2018 Dec 21;9(100):37393-37406. PMID: 30647840
83. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/208065s033lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/208065s033lbl.pdf)
84. Niederst et al. The Allelic Context of the C797S Mutation Acquired upon Treatment with Third-Generation EGFR Inhibitors Impacts Sensitivity to Subsequent Treatment Strategies. *Clin. Cancer Res.* 2015 Sep 1;21(17):3924-33. PMID: 25964297
85. Wang et al. Lung Adenocarcinoma Harboring EGFR T790M and In Trans C797S Responds to Combination Therapy of First- and Third-Generation EGFR TKIs and Shifts Allelic Configuration at Resistance. *J Thorac Oncol.* 2017 Nov;12(11):1723-1727. PMID: 28662863
86. <https://investors.blackdiamondtherapeutics.com/news-releases/news-release-details/black-diamond-therapeutics-announces-corporate-update-and>
87. Ciardiello et al. The role of anti-EGFR therapies in EGFR-TKI-resistant advanced non-small cell lung cancer. *Cancer Treat Rev.* 2024 Jan;122:102664. PMID: 38064878
88. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/761210s008lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/761210s008lbl.pdf)
89. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/219008s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/219008s000lbl.pdf)
90. <https://iis.aastocks.com/20231227/11015917-0.PDF>
91. <https://www1.hkexnews.hk/listedco/listconews/sehk/2024/1008/2024100800433.pdf>

## References (continued)

92. <https://www.genprex.com/news/genprex-receives-u-s-fda-fast-track-designation-for-gene-therapy-that-targets-lung-cancer/>
93. NCCN Guidelines® - NCCN-Pediatric Central Nervous System Cancers [Version 3.2025]
94. Buccoliero et al. Pediatric High Grade Glioma Classification Criteria and Molecular Features of a Case Series. *Genes (Basel)*. 2022 Mar 31;13(4). PMID: 35456430
95. Louis et al. cIMPACT-NOW update 6: new entity and diagnostic principle recommendations of the cIMPACT-Utrecht meeting on future CNS tumor classification and grading. *Brain Pathol*. 2020 Jul;30(4):844-856. PMID: 32307792