

**Patient Name:** 박한나  
**Gender:** F  
**Sample ID:** N25-306

**Primary Tumor Site:**  
**Collection Date:** 2025.10.30

## Sample Cancer Type: Colon Cancer

Table of Contents	Page	Report Highlights
Variant Details	2	4 Relevant Biomarkers
Biomarker Descriptions	3	6 Therapies Available
Alert Details	7	37 Clinical Trials
Relevant Therapy Summary	13	

## Relevant Colon Cancer Findings

Gene	Finding	Gene	Finding
BRAF	None detected	NTRK3	None detected
ERBB2	None detected	PIK3CA	<b>PIK3CA p.(E545G) c.1634A&gt;G</b>
KRAS	<b>KRAS p.(G12D) c.35G&gt;A</b>	POLD1	None detected
NRAS	None detected	POLE	None detected
NTRK1	None detected	RET	None detected
NTRK2	None detected		

Genomic Alteration	Finding
Tumor Mutational Burden	<b>7.58 Mut/Mb measured</b>

HRD Status: **HR Proficient (HRD-)**

## Relevant Biomarkers

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IA	<b>PIK3CA p.(E545G) c.1634A&gt;G</b> phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha Allele Frequency: 24.66% Locus: chr3:178936092 Transcript: NM_006218.4	aspirin II+	<b>inavolisib + palbociclib + hormone therapy</b> <sup>1 / I</sup> <b>alpelisib + hormone therapy</b> <sup>1, 2 / II+</sup> <b>capiwasertib + hormone therapy</b> <sup>1, 2 / II</sup> + aspirin II+	4
IA	<b>KRAS p.(G12D) c.35G&gt;A</b> KRAS proto-oncogene, GTPase Allele Frequency: 38.30% Locus: chr12:25398284 Transcript: NM_033360.4	bevacizumab + chemotherapy <sup>I</sup>	<b>avutometinib + defactinib</b> <sup>1 / II+</sup>	33

\* Public data sources included in relevant therapies: FDA<sup>1</sup>, NCCN, EMA<sup>2</sup>, ESMO

\* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO

Line of therapy: I: First-line therapy, II+: Other line of therapy

Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.




Relevant Biomarkers (continued)

Tier	Genomic Alteration	Relevant Therapies (In this cancer type)	Relevant Therapies (In other cancer type)	Clinical Trials
IIC	SMAD4 deletion SMAD family member 4 Locus: chr18:48573387	None*	None*	1
IIC	TP53 p.(K132R) c.395A>G tumor protein p53 Allele Frequency: 39.86% Locus: chr17:7578535 Transcript: NM_000546.6	None*	None*	1

\* Public data sources included in relevant therapies: FDA1, NCCN, EMA2, ESMO  
\* Public data sources included in prognostic and diagnostic significance: NCCN, ESMO  
Line of therapy: I: First-line therapy, II+: Other line of therapy  
Tier Reference: Li et al. Standards and Guidelines for the Interpretation and Reporting of Sequence Variants in Cancer: A Joint Consensus Recommendation of the Association for Molecular Pathology, American Society of Clinical Oncology, and College of American Pathologists. J Mol Diagn. 2017 Jan;19(1):4-23.

 Alerts informed by public data sources:  Contraindicated,  Resistance,  Breakthrough,  Fast Track

KRAS p.(G12D) c.35G>A  cetuximab<sup>1,2</sup>, cetuximab + chemotherapy<sup>2</sup>, panitumumab<sup>1</sup>, panitumumab + chemotherapy<sup>2</sup>

Public data sources included in alerts: FDA1, NCCN, EMA2, ESMO

Prevalent cancer biomarkers without relevant evidence based on included data sources  
APC p.(Q1062\*) c.3184C>T, NF1 p.(K2664Dfs\*2) c.7989\_7990insGATT, HLA-A deletion, HLA-B deletion, NQO1 p.(P187S) c.559C>T, Tumor Mutational Burden

Variant Details

DNA Sequence Variants							
Gene	Amino Acid Change	Coding	Variant ID	Locus	Allele Frequency	Transcript	Variant Effect
PIK3CA	p.(E545G)	c.1634A>G	COSM764	chr3:178936092	24.66%	NM_006218.4	missense
KRAS	p.(G12D)	c.35G>A	COSM521	chr12:25398284	38.30%	NM_033360.4	missense
TP53	p.(K132R)	c.395A>G	COSM11582	chr17:7578535	39.86%	NM_000546.6	missense
APC	p.(Q1062*)	c.3184C>T	.	chr5:112174475	55.21%	NM_000038.6	nonsense
NF1	p.(K2664Dfs*2)	c.7989_7990insGATT	.	chr17:29685516	40.70%	NM_001042492.3	frameshift Insertion
NQO1	p.(P187S)	c.559C>T	.	chr16:69745145	51.38%	NM_000903.3	missense
PRDM9	p.(S238T)	c.712T>A	.	chr5:23522824	11.66%	NM_020227.4	missense
MSH3	p.(A57_A62del)	c.162_179delTGCAGC GGCCGCAGCGGC	.	chr5:79950707	48.11%	NM_002439.5	nonframeshift Deletion
ZER1	p.(P197L)	c.590C>T	.	chr9:131515599	29.35%	NM_006336.4	missense
PARP4	p.(?)	c.3285_3285+5delinsA GT	.	chr13:25021149	100.00%	NM_006437.4	unknown
SMAD4	p.(V350F)	c.1048G>T	.	chr18:48591885	27.33%	NM_005359.6	missense



Variant Details (continued)

Copy Number Variations			
Gene	Locus	Copy Number	CNV Ratio
SMAD4	chr18:48573387	0	0.58
HLA-A	chr6:29910229	0.1	0.62
HLA-B	chr6:31322252	0	0.57
BCL2	chr18:60795830	0.28	0.65

Biomarker Descriptions

PIK3CA p.(E545G) c.1634A>G

*phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha*

**Background:** The PIK3CA gene encodes the phosphatidylinositol-4,5-bisphosphate 3-kinase catalytic subunit alpha of the class I phosphatidylinositol 3-kinase (PI3K) enzyme<sup>75</sup>. PI3K is a heterodimer that contains a p85 regulatory subunit, which couples one of four p110 catalytic subunits to activated tyrosine protein kinases<sup>76,77</sup>. The p110 catalytic subunits include p110α, β, δ, γ and are encoded by genes PIK3CA, PIK3CB, PIK3CD, and PIK3CG, respectively<sup>76</sup>. PI3K catalyzes the conversion of phosphatidylinositol (4,5)-bisphosphate (PI(4,5)P2) into phosphatidylinositol (3,4,5)-trisphosphate (PI(3,4,5)P3) while the phosphatase and tensin homolog (PTEN) catalyzes the reverse reaction<sup>78,79</sup>. The reversible phosphorylation of inositol lipids regulates diverse aspects of cell growth and metabolism<sup>78,79,80,81</sup>. Recurrent somatic alterations in PIK3CA are frequent in cancer and result in the activation of PI3K/AKT/MTOR pathway, which can influence several hallmarks of cancer including cell proliferation, apoptosis, cancer cell metabolism and invasion, and genetic instability<sup>82,83,84</sup>.

**Alterations and prevalence:** Recurrent somatic activating mutations in PIK3CA are common in diverse cancers and are observed in 20-30% of breast, cervical, and uterine cancers and 10-20% of bladder, gastric, head and neck, and colorectal cancers<sup>8,9</sup>. Activating mutations in PIK3CA commonly occur in exons 10 and 21 (previously referred to as exons 9 and 20 due to exon 1 being untranslated)<sup>85,86</sup>. These mutations typically cluster in the exon 10 helical (codons E542/E545) and exon 21 kinase (codon H1047) domains, each having distinct mechanisms of activation<sup>87,88,89</sup>. PIK3CA resides in the 3q26 cytoband, a region frequently amplified (10-30%) in diverse cancers including squamous carcinomas of the lung, cervix, head and neck, and esophagus, and in serous ovarian and uterine cancers<sup>8,9</sup>.

**Potential relevance:** The PI3K inhibitor, alpelisib<sup>90</sup>, is FDA-approved (2019) in combination with fulvestrant for the treatment of patients with PIK3CA-mutated, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, advanced or metastatic breast cancer. Additionally, a phase Ib study of alpelisib with letrozole in patients with metastatic estrogen receptor (ER)-positive breast cancer showed the clinical benefit rate, defined as lack of disease progression ≥ 6 months, was 44% (7/16) in PIK3CA-mutated tumors and 20% (2/20) in PIK3CA wild-type tumors<sup>91</sup>. Specifically, exon 21 H1047R mutations were associated with more durable clinical responses in comparison to exon 10 E545K mutations<sup>91</sup>. However, alpelisib did not improve response when administered with letrozole in patients with ER+ early breast cancer with PIK3CA mutations<sup>92</sup>. The FDA also approved the kinase inhibitor, capivasertib (2023)<sup>93</sup> in combination with fulvestrant for locally advanced or metastatic HR-positive, HER2-negative breast cancer with one or more PIK3CA/AKT1/PTEN-alterations following progression after endocrine treatment. The kinase inhibitor, inavolisib<sup>94</sup>, is also FDA-approved (2024) in combination with palbociclib and fulvestrant for the treatment of adults with endocrine-resistant, PIK3CA-mutated, HR-positive, and HER2-negative breast cancer. Case studies with mTOR inhibitors sirolimus and temsirolimus report isolated cases of clinical response in PIK3CA mutated refractory cancers<sup>95,96</sup>. In colorectal cancers, PIK3CA mutations predict significantly improved survival and reduced disease recurrence with adjuvant aspirin therapy, compared to no benefit in wild-type PIK3CA tumors<sup>97,98,99,100</sup>.

KRAS p.(G12D) c.35G>A

*KRAS proto-oncogene, GTPase*

**Background:** The KRAS proto-oncogene encodes a GTPase that functions in signal transduction and is a member of the RAS superfamily which also includes NRAS and HRAS. RAS proteins mediate the transmission of growth signals from the cell surface to the nucleus via the PI3K/AKT/MTOR and RAS/RAF/MEK/ERK pathways, which regulate cell division, differentiation, and survival<sup>10,11,12</sup>.

**Alterations and prevalence:** Recurrent mutations in RAS oncogenes cause constitutive activation and are found in 20-30% of cancers. KRAS mutations are observed in up to 10-20% of uterine cancer, 30-35% of lung adenocarcinoma and colorectal cancer, and about 60%



## Biomarker Descriptions (continued)

of pancreatic cancer<sup>8</sup>. The majority of KRAS mutations consist of point mutations occurring at G12, G13, and Q61<sup>8,13,14</sup>. Mutations at A59, K117, and A146 have also been observed but are less frequent<sup>9,15</sup>.

**Potential relevance:** The FDA has approved the small molecule inhibitors, sotorasib<sup>16</sup> (2021) and adagrasib<sup>17</sup> (2022), for the treatment of adult patients with KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC). Sotorasib and adagrasib are also useful in certain circumstances for KRAS G12C-mutated pancreatic adenocarcinoma<sup>18</sup>. The FDA has approved the combination of kinase inhibitors, avutometinib and defactinib<sup>19</sup> (2025), for the treatment of adult patients with KRAS-mutated recurrent low-grade serous ovarian cancer (LGSOC) after prior systemic therapy. The FDA has granted breakthrough therapy designation (2022) to the KRAS G12C inhibitor, GDC-6036<sup>20</sup>, for KRAS G12C-mutated NSCLC. The KRAS-G12C/NRAS-G12C dual inhibitor, elironrasib<sup>21</sup>, and the KRAS G12C inhibitor, D3S-001<sup>22</sup>, were both granted breakthrough therapy designation (2025) for KRAS G12C-mutated locally advanced or metastatic NSCLC in adults previously treated with chemotherapy and immunotherapy, excluding KRAS G12C inhibitors. The KRAS-G12C inhibitor, olomorasib<sup>23</sup>, was granted breakthrough designation (2025) in combination with pembrolizumab<sup>24</sup> for unresectable advanced or metastatic NSCLC with a KRAS G12C mutation and PD-L1 expression  $\geq 50\%$ . The SHP2 inhibitor, BBP-398<sup>25</sup> was granted fast track designation (2022) in combination with sotorasib for previously treated patients with KRAS G12C-mutated metastatic NSCLC. The RAF/MEK clamp, avutometinib<sup>26</sup> was also granted fast track designation (2024) in combination with sotorasib for KRAS G12C-mutated metastatic NSCLC in patients who have received at least one prior systemic therapy and have not been previously treated with a KRAS G12C inhibitor. The KRAS G12C inhibitor, BBO-8520<sup>27</sup>, was granted fast track designation in 2025 for previously treated KRAS G12C-mutated patients with metastatic NSCLC. The RAS inhibitor, daraxonrasib<sup>28</sup>, was granted breakthrough designation (2025) for previously treated metastatic pancreatic cancer with KRAS G12 mutations. The KRAS G12D (ON/OFF) inhibitor, GFH-375<sup>29</sup>, was also granted fast track designation (2025) for first-line and previously treated KRAS G12D-mutated locally advanced or metastatic pancreatic adenocarcinoma. The KRAS G12C inhibitor, D3S-001<sup>30</sup>, was granted fast track designation in 2024 for KRAS G12C-mutated patients with advanced unresectable or metastatic colorectal cancers. The PLK1 inhibitor, onvansertib<sup>31</sup>, was granted fast track designation (2020) in combination with bevacizumab and FOLFIRI for second-line treatment of patients with KRAS-mutated metastatic colorectal cancer (mCRC). The EGFR antagonists, cetuximab<sup>32</sup> and panitumumab<sup>33</sup>, are contraindicated for treatment of colorectal cancer patients with KRAS mutations in exon 2 (codons 12 and 13), exon 3 (codons 59 and 61), and exon 4 (codons 117 and 146)<sup>15</sup>. Additionally, KRAS mutations are associated with poor prognosis in NSCLC<sup>34</sup>.

### SMAD4 deletion

#### SMAD family member 4

**Background:** The SMAD4 gene encodes the SMAD family member 4, a transcription factor that belongs to a family of 8 SMAD genes that can be divided into three main classes. SMAD4 (also known as DPC4) belongs to the common mediator SMAD (co-SMAD) class while SMAD1, SMAD2, SMAD3, SMAD5, and SMAD8 are part of the regulator SMAD (R-SMAD) class. The inhibitory SMAD (I-SMAD) class includes both SMAD6 and SMAD7<sup>35,36</sup>. SMAD4 is a tumor suppressor gene and functions as a mediator of the TGF- $\beta$  and BMP signaling pathways that are implicated in cancer initiation and progression<sup>36,37,38</sup>. Loss of SMAD4 does not drive oncogenesis, but is associated with progression of cancers initiated by driver genes such as KRAS and APC<sup>35,36</sup>.

**Alterations and prevalence:** Inactivation of SMAD4 can occur due to mutations, allelic loss, homozygous deletions, and 18q loss of heterozygosity (LOH)<sup>35</sup>. Somatic mutations in SMAD4 occur in up to 20% of pancreatic, 12% of colorectal, and 8% of stomach cancers. Recurrent hotspot mutations including R361 and P356 occur in the mad homology 2 (MH2) domain leading to the disruption of the TGF- $\beta$  signaling<sup>9,38,39</sup>. Copy number deletions occur in up to 12% of pancreatic, 10% of esophageal, and 13% of stomach cancers<sup>8,9,40</sup>.

**Potential relevance:** Currently, no therapies are approved for SMAD4 aberrations. Clinical studies and meta-analyses have demonstrated that loss of SMAD4 expression confers poor prognosis and poor overall survival (OS) in colorectal and pancreatic cancers<sup>36,38,41,42,43</sup>. Importantly, SMAD4 is a predictive biomarker to fluorouracil based chemotherapy<sup>44,45</sup>. In a retrospective analysis of 241 colorectal cancer patients treated with fluorouracil, 21 patients with SMAD4 loss demonstrated significantly poor median OS when compared to SMAD4 positive patients (31 months vs 89 months)<sup>45</sup>. In another clinical study of 173 newly diagnosed and recurrent head and neck squamous cell carcinoma (HNSCC) patients, SMAD4 loss is correlated with cetuximab resistance in HPV-negative HNSCC tumors<sup>46</sup>.

### TP53 p.(K132R) c.395A>G

#### tumor protein p53

**Background:** The TP53 gene encodes the tumor suppressor protein p53, which binds to DNA and activates transcription in response to diverse cellular stresses to induce cell cycle arrest, apoptosis, or DNA repair<sup>1</sup>. In unstressed cells, TP53 is kept inactive by targeted degradation via MDM2, a substrate recognition factor for ubiquitin-dependent proteolysis<sup>53</sup>. Alterations in TP53 are required for oncogenesis as they result in loss of protein function and gain of transforming potential<sup>54</sup>. Germline mutations in TP53 are the underlying cause of Li-Fraumeni syndrome, a complex hereditary cancer predisposition disorder associated with early-onset cancers<sup>55,56</sup>.



## Biomarker Descriptions (continued)

**Alterations and prevalence:** TP53 is the most frequently mutated gene in the cancer genome with approximately half of all cancers experiencing TP53 mutations. Ovarian, head and neck, esophageal, and lung squamous cancers have particularly high TP53 mutation rates (60-90%)<sup>8,9,57,58,59,60</sup>. Approximately two-thirds of TP53 mutations are missense mutations and several recurrent missense mutations are common, including substitutions at codons R158, R175, Y220, R248, R273, and R282<sup>8,9</sup>. Invariably, recurrent missense mutations in TP53 inactivate its ability to bind DNA and activate transcription of target genes<sup>61,62,63,64</sup>. Alterations in TP53 are also observed in pediatric cancers<sup>8,9</sup>. Somatic mutations are observed in 53% of non-Hodgkin lymphoma, 24% of soft tissue sarcoma, 19% of glioma, 13% of bone cancer, 9% of B-lymphoblastic leukemia/lymphoma, 4% of embryonal tumors, 3% of Wilms tumor and leukemia, 2% of T-lymphoblastic leukemia/lymphoma, and less than 1% of peripheral nervous system cancers (5 in 1158 cases)<sup>8,9</sup>. Biallelic loss of TP53 is observed in 10% of bone cancer, 2% of Wilms tumor, and less than 1% of B-lymphoblastic leukemia/lymphoma (2 in 731 cases) and leukemia (1 in 250 cases)<sup>8,9</sup>.

**Potential relevance:** The small molecule p53 reactivator, PC14586<sup>65</sup> (2020), received a fast track designation by the FDA for advanced tumors harboring a TP53 Y220C mutation. In addition to investigational therapies aimed at restoring wild-type TP53 activity, compounds that induce synthetic lethality are also under clinical evaluation<sup>66,67</sup>. TP53 mutation are a diagnostic marker of SHH-activated, TP53-mutant medulloblastoma<sup>68</sup>. TP53 mutations confer poor prognosis and poor risk in multiple blood cancers including AML, MDS, myeloproliferative neoplasms (MPN), and chronic lymphocytic leukemia (CLL), and acute lymphoblastic leukemia (ALL)<sup>50,69,70,71,72</sup>. In mantle cell lymphoma, TP53 mutations are associated with poor prognosis when treated with conventional therapy including hematopoietic cell transplant<sup>73</sup>. Mono- and bi-allelic mutations in TP53 confer unique characteristics in MDS, with multi-hit patients also experiencing associations with complex karyotype, few co-occurring mutations, and high-risk disease presentation as well as predicted death and leukemic transformation independent of the IPSS-R staging system<sup>74</sup>.

### APC p.(Q1062\*) c.3184C>T

*APC, WNT signaling pathway regulator*

**Background:** The APC gene encodes the adenomatous polyposis coli tumor suppressor protein that plays a crucial role in regulating the  $\beta$ -catenin/WNT signaling pathway which is involved in cell migration, adhesion, proliferation, and differentiation<sup>101</sup>. APC is an antagonist of WNT signaling as it targets  $\beta$ -catenin for proteasomal degradation<sup>102,103</sup>. Germline mutations in APC are predominantly inactivating and result in an autosomal dominant predisposition for familial adenomatous polyposis (FAP) which is characterized by numerous polyps in the intestine<sup>101,104</sup>. Acquiring a somatic mutation in APC is considered to be an early and possibly initiating event in colorectal cancer<sup>105</sup>.

**Alterations and prevalence:** Somatic mutations in APC are observed in up to 65% of colorectal cancer, and in up to 15% of stomach adenocarcinoma and uterine corpus endometrial carcinoma<sup>8,9,40</sup>. In colorectal cancer, ~60% of somatic APC mutations have been reported to occur in a mutation cluster region (MCR) resulting in C-terminal protein truncation and APC inactivation<sup>106,107</sup>.

**Potential relevance:** Currently, no therapies are approved for APC aberrations.

### NF1 p.(K2664Dfs\*2) c.7989\_7990insGATT

*neurofibromin 1*

**Background:** The NF1 gene encodes the neurofibromin protein, a tumor suppressor within the Ras-GTPase-activating protein (GAP) family<sup>47</sup>. NF1 regulates cellular levels of activated RAS proteins including KRAS, NRAS, and HRAS, by down regulating the active GTP-bound state to an inactive GDP-bound state<sup>47,48</sup>. Inactivation of NF1 due to missense mutations results in sustained intracellular levels of RAS-GTP and prolonged activation of the RAS/RAF/MAPK and PI3K/AKT/mTOR signaling pathways leading to increased proliferation and survival<sup>47</sup>. Constitutional mutations in NF1 are associated with neurofibromatosis type 1, a RASopathy autosomal dominant tumor syndrome with predisposition to myeloid malignancies such as juvenile myelomonocytic leukemia (JMML) and myeloproliferative neoplasms (MPN)<sup>47,49,50</sup>.

**Alterations and prevalence:** NF1 aberrations include missense mutations, insertions, indels, aberrant splicing, microdeletions, and rearrangements<sup>47</sup>. The majority of NF1 mutated tumors exhibit biallelic inactivation of NF1, supporting the 'two-hit' hypothesis of carcinogenesis<sup>47,51</sup>. Somatic mutations in NF1 have been identified in over 30% of ovarian serous carcinoma, 12-30% of melanoma, 10-20% of chronic myelomonocytic leukemia (CMML), and 7% of acute myeloid leukemia (AML)<sup>47,50</sup>.

**Potential relevance:** Currently, no therapies are approved for NF1 aberrations. Somatic mutation of NF1 is useful as an ancillary diagnostic marker for malignant peripheral nerve sheath tumor (MPNST)<sup>52</sup>.



## Biomarker Descriptions (continued)

### HLA-A deletion

*major histocompatibility complex, class I, A*

**Background:** The HLA-A gene encodes the major histocompatibility complex, class I, A<sup>1</sup>. MHC (major histocompatibility complex) class I molecules are located on the cell surface of nucleated cells and present antigens from within the cell for recognition by cytotoxic T cells<sup>2</sup>. MHC class I molecules are heterodimers composed of two polypeptide chains,  $\alpha$  and B2M<sup>3</sup>. The classical MHC class I genes include HLA-A, HLA-B, and HLA-C and encode the  $\alpha$  polypeptide chains, which present short polypeptide chains, of 7 to 11 amino acids, to the immune system to distinguish self from non-self<sup>4,5,6</sup>. Downregulation of MHC class I promotes tumor evasion of the immune system, suggesting a tumor suppressor role for HLA-A<sup>7</sup>.

**Alterations and prevalence:** Somatic mutations in HLA-A are observed in 7% of diffuse large B-cell lymphoma (DLBCL), 4% of cervical squamous cell carcinoma and head and neck squamous cell carcinoma, 3% of colorectal adenocarcinoma, and 2% of uterine corpus endometrial carcinoma and stomach adenocarcinoma<sup>8,9</sup>. Biallelic loss of HLA-A is observed in 4% of DLBCL<sup>8,9</sup>.

**Potential relevance:** Currently, no therapies are approved for HLA-A aberrations.

### HLA-B deletion

*major histocompatibility complex, class I, B*

**Background:** The HLA-B gene encodes the major histocompatibility complex, class I, B<sup>1</sup>. MHC (major histocompatibility complex) class I molecules are located on the cell surface of nucleated cells and present antigens from within the cell for recognition by cytotoxic T cells<sup>2</sup>. MHC class I molecules are heterodimers composed of two polypeptide chains,  $\alpha$  and B2M<sup>3</sup>. The classical MHC class I genes include HLA-A, HLA-B, and HLA-C and encode the  $\alpha$  polypeptide chains, which present short polypeptide chains, of 7 to 11 amino acids, to the immune system to distinguish self from non-self<sup>4,5,6</sup>. Downregulation of MHC class I promotes tumor evasion of the immune system, suggesting a tumor suppressor role for HLA-B<sup>7</sup>.

**Alterations and prevalence:** Somatic mutations in HLA-B are observed in 10% of diffuse large B-cell lymphoma (DLBCL), 5% of cervical squamous cell carcinoma and stomach adenocarcinoma, 4% of head and neck squamous cell carcinoma and colorectal adenocarcinoma, 3% of uterine cancer, and 2% of esophageal adenocarcinoma and skin cutaneous melanoma<sup>8,9</sup>. Biallelic loss of HLA-B is observed in 5% of DLBCL<sup>8,9</sup>.

**Potential relevance:** Currently, no therapies are approved for HLA-B aberrations.



Alerts Informed By Public Data Sources

Current FDA Information

Contraindicated   Not recommended   Resistance   Breakthrough   Fast Track

FDA information is current as of 2025-09-17. For the most up-to-date information, search [www.fda.gov](https://www.fda.gov).

KRAS p.(G12D) c.35G>A

cetuximab

Cancer type: Colorectal Cancer   Label as of: 2021-09-24   Variant class: KRAS G12 mutation

Indications and usage:

Erbixux® is an epidermal growth factor receptor (EGFR) antagonist indicated for treatment of:

Head and Neck Cancer

- Locally or regionally advanced squamous cell carcinoma of the head and neck in combination with radiation therapy.
- Recurrent locoregional disease or metastatic squamous cell carcinoma of the head and neck in combination with platinum-based therapy with fluorouracil.
- Recurrent or metastatic squamous cell carcinoma of the head and neck progressing after platinum-based therapy.

Colorectal Cancer

K-Ras wild-type, EGFR-expressing, metastatic colorectal cancer as determined by FDA-approved test

- in combination with FOLFIRI for first-line treatment,
- in combination with irinotecan in patients who are refractory to irinotecan-based chemotherapy,
- as a single agent in patients who have failed oxaliplatin- and irinotecan-based chemotherapy or who are intolerant to irinotecan.

Limitations of Use: Erbixux® is not indicated for treatment of Ras-mutant colorectal cancer or when the results of the Ras mutation tests are unknown.

BRAF V600E Mutation-Positive Metastatic Colorectal Cancer (CRC)

- in combination with encorafenib, for the treatment of adult patients with metastatic colorectal cancer (CRC) with a BRAF V600E mutation, as detected by an FDA-approved test, after prior therapy.

Reference:

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125084s279lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf)



## KRAS p.(G12D) c.35G>A (continued)

### panitumumab

**Cancer type:** Colorectal Cancer

**Label as of:** 2025-01-16

**Variant class:** KRAS G12 mutation

**Indications and usage:**

VECTIBIX® is an epidermal growth factor receptor (EGFR) antagonist indicated for the treatment of:

Adult patients with wild-type RAS (defined as wild-type in both KRAS and NRAS as determined by an FDA-approved test) Metastatic Colorectal Cancer (mCRC)\*:

- In combination with FOLFOX for first-line treatment.
- As monotherapy following disease progression after prior treatment with fluoropyrimidine, oxaliplatin, and irinotecan-containing chemotherapy.

KRAS G12C-mutated Metastatic Colorectal Cancer (mCRC)\*

- In combination with sotorasib, for the treatment of adult patients with KRAS G12C-mutated mCRC, as determined by an FDA-approved test, who have received prior treatment with fluoropyrimidine-, oxaliplatin-, and irinotecan-based chemotherapy.

\*Limitations of Use: VECTIBIX® is not indicated for the treatment of patients with RAS-mutant mCRC unless used in combination with sotorasib in KRAS G12C-mutated mCRC. VECTIBIX® is not indicated for the treatment of patients with mCRC for whom RAS mutation status is unknown.

**Reference:**

[https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125147s213lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125147s213lbl.pdf)

### daraxonrasib

**Cancer type:** Pancreatic Cancer

**Variant class:** KRAS G12 mutation

**Supporting Statement:**

The FDA has granted Breakthrough designation to the RAS inhibitor, daraxonrasib, for previously treated metastatic pancreatic adenocarcinoma (PDAC) in patients with KRAS G12 mutations.

**Reference:**

<https://ir.revmed.com/news-releases/news-release-details/revolution-medicines-announces-fda-breakthrough-therapy>

### GFH-375

**Cancer type:** Pancreatic Cancer

**Variant class:** KRAS G12D mutation

**Supporting Statement:**


The FDA has granted Fast Track designation to an oral KRAS G12D (ON/OFF) inhibitor, GFH-375 (VS-7375), for the first-line treatment of patients with KRAS G12D-mutated locally advanced or metastatic adenocarcinoma of the pancreas (PDAC) and for the treatment of patients with KRAS G12D-mutated locally advanced or metastatic PDAC who have received at least one prior line of standard systemic therapy.


**Reference:**

<https://investor.verastem.com/news-releases/news-release-details/verastem-oncology-granted-fast-track-designation-vs-7375>





## Current NCCN Information

 Contraindicated

 Not recommended

 Resistance

 Breakthrough

 Fast Track

NCCN information is current as of 2025-09-02. To view the most recent and complete version of the guideline, go online to [NCCN.org](https://www.nccn.org).

For NCCN International Adaptations & Translations, search [www.nccn.org/global/what-we-do/international-adaptations](https://www.nccn.org/global/what-we-do/international-adaptations).

Some variant specific evidence in this report may be associated with a broader set of alterations from the NCCN Guidelines. Specific variants listed in this report were sourced from approved therapies or scientific literature. These therapeutic options are appropriate for certain population segments with cancer. Refer to the NCCN Guidelines® for full recommendation.

All guidelines cited below are referenced with permission from the NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®) National Comprehensive Cancer Network, Inc. 2023. All rights reserved. NCCN makes no warranties regarding their content.

### KRAS p.(G12D) c.35G>A

#### cetuximab

**Cancer type:** Colon Cancer

**Variant class:** KRAS G12 mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

**Reference:** NCCN Guidelines® - NCCN-Colon Cancer [Version 4.2025]

#### panitumumab

**Cancer type:** Colon Cancer

**Variant class:** KRAS G12 mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exon 2, 3, 4) or NRAS mutation (exon 2, 3, 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

**Reference:** NCCN Guidelines® - NCCN-Colon Cancer [Version 4.2025]

#### cetuximab

**Cancer type:** Rectal Cancer

**Variant class:** KRAS G12 mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exons 2, 3, and 4) or NRAS mutation (exons 2, 3, and 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

**Reference:** NCCN Guidelines® - NCCN-Rectal Cancer [Version 3.2025]



**KRAS p.(G12D) c.35G>A (continued)****🚫 panitumumab**

Cancer type: Rectal Cancer

Variant class: KRAS G12 mutation

**Summary:**

NCCN Guidelines® include the following supporting statement(s):

- "Patients with any known KRAS mutation (exons 2, 3, and 4) or NRAS mutation (exons 2, 3, and 4) should not be treated with either cetuximab or panitumumab, unless given as part of a regimen targeting a KRAS G12C mutation."

Reference: NCCN Guidelines® - NCCN-Rectal Cancer [Version 3.2025]

**Current EMA Information****🚫** Contraindicated**⊖** Not recommended**🛡️** Resistance**🚀** Breakthrough**⚡** Fast TrackEMA information is current as of 2025-09-17. For the most up-to-date information, search [www.ema.europa.eu](http://www.ema.europa.eu).**KRAS p.(G12D) c.35G>A****🚫 cetuximab, cetuximab + oxaliplatin**

Cancer type: Colorectal Cancer

Label as of: 2025-01-16

Variant class: KRAS G12 mutation

**Reference:**[https://www.ema.europa.eu/en/documents/product-information/erbitux-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/erbitux-epar-product-information_en.pdf)**🚫 panitumumab + oxaliplatin**

Cancer type: Colorectal Cancer

Label as of: 2025-05-07

Variant class: KRAS G12 mutation

**Reference:**[https://www.ema.europa.eu/en/documents/product-information/vectibix-epar-product-information\\_en.pdf](https://www.ema.europa.eu/en/documents/product-information/vectibix-epar-product-information_en.pdf)



## Current ESMO Information

 Contraindicated  Not recommended  Resistance  Breakthrough  Fast Track

ESMO information is current as of 2025-09-02. For the most up-to-date information, search [www.esmo.org](http://www.esmo.org).

### KRAS p.(G12D) c.35G>A

#### cetuximab

Cancer type: Colorectal Cancer

Variant class: KRAS G12 mutation

##### Summary:

ESMO Clinical Practice Guidelines include the following supporting statement:

- "The presence of RAS mutations is associated with resistance to anti-EGFR mAbs and knowing the expanded RAS mutational status is mandatory for use of both cetuximab and panitumumab, avoiding anti-EGFR mAb treatment when a RAS mutation is confirmed".
- "RAS testing is mandatory before treatment with anti-EGFR mAbs and can be carried out on either the primary tumor or other metastatic sites [III, A]".

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2022.10.003> (published)]

#### panitumumab

Cancer type: Colorectal Cancer

Variant class: KRAS G12 mutation

##### Summary:

ESMO Clinical Practice Guidelines include the following supporting statement:

- "The presence of RAS mutations is associated with resistance to anti-EGFR mAbs and knowing the expanded RAS mutational status is mandatory for use of both cetuximab and panitumumab, avoiding anti-EGFR mAb treatment when a RAS mutation is confirmed".
- "RAS testing is mandatory before treatment with anti-EGFR mAbs and can be carried out on either the primary tumor or other metastatic sites [III, A]".

**Reference:** ESMO Clinical Practice Guidelines - ESMO-Metastatic Colorectal Cancer [Ann Oncol (2023); <https://doi.org/10.1016/j.annonc.2022.10.003> (published)]

## Genes Assayed

### Genes Assayed for the Detection of DNA Sequence Variants

ABL1, ABL2, ACVR1, AKT1, AKT2, AKT3, ALK, AR, ARAF, ATP1A1, AURKA, AURKB, AURKC, AXL, BCL2, BCL2L12, BCL6, BCR, BMP5, BRAF, BTK, CACNA1D, CARD11, CBL, CCND1, CCND2, CCND3, CCNE1, CD79B, CDK4, CDK6, CHD4, CSF1R, CTNNB1, CUL1, CYSLTR2, DDR2, DGCR8, DROSHA, E2F1, EGFR, EIF1AX, EPAS1, ERBB2, ERBB3, ERBB4, ESR1, EZH2, FAM135B, FGF7, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FOXL2, FOXO1, GATA2, GLI1, GNA11, GNAQ, GNAS, HIF1A, HRAS, IDH1, IDH2, IKBKB, IL6ST, IL7R, IRF4, IRS4, KCNJ5, KDR, KIT, KLF4, KLF5, KNSTRN, KRAS, MAGOH, MAP2K1, MAP2K2, MAPK1, MAX, MDM4, MECOM, MED12, MEF2B, MET, MITF, MPL, MTOR, MYC, MYCN, MYD88, MYO10, NFE2L2, NRAS, NSD2, NT5C2, NTRK1, NTRK2, NTRK3, NUP93, PAX5, PCBP1, PDGFRA, PDGFRB, PIK3C2B, PIK3CA, PIK3CB, PIK3CD, PIK3CG, PIK3R2, PIM1, PLCG1, PPP2R1A, PPP6C, PRKACA, PTPN11, PTPRD, PXDN, RAC1, RAF1, RARA, RET, RGS7, RHEB, RHOA, RICTOR, RIT1, ROS1, RPL10, SETBP1, SF3B1, SIX1, SIX2, SLC01B3, SMC1A, SMO, SNCAIP, SOS1, SOX2, SPOP, SRC, SRSF2, STAT3, STAT5B, STAT6, TAF1, TERT, TGFBR1, TOP1, TOP2A, TPM1, TRRAP, TSHR, U2AF1, USP8, WAS, XPO1, ZNF217, ZNF429



## Genes Assayed (continued)

### Genes Assayed for the Detection of Copy Number Variations

ABCB1, ABL1, ABL2, ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AKT1, AKT2, AKT3, ALK, AMER1, APC, AR, ARAF, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AURKA, AURKC, AXIN1, AXIN2, AXL, B2M, BAP1, BARD1, BCL2, BCL2L12, BCL6, BCOR, BLM, BMPR2, BRAF, BRCA1, BRCA2, BRIP1, CARD11, CASP8, CBFB, CBL, CCND1, CCND2, CCND3, CCNE1, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDK4, CDK6, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHD4, CHEK1, CHEK2, CIC, CREBBP, CSMD3, CTCF, CTLA4, CTNND2, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, DAXX, DDR1, DDR2, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, EGFR, EIF1AX, ELF3, EMSY, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERBB2, ERBB3, ERBB4, ERCC2, ERCC4, ERRFI1, ESR1, ETV6, EZH2, FAM135B, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAT1, FBXW7, FGF19, FGF23, FGF3, FGF4, FGF9, FGFR1, FGFR2, FGFR3, FGFR4, FLT3, FLT4, FOXA1, FUBP1, FYN, GATA2, GATA3, GLI3, GNA13, GNAS, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, IDH2, IGF1R, IKBKB, IL7R, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KDR, KEAP1, KIT, KLF5, KMT2A, KMT2B, KMT2C, KMT2D, KRAS, LARP4B, LATS1, LATS2, MAGOH, MAP2K1, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK1, MAPK8, MAX, MCL1, MDM2, MDM4, MECOM, MEF2B, MEN1, MET, MGA, MITF, MLH1, MLH3, MPL, MRE11, MSH2, MSH3, MSH6, MTAP, MTOR, MUTYH, MYC, MYCL, MYCN, MYD88, NBN, NCOR1, NF1, NF2, NFE2L2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, NRAS, NTRK1, NTRK3, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PCBP1, PDCD1, PDCD1LG2, PDGFRA, PDGFRB, PDIA3, PGD, PHF6, PIK3C2B, PIK3CA, PIK3CB, PIK3R1, PIK3R2, PIM1, PLCG1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R1A, PPP2R2A, PPP6C, PRDM1, PRDM9, PRKACA, PRKAR1A, PTCH1, PTEN, PTPN11, PTPRT, PXDN, RAC1, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RAF1, RARA, RASA1, RASA2, RB1, RBM10, RECQL4, RET, RHEB, RICTOR, RIT1, RNASEH2A, RNASEH2B, RNF43, ROS1, RPA1, RPS6KB1, RPTOR, RUNX1, SDHA, SDHB, SDHD, SETBP1, SETD2, SF3B1, SLC01B3, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SMC1A, SMO, SOX9, SPEN, SPOP, SRC, STAG2, STAT3, STAT6, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TERT, TET2, TGFB2, TNFAIP3, TNFRSF14, TOP1, TP53, TP63, TPMT, TPP2, TSC1, TSC2, U2AF1, USP8, USP9X, VHL, WT1, XPO1, XRCC2, XRCC3, YAP1, YES1, ZFH3, ZMYM3, ZNF217, ZNF429, ZRSR2

### Genes Assayed for the Detection of Fusions

AKT2, ALK, AR, AXL, BRAF, BRCA1, BRCA2, CDKN2A, EGFR, ERBB2, ERBB4, ERG, ESR1, ETV1, ETV4, ETV5, FGFR1, FGFR2, FGFR3, FGR, FLT3, JAK2, KRAS, MDM4, MET, MYB, MYBL1, NF1, NOTCH1, NOTCH4, NRG1, NTRK1, NTRK2, NTRK3, NUTM1, PDGFRA, PDGFRB, PIK3CA, PPARG, PRKACA, PRKACB, PTEN, RAD51B, RAF1, RB1, RELA, RET, ROS1, RSPO2, RSPO3, TERT

### Genes Assayed with Full Exon Coverage

ABRAXAS1, ACVR1B, ACVR2A, ADAMTS12, ADAMTS2, AMER1, APC, ARHGAP35, ARID1A, ARID1B, ARID2, ARID5B, ASXL1, ASXL2, ATM, ATR, ATRX, AXIN1, AXIN2, B2M, BAP1, BARD1, BCOR, BLM, BMPR2, BRCA1, BRCA2, BRIP1, CALR, CASP8, CBFB, CD274, CD276, CDC73, CDH1, CDH10, CDK12, CDKN1A, CDKN1B, CDKN2A, CDKN2B, CDKN2C, CHEK1, CHEK2, CIC, CIITA, CREBBP, CSMD3, CTCF, CTLA4, CUL3, CUL4A, CUL4B, CYLD, CYP2C9, CYP2D6, DAXX, DDX3X, DICER1, DNMT3A, DOCK3, DPYD, DSC1, DSC3, ELF3, ENO1, EP300, EPCAM, EPHA2, ERAP1, ERAP2, ERCC2, ERCC4, ERCC5, ERRFI1, ETV6, FANCA, FANCC, FANCD2, FANCE, FANCF, FANCG, FANCI, FANCL, FANCM, FAS, FAT1, FBXW7, FUBP1, GATA3, GNA13, GPS2, HDAC2, HDAC9, HLA-A, HLA-B, HNF1A, ID3, INPP4B, JAK1, JAK2, JAK3, KDM5C, KDM6A, KEAP1, KLHL13, KMT2A, KMT2B, KMT2C, KMT2D, LARP4B, LATS1, LATS2, MAP2K4, MAP2K7, MAP3K1, MAP3K4, MAPK8, MEN1, MGA, MLH1, MLH3, MRE11, MSH2, MSH3, MSH6, MTAP, MTUS2, MUTYH, NBN, NCOR1, NF1, NF2, NOTCH1, NOTCH2, NOTCH3, NOTCH4, PALB2, PARP1, PARP2, PARP3, PARP4, PBRM1, PDCD1, PDCD1LG2, PDIA3, PGD, PHF6, PIK3R1, PMS1, PMS2, POLD1, POLE, POT1, PPM1D, PPP2R2A, PRDM1, PRDM9, PRKAR1A, PSMB10, PSMB8, PSMB9, PTCH1, PTEN, PTPRT, RAD50, RAD51, RAD51B, RAD51C, RAD51D, RAD52, RAD54L, RASA1, RASA2, RB1, RBM10, RECQL4, RNASEH2A, RNASEH2B, RNASEH2C, RNF43, RPA1, RPL22, RPL5, RUNX1, RUNX1T1, SDHA, SDHB, SDHC, SDHD, SETD2, SLX4, SMAD2, SMAD4, SMARCA4, SMARCB1, SOCS1, SOX9, SPEN, STAG2, STAT1, STK11, SUFU, TAP1, TAP2, TBX3, TCF7L2, TET2, TGFB2, TMEM132D, TNFAIP3, TNFRSF14, TP53, TP63, TPP2, TSC1, TSC2, UGT1A1, USP9X, VHL, WT1, XRCC2, XRCC3, ZBTB20, ZFH3, ZMYM3, ZRSR2



## Relevant Therapy Summary

● In this cancer type    
 ○ In other cancer type    
 ⓘ In this cancer type and other cancer types    
 × No evidence

### PIK3CA p.(E545G) c.1634A>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
alpelisib + fulvestrant	○	○	○	○	×
capiwasertib + fulvestrant	○	○	○	×	×
inavolisib + palbociclib + fulvestrant	○	○	×	×	×
aspirin	×	ⓘ	×	×	×
ETX-636	×	×	×	×	● (I/II)
HTL-0039732, atezolizumab	×	×	×	×	● (I/II)
JS-105	×	×	×	×	● (I)
SNV-4818, hormone therapy	×	×	×	×	● (I)

### KRAS p.(G12D) c.35G>A

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
avutometinib + defactinib	○	○	×	×	×
bevacizumab + CAPOX	×	×	×	●	×
bevacizumab + FOLFIRI	×	×	×	●	×
bevacizumab + FOLFOX	×	×	×	●	×
bevacizumab + FOLFOXIRI	×	×	×	●	×
bevacizumab, chemotherapy	×	×	×	×	● (III)
fruquintinib, chemotherapy	×	×	×	×	● (II)
regorafenib	×	×	×	×	● (II)
tunlametinib, vemurafenib	×	×	×	×	● (II)
almonertinib, palbociclib	×	×	×	×	● (I/II)
anti-KRAS G12D mTCR	×	×	×	×	● (I/II)
APR-1051	×	×	×	×	● (I/II)
ARV-806	×	×	×	×	● (I/II)
DN-022150	×	×	×	×	● (I/II)
ERAS-0015	×	×	×	×	● (I/II)
GFH-375	×	×	×	×	● (I/II)
QLC-1101, QL1203, pembrolizumab (Qilu Pharmaceutical), iparomlimab and tuvonralimab, chemotherapy	×	×	×	×	● (I/II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



## Relevant Therapy Summary (continued)

☒ In this cancer type    
 ☐ In other cancer type    
 ☒ In this cancer type and other cancer types    
 ✕ No evidence

### KRAS p.(G12D) c.35G>A (continued)

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
RNK-08954	✕	✕	✕	✕	● (I/II)
TSN-1611	✕	✕	✕	✕	● (I/II)
YL-15293	✕	✕	✕	✕	● (I/II)
ASP-4396	✕	✕	✕	✕	● (I)
ASP-5834	✕	✕	✕	✕	● (I)
AST-NS2101	✕	✕	✕	✕	● (I)
BPI-442096	✕	✕	✕	✕	● (I)
daraxonrasib	✕	✕	✕	✕	● (I)
GDC-7035	✕	✕	✕	✕	● (I)
imatinib, trametinib	✕	✕	✕	✕	● (I)
IX-001	✕	✕	✕	✕	● (I)
JAB-3312	✕	✕	✕	✕	● (I)
KRAS TCR, aldesleukin, SLATE 001, chemotherapy	✕	✕	✕	✕	● (I)
KRAS-EphA-2-CAR-DC, anti-PD-1, ipilimumab	✕	✕	✕	✕	● (I)
Nest-1	✕	✕	✕	✕	● (I)
NT-112, AZD-0240	✕	✕	✕	✕	● (I)
NW-301D	✕	✕	✕	✕	● (I)
PT-0253	✕	✕	✕	✕	● (I)
QLC-1101	✕	✕	✕	✕	● (I)
RMC-9805, daraxonrasib	✕	✕	✕	✕	● (I)
toripalimab, chemotherapy, KRAS peptide vaccine	✕	✕	✕	✕	● (I)

### SMAD4 deletion

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
regorafenib	✕	✕	✕	✕	● (II)

### TP53 p.(K132R) c.395A>G

Relevant Therapy	FDA	NCCN	EMA	ESMO	Clinical Trials*
APR-1051	✕	✕	✕	✕	● (I/II)

\* Most advanced phase (IV, III, II/III, II, I/II, I) is shown and multiple clinical trials may be available.



Thermo Fisher Scientific's Ion Torrent OncoPrint Reporter software was used in generation of this report. Software was developed and designed internally by Thermo Fisher Scientific. The analysis was based on OncoPrint Reporter (6.1.1 data version 2025.10(006)). The data presented here are from a curated knowledge base of publicly available information, but may not be exhaustive. FDA information was sourced from [www.fda.gov](http://www.fda.gov) and is current as of 2025-09-17. NCCN information was sourced from [www.nccn.org](http://www.nccn.org) and is current as of 2025-09-02. EMA information was sourced from [www.ema.europa.eu](http://www.ema.europa.eu) and is current as of 2025-09-17. ESMO information was sourced from [www.esmo.org](http://www.esmo.org) and is current as of 2025-09-02. Clinical Trials information is current as of 2025-09-02. For the most up-to-date information regarding a particular trial, search [www.clinicaltrials.gov](http://www.clinicaltrials.gov) by NCT ID or search local clinical trials authority website by local identifier listed in 'Other identifiers.' Variants are reported according to HGVS nomenclature and classified following AMP/ASCO/CAP guidelines (Li et al. 2017). Based on the data sources selected, variants, therapies, and trials listed in this report are listed in order of potential clinical significance but not for predicted efficacy of the therapies.



## References

1. O'Leary et al. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. *Nucleic Acids Res.* 2016 Jan 4;44(D1):D733-45. PMID: 26553804
2. Hulpke et al. The MHC I loading complex: a multitasking machinery in adaptive immunity. *Trends Biochem Sci.* PMID: 23849087
3. Adams et al. The adaptable major histocompatibility complex (MHC) fold: structure and function of nonclassical and MHC class I-like molecules. *Annu Rev Immunol.* 2013;31:529-61. PMID: 23298204
4. Rossjohn et al. T cell antigen receptor recognition of antigen-presenting molecules. *Annu Rev Immunol.* 2015;33:169-200. PMID: 25493333
5. Parham. MHC class I molecules and KIRs in human history, health and survival. *Nat Rev Immunol.* 2005 Mar;5(3):201-14. PMID: 15719024
6. Sidney et al. HLA class I supertypes: a revised and updated classification. *BMC Immunol.* 2008 Jan 22;9:1. PMID: 18211710
7. Cornel et al. MHC Class I Downregulation in Cancer: Underlying Mechanisms and Potential Targets for Cancer Immunotherapy. *Cancers (Basel).* 2020 Jul 2;12(7). PMID: 32630675
8. Weinstein et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat. Genet.* 2013 Oct;45(10):1113-20. PMID: 24071849
9. Cerami et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov.* 2012 May;2(5):401-4. PMID: 22588877
10. Pylayeva-Gupta et al. RAS oncogenes: weaving a tumorigenic web. *Nat. Rev. Cancer.* 2011 Oct 13;11(11):761-74. PMID: 21993244
11. Karnoub et al. Ras oncogenes: split personalities. *Nat. Rev. Mol. Cell Biol.* 2008 Jul;9(7):517-31. PMID: 18568040
12. Scott et al. Therapeutic Approaches to RAS Mutation. *Cancer J.* 2016 May-Jun;22(3):165-74. doi: 10.1097/PPO.000000000000187. PMID: 27341593
13. Román et al. KRAS oncogene in non-small cell lung cancer: clinical perspectives on the treatment of an old target. *Mol Cancer.* 2018 Feb 19;17(1):33. doi: 10.1186/s12943-018-0789-x. PMID: 29455666
14. Dinu et al. Prognostic significance of KRAS gene mutations in colorectal cancer—preliminary study. *J Med Life.* 2014 Oct-Dec;7(4):581-7. PMID: 25713627
15. Allegra et al. Extended RAS Gene Mutation Testing in Metastatic Colorectal Carcinoma to Predict Response to Anti-Epidermal Growth Factor Receptor Monoclonal Antibody Therapy: American Society of Clinical Oncology Provisional Clinical Opinion Update 2015. *J. Clin. Oncol.* 2016 Jan 10;34(2):179-85. PMID: 26438111
16. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/214665Orig1s009correctedlbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/214665Orig1s009correctedlbl.pdf)
17. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/216340s005lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/216340s005lbl.pdf)
18. NCCN Guidelines® - NCCN-Pancreatic Adenocarcinoma [Version 2.2025]
19. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/219616s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/219616s000lbl.pdf)
20. <https://assets.cwp.roche.com/f/126832/x/5738a7538b/irp230202.pdf>
21. <https://ir.revmed.com/node/11881/pdf>
22. <https://www.prnewswire.com/news-releases/d3-bio-inc-announces-fda-breakthrough-therapy-designation-and-orphan-drug-designation-for-d3s-001-for-the-treatment-of-patients-with-kras-g12c-mutated-cancers-302540808.html>
23. <https://www.prnewswire.com/news-releases/lillys-olomorasib-receives-us-fdas-breakthrough-therapy-designation-for-the-treatment-of-certain-newly-diagnosed-metastatic-kras-g12c-mutant-lung-cancers-302545643.html>
24. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125514s178lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125514s178lbl.pdf)
25. <https://bridgebio.com/news/bridgebio-pharma-announces-first-lung-cancer-patient-dosed-in-phase-1-2-trial-and-us-fda-fast-track-designation-for-shp2-inhibitor-bbp-398-in-combination-with-amgens-lumakras-sotorasib/>
26. <https://investor.verastem.com/news-releases/news-release-details/verastem-oncology-granted-fast-track-designation-combination>
27. <https://www.businesswire.com/news/home/20250109170439/en/>
28. <https://ir.revmed.com/news-releases/news-release-details/revolution-medicines-announces-fda-breakthrough-therapy>
29. <https://investor.verastem.com/news-releases/news-release-details/verastem-oncology-granted-fast-track-designation-vs-7375>
30. <https://www.d3bio.com/press-releases/d3-bios-d3s-001-receives-u-s-fda-fast-track-designation-for-the-treatment-of-colorectal-cancer-with-kras-g12c-mutation>
31. [https://cardiffoncology.com/wp-content/uploads/2021/07/Cardiff\\_Oncology\\_Investor\\_Presentation\\_July\\_2021.pdf](https://cardiffoncology.com/wp-content/uploads/2021/07/Cardiff_Oncology_Investor_Presentation_July_2021.pdf)
32. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/125084s279lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/125084s279lbl.pdf)
33. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/125147s213lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/125147s213lbl.pdf)



## References (continued)

34. Slebos et al. K-ras oncogene activation as a prognostic marker in adenocarcinoma of the lung. *N. Engl. J. Med.* 1990 Aug 30;323(9):561-5. PMID: 2199829
35. Ahmed et al. The TGF- $\beta$ /Smad4 Signaling Pathway in Pancreatic Carcinogenesis and Its Clinical Significance. *J Clin Med.* 2017 Jan 5;6(1). PMID: 28067794
36. Zhao et al. The role of TGF- $\beta$ /SMAD4 signaling in cancer. *Int. J. Biol. Sci.* 2018;14(2):111-123. PMID: 29483830
37. Cicens et al. KRAS, TP53, CDKN2A, SMAD4, BRCA1, and BRCA2 Mutations in Pancreatic Cancer. *Cancers (Basel).* 2017 Apr 28;9(5). PMID: 28452926
38. Miyaki et al. Role of Smad4 (DPC4) inactivation in human cancer. *Biochem. Biophys. Res. Commun.* 2003 Jul 11;306(4):799-804. PMID: 12821112
39. Mehrvarz et al. Association of SMAD4 mutation with patient demographics, tumor characteristics, and clinical outcomes in colorectal cancer. *PLoS ONE.* 2017;12(3):e0173345. PMID: 28267766
40. Cancer Genome Atlas Research Network. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature.* 2014 Sep 11;513(7517):202-9. doi: 10.1038/nature13480. Epub 2014 Jul 23. PMID: 25079317
41. Yan et al. Reduced Expression of SMAD4 Is Associated with Poor Survival in Colon Cancer. *Clin. Cancer Res.* 2016 Jun 15;22(12):3037-47. PMID: 26861460
42. Voorneveld et al. A Meta-Analysis of SMAD4 Immunohistochemistry as a Prognostic Marker in Colorectal Cancer. *Transl Oncol.* 2015 Feb;8(1):18-24. PMID: 25749173
43. Shugang et al. Prognostic Value of SMAD4 in Pancreatic Cancer: A Meta-Analysis. *Transl Oncol.* 2016 Feb;9(1):1-7. PMID: 26947875
44. Boulay et al. SMAD4 is a predictive marker for 5-fluorouracil-based chemotherapy in patients with colorectal cancer. *Br. J. Cancer.* 2002 Sep 9;87(6):630-4. PMID: 12237773
45. Kozak et al. Smad4 inactivation predicts for worse prognosis and response to fluorouracil-based treatment in colorectal cancer. *J. Clin. Pathol.* 2015 May;68(5):341-5. PMID: 25681512
46. Ozawa et al. SMAD4 Loss Is Associated with Cetuximab Resistance and Induction of MAPK/JNK Activation in Head and Neck Cancer Cells. *Clin. Cancer Res.* 2017 Sep 1;23(17):5162-5175. PMID: 28522603
47. Philpott et al. The NF1 somatic mutational landscape in sporadic human cancers. 2017 Jun 21;11(1):13. doi: 10.1186/s40246-017-0109-3. PMID: 28637487
48. Scheffzek et al. The Ras-RasGAP complex: structural basis for GTPase activation and its loss in oncogenic Ras mutants. *Science.* 1997 Jul 18;277(5324):333-8. PMID: 9219684
49. Fioretos et al. Isochromosome 17q in blast crisis of chronic myeloid leukemia and in other hematologic malignancies is the result of clustered breakpoints in 17p11 and is not associated with coding TP53 mutations. *Blood.* 1999 Jul 1;94(1):225-32. PMID: 10381517
50. NCCN Guidelines® - NCCN-Myelodysplastic Syndromes [Version 2.2025]
51. Brems et al. Mechanisms in the pathogenesis of malignant tumours in neurofibromatosis type 1. *Lancet Oncol.* 2009 May;10(5):508-15. PMID: 19410195
52. NCCN Guidelines® - NCCN-Soft Tissue Sarcoma [Version 1.2025]
53. Nag et al. The MDM2-p53 pathway revisited. *J Biomed Res.* 2013 Jul;27(4):254-71. PMID: 23885265
54. Muller et al. Mutant p53 in cancer: new functions and therapeutic opportunities. *Cancer Cell.* 2014 Mar 17;25(3):304-17. PMID: 24651012
55. Olivier et al. TP53 mutations in human cancers: origins, consequences, and clinical use. *Cold Spring Harb Perspect Biol.* 2010 Jan;2(1):a001008. PMID: 20182602
56. Guha et al. Inherited TP53 Mutations and the Li-Fraumeni Syndrome. *Cold Spring Harb Perspect Med.* 2017 Apr 3;7(4). PMID: 28270529
57. Peter et al. Comprehensive genomic characterization of squamous cell lung cancers. *Nature.* 2012 Sep 27;489(7417):519-25. PMID: 22960745
58. Cancer Genome Atlas Network. Comprehensive genomic characterization of head and neck squamous cell carcinomas. *Nature.* 2015 Jan 29;517(7536):576-82. PMID: 25631445
59. Campbell et al. Distinct patterns of somatic genome alterations in lung adenocarcinomas and squamous cell carcinomas. *Nat. Genet.* 2016 Jun;48(6):607-16. PMID: 27158780
60. Cancer Genome Atlas Research Network. Integrated genomic characterization of oesophageal carcinoma. *Nature.* 2017 Jan 12;541(7636):169-175. doi: 10.1038/nature20805. Epub 2017 Jan 4. PMID: 28052061



## References (continued)

61. Olivier et al. The IARC TP53 database: new online mutation analysis and recommendations to users. *Hum. Mutat.* 2002 Jun;19(6):607-14. PMID: 12007217
62. Rivlin et al. Mutations in the p53 Tumor Suppressor Gene: Important Milestones at the Various Steps of Tumorigenesis. *Genes Cancer.* 2011 Apr;2(4):466-74. PMID: 21779514
63. Petitjean et al. TP53 mutations in human cancers: functional selection and impact on cancer prognosis and outcomes. *Oncogene.* 2007 Apr 2;26(15):2157-65. PMID: 17401424
64. Soussi et al. Recommendations for analyzing and reporting TP53 gene variants in the high-throughput sequencing era. *Hum. Mutat.* 2014 Jun;35(6):766-78. PMID: 24729566
65. <https://www.globenewswire.com/news-release/2020/10/13/2107498/0/en/PMV-Pharma-Granted-FDA-Fast-Track-Designation-of-PC14586-for-the-Treatment-of-Advanced-Cancer-Patients-that-have-Tumors-with-a-p53-Y220C-Mutation.html>
66. Parrales et al. Targeting Oncogenic Mutant p53 for Cancer Therapy. *Front Oncol.* 2015 Dec 21;5:288. doi: 10.3389/fonc.2015.00288. eCollection 2015. PMID: 26732534
67. Zhao et al. Molecularly targeted therapies for p53-mutant cancers. *Cell. Mol. Life Sci.* 2017 Nov;74(22):4171-4187. PMID: 28643165
68. Louis et al. The 2021 WHO Classification of Tumors of the Central Nervous System: a summary. *Neuro Oncol.* 2021 Aug 2;23(8):1231-1251. PMID: 34185076
69. Döhner et al. Diagnosis and management of AML in adults: 2022 recommendations from an international expert panel on behalf of the ELN. *Blood.* 2022 Sep 22;140(12):1345-1377. PMID: 35797463
70. NCCN Guidelines® - NCCN-Myeloproliferative Neoplasms [Version 2.2025]
71. NCCN Guidelines® - NCCN-Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma [Version 3.2025]
72. NCCN Guidelines® - NCCN-Acute Lymphoblastic Leukemia [Version 2.2025]
73. NCCN Guidelines® - NCCN-B-Cell Lymphomas [Version 3.2025]
74. Bernard et al. Implications of TP53 allelic state for genome stability, clinical presentation and outcomes in myelodysplastic syndromes. *Nat. Med.* 2020 Aug 3. PMID: 32747829
75. Volinia et al. Molecular cloning, cDNA sequence, and chromosomal localization of the human phosphatidylinositol 3-kinase p110 alpha (PIK3CA) gene. *Genomics.* 1994 Dec;24(3):472-7. PMID: 7713498
76. Whale et al. Functional characterization of a novel somatic oncogenic mutation of PIK3CB. *Signal Transduct Target Ther.* 2017;2:17063. PMID: 29279775
77. Osaki et al. PI3K-Akt pathway: its functions and alterations in human cancer. *Apoptosis.* 2004 Nov;9(6):667-76. PMID: 15505410
78. Cantley. The phosphoinositide 3-kinase pathway. *Science.* 2002 May 31;296(5573):1655-7. PMID: 12040186
79. Fruman et al. The PI3K Pathway in Human Disease. *Cell.* 2017 Aug 10;170(4):605-635. PMID: 28802037
80. Engelman et al. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat. Rev. Genet.* 2006 Aug;7(8):606-19. PMID: 16847462
81. Vanhaesebroeck et al. PI3K signalling: the path to discovery and understanding. *Nat. Rev. Mol. Cell Biol.* 2012 Feb 23;13(3):195-203. PMID: 22358332
82. Yuan et al. PI3K pathway alterations in cancer: variations on a theme. *Oncogene.* 2008 Sep 18;27(41):5497-510. PMID: 18794884
83. Liu et al. Targeting the phosphoinositide 3-kinase pathway in cancer. *Nat Rev Drug Discov.* 2009 Aug;8(8):627-44. PMID: 19644473
84. Hanahan et al. Hallmarks of cancer: the next generation. *Cell.* 2011 Mar 4;144(5):646-74. PMID: 21376230
85. Brito et al. PIK3CA Mutations in Diffuse Gliomas: An Update on Molecular Stratification, Prognosis, Recurrence, and Aggressiveness. *Clin Med Insights Oncol.* 2022;16:11795549211068804. PMID: 35023985
86. Huret et al. Atlas of genetics and cytogenetics in oncology and haematology in 2013. *Nucleic Acids Res.* 2013 Jan;41(Database issue):D920-4. PMID: 23161685
87. Miled et al. Mechanism of two classes of cancer mutations in the phosphoinositide 3-kinase catalytic subunit. *Science.* 2007 Jul 13;317(5835):239-42. PMID: 17626883
88. Burke et al. Synergy in activating class I PI3Ks. *Trends Biochem. Sci.* 2015 Feb;40(2):88-100. PMID: 25573003
89. Burke et al. Oncogenic mutations mimic and enhance dynamic events in the natural activation of phosphoinositide 3-kinase p110α (PIK3CA). *Proc. Natl. Acad. Sci. U.S.A.* 2012 Sep 18;109(38):15259-64. PMID: 22949682
90. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2024/212526s009lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2024/212526s009lbl.pdf)



## References (continued)

91. Mayer et al. A Phase Ib Study of Alpelisib (BYL719), a PI3K $\alpha$ -Specific Inhibitor, with Letrozole in ER+/HER2- Metastatic Breast Cancer. *Clin. Cancer Res.* 2017 Jan 1;23(1):26-34. PMID: 27126994
92. Mayer et al. A Phase II Randomized Study of Neoadjuvant Letrozole Plus Alpelisib for Hormone Receptor-Positive, Human Epidermal Growth Factor Receptor 2-Negative Breast Cancer (NEO-ORB). *Clin. Cancer Res.* 2019 Feb 5. PMID: 30723140
93. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/218197s002lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/218197s002lbl.pdf)
94. [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2025/219249s002lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2025/219249s002lbl.pdf)
95. Jung et al. Pilot study of sirolimus in patients with PIK3CA mutant/amplified refractory solid cancer. *Mol Clin Oncol.* 2017 Jul;7(1):27-31. PMID: 28685070
96. Janku et al. PIK3CA mutations in patients with advanced cancers treated with PI3K/AKT/mTOR axis inhibitors. *Mol. Cancer Ther.* 2011 Mar;10(3):558-65. PMID: 21216929
97. NCCN Guidelines® - NCCN-Colon Cancer [Version 4.2025]
98. NCCN Guidelines® - NCCN-Rectal Cancer [Version 3.2025]
99. Liao et al. Aspirin use, tumor PIK3CA mutation, and colorectal-cancer survival. *N Engl J Med.* 2012 Oct 25;367(17):1596-606. PMID: 23094721
100. Domingo et al. Evaluation of PIK3CA mutation as a predictor of benefit from nonsteroidal anti-inflammatory drug therapy in colorectal cancer. *J Clin Oncol.* 2013 Dec 1;31(34):4297-305. PMID: 24062397
101. Wang et al. Loss of Tumor Suppressor Gene Function in Human Cancer: An Overview. *Cell. Physiol. Biochem.* 2018;51(6):2647-2693. PMID: 30562755
102. Stamos et al. The  $\beta$ -catenin destruction complex. *Cold Spring Harb Perspect Biol.* 2013 Jan 1;5(1):a007898. PMID: 23169527
103. Minde et al. Messing up disorder: how do missense mutations in the tumor suppressor protein APC lead to cancer?. *Mol Cancer.* 2011 Aug 22;10:101. doi: 10.1186/1476-4598-10-101. PMID: 21859464
104. Aoki et al. Adenomatous polyposis coli (APC): a multi-functional tumor suppressor gene. *J. Cell. Sci.* 2007 Oct 1;120(Pt 19):3327-35. PMID: 17881494
105. Miyoshi et al. Somatic mutations of the APC gene in colorectal tumors: mutation cluster region in the APC gene. *Hum. Mol. Genet.* 1992 Jul;1(4):229-33. PMID: 1338904
106. Rowan et al. APC mutations in sporadic colorectal tumors: A mutational "hotspot" and interdependence of the "two hits". *Proc. Natl. Acad. Sci. U.S.A.* 2000 Mar 28;97(7):3352-7. PMID: 10737795
107. Laurent-Puig et al. APC gene: database of germline and somatic mutations in human tumors and cell lines. *Nucleic Acids Res.* 1998 Jan 1;26(1):269-70. PMID: 9399850